

Durham E-Theses

Development of nanoparticle catalysts and total internal reflection (TIR) Raman spectroscopy for improved understanding of heterogeneous catalysis

LAURA MARIA BINGHAM

How to cite:

BINGHAM, LAURA MARIA (2017) Development of nanoparticle catalysts and total internal reflection (TIR) Raman spectroscopy for improved understanding of heterogeneous catalysis. Doctoral thesis, Durham University.

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a <https://etheses.durham.ac.uk/id/eprint/12445/> is made to the metadata record in Durham E-Theses
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full Durham E-Theses policy](#) for further details.

Appendix: Chapter 1

The calculation of probing distances for infrared (IR) and Raman spectroscopy. The relationship between probing distance (d) and wavelength (λ) is given by Equation 1A.1. Values of $\theta=73^\circ$, and $n_1= 1.461$ (the refractive index for silicon) and $n_2= 1$ (the refractive index for air) were used (representative of conditions used within our system). This allowed for the plotting of probing depth as a function of wavelength (Figure 1A.1). A linear trend was seen. Typical Raman excitation wavelengths (such as 532 nm) gave lower probing depths (in the region of 40 nm), than for the higher wavelengths associated with IR spectroscopy.

$$d = \frac{\lambda_0}{4\pi} (n_1^2 \sin^2(\theta) - n_2^2)^{-\frac{1}{2}} \quad \text{Equation 1A.1}$$




Figure 1A.1. Graph showing the linear relation of probing depth against excitation wavelength for spectroscopic techniques.