

Durham E-Theses

*Some amine hydrofluorides and amines in
organofluorine chemistry*

Graham Sandford

How to cite:

Sandford, Graham (1991) Some amine hydrofluorides and amines in organofluorine chemistry. Doctoral thesis, Durham University.

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a <https://etheses.durham.ac.uk/id/eprint/6209/> is made to the metadata record in Durham E-Theses
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full Durham E-Theses policy](#) for further details.

UNIVERSITY OF DURHAM

**A THESIS
entitled**

**SOME AMINE HYDROFLUORIDES AND AMINES IN ORGANOFLUORINE
CHEMISTRY**

submitted by

GRAHAM SANDFORD B. Sc.
(Van Mildert College)

The copyright of this thesis rests with the author.
No quotation from it should be published without
his prior written consent and information derived
from it should be acknowledged.

A candidate for the degree of Doctor of Philosophy

1991



- 9 JUL 1992

To Mum, Dad and Aly

Acknowledgements

I would like to thank Professor R.D. Chambers for his help and advice throughout this period of research.

Thanks to my industrial supervisor, Dr. Stewart Korn (ICI Specialties), for his useful discussions and willingness to carry out computer searches, provide chemicals, set up biological tests, etc., at ICI. Also, thanks to ICI for providing the grant.

This thesis could not have been completed without the help of the technical staff in the department who are: Dr. Mike Jones and Vince McNeilly (Mass Spectrometry); Dr. Ray Matthews, Dr. Alan Kenwright and Mrs. Julia Say (NMR); Lenny Lauchlan (Chemicals and Chromatography); Mrs. Mollie Cocks (Elemental Analysis); Ray Hart and Gordon Haswell (Glass Blowing), and thanks to Tom Holmes for endless help with practical chemistry and tasteless jokes.

Finally, thanks to the lads in the lab for making this period of study so enjoyable and to the New Inn for never throwing me out.

MEMORANDUM

The work described in this thesis was carried out in the University of Durham between October 1988 and September 1991. This thesis is the work of the author, except where acknowledged by reference, and has not been submitted for any other degree.

The work has been presented, in part, by the author at:

13th. International Symposium on Fluorine Chemistry, Ruhr Universitat, Bochum, Germany, September 1991.

NOMENCLATURE

Throughout this thesis an "F" in the centre of a ring is used to denote that all bonds are to fluorines.

CONTENTS**Chapter One****Sources of Fluoride Ion**

1.1	General Introduction	1
1.2	Formation of C-F Bonds using Fluoride Ion	2
1.2.1	Nucleophilic Substitution at Saturated Carbon	2
1.2.2	Nucleophilic Substitution Involving Unsaturated Carbon	2
1.2.3	Nucleophilic Substitution at Aromatic Carbon	3
1.3	Sources of Fluoride Ion	3
1.3.1	Alkali Metal Fluorides	3
1.3.1.1	Crown Ether Activation	4
1.3.1.2	Phase Transfer Catalysis	5
1.3.1.3	Additives to Alkali Metals	5
1.3.1.4	Spray-dried and Freeze-dried KF	6
1.3.1.5	KF on Support Reagents	6
1.3.1.6	Ion Exchange Resins	7
1.3.2	Alternatives to Alkali Metal Fluorides	7
1.3.2.1	Tetraalkylammonium and phosphonium Salts	8
1.3.2.2	Phosphoranes	9
1.3.2.3	Amine Hydrofluorides	9
1.3.2.4	DAST and TAS-F	9
1.3.2.5	Tris(morpholino)selenonium fluoride	10
1.4	Hydrogen Fluoride as a Fluorinating Reagent	11
1.4.1	Classical Processes	11
1.4.2	HF used in Conjunction with Lewis Bases	11
1.5	Pyridine Poly(Hydrogen Fluoride) - Olah's Reagent	13
1.5.1	Structure of Pyridine Poly(Hydrogen Fluoride)	13
1.5.2	Reactions of Pyridine Poly(Hydrogen Fluoride)	15
1.6	Amine Hydrofluoride Salts	18
1.6.1	Introduction	18
1.6.2	Spectroscopy of Amine/HF Systems	18
1.6.2.1	Nuclear Magnetic Resonance Spectroscopy	18
1.6.2.1.1	R ₃ N.nHF Systems	20
1.6.2.2	Infra Red Spectroscopy	22
1.6.2.2.1	Hydrogen Fluoride	22
1.6.2.2.2	Hydrogen Difluoride	22
1.6.2.2.3	Trialkylamine Hydrogen Fluoride Systems	23

1.6.3	Reactions of Amine Hydrofluorides as Fluorinating Agents	24
1.6.3.1	Halogen Exchange Reactions (Halex)	24
1.6.3.2	Halofluorination	24
1.6.3.3	Sulphenylation Fluorination	26
1.6.3.4	Phenylselenofluorination	26
1.6.3.5	Ring Opening Reactions	26
1.6.3.5.1	Aziridines	26
1.6.3.5.2	Aziridinium Ion	27
1.6.3.5.3	Epoxides	27
1.6.3.6	Oxidative Fluorination	28
1.6.3.7	Other Systems	29
1.7	Summary	

Chapter Two

1,8-Bis(dimethylamino)naphthalene Hydrofluoride (PS/HF) as a Potential Source of Fluoride Ion

2.1	Introduction	30
2.2	1,8-Bis(dimethylamino)naphthalene (Proton Sponge)	31
2.2.1	Preparation	31
2.2.2	Structure	31
2.2.3	Structure of 1,8-Bis(dimethylamino)naphthalene Salts	32
2.2.4	Basicity	33
2.2.5	Spectroscopy	34
2.2.5.1	UV	34
2.2.5.2	Infra Red	34
2.2.5.3	NMR	35
2.2.6	Reactions	35
2.3	The Use of the Hydrofluoride Complex of Proton Sponge as a Potential Source of Soluble Fluoride Ion	37
2.3.1	Preparation of PS/HF	37
2.3.2	Spectroscopical Studies of PS/HF	37
2.3.2.1	NMR	37
2.3.2.2	Infra Red	39
2.3.2.3	Mass Spectroscopy	40
2.4	Reactions of PS/HF	41
2.4.1	PS/HF as a Catalyst in C-C Bond Forming Reactions	41
2.4.1.1	Oligomerisation Reactions	41
2.4.1.2	Perfluoroalkylation Reactions	43

2.4.2	PS/HF in C-F Bond Forming Reactions	44
2.4.2.1	At Unsaturated Carbon	44
2.4.2.2	At Saturated Carbon	45
2.5	Summary	46

Chapter Three

Organic Base Hydrogen Fluoride Complexes as Sources of Soluble Fluoride Ion

3.1	Introduction	48
3.2	Trialkylamine Hydrogen Fluoride Complexes	48
3.2.1	NMR Spectroscopy	48
3.2.2	Low Temperature NMR Study of Bu ₃ N.HF	49
3.2.3	Reactions	51
3.3	Tetramethylpiperidinium Hydrofluoride Complexes	53
3.3.1	Preparation of Penta-alkyl piperidine Bases and their Hydrofluoride Complexes	53
3.3.2	NMR Spectroscopy	54
3.3.3	Reactions	55
3.4	Tetramethylguanidine Hydrofluoride Complex	56
3.5	Conclusions	56

Chapter Four

Annelation Reactions Between 1,8-Bis(dimethylamino)Naphthalene (Proton Sponge) and Fluorinated Alkenes

4.1	Introduction	58
4.2	Electrophilic Substitution Reactions of 1,8-Bis(dimethylamino)-Naphthalene (PS)	58
4.2.1	Nitration	58
4.2.2	Bromination	59
4.2.3	Alkylsulphination	59
4.2.4	Formylation	59
4.2.5	Reaction with Dinitrobenzofurazan	60
4.3	Reaction of 1,8-Bis(dimethylamino)naphthalene With Fluorinated Alkenes (3) and (43)	61
4.3.1	Structure Elucidation	62
4.3.1.1	NMR Spectra of (4) and (44)	62

4.3.2	Molecular Modelling of Annelation Product (4)	67
4.4	Mechanism of the Annelation Reaction	69
4.4.1	Orientation of Initial Nucleophilic Attack	69
4.5	Reaction of 1,8-Bis(dimethylamino)naphthalene With Perfluoro-bicyclopentylidene	72
4.5.1	At Low Dilution	72
4.5.2	At High Dilution	73
4.5.2.1	Green Crystals (60)	74
4.5.2.2	Purple Crystals (61)	77
4.5.2.3	Mechanism of Hydrolysis to Form (60) and (61)	80
4.6	Reactions of Perfluorobicyclobutylidene With Tertiary Aromatic Amines	81
4.6.1	With N,N-Dimethylaniline	81
4.6.2	With N-Methylindole	82
4.6.3	With Proton Sponge	83
4.7	Summary	84

Chapter Five

Reactions Between Tertiary Aromatic Amines Acting as Soft Carbon Nucleophiles With Fluorinated Heterocycles

5.1	Introduction	85
5.2	Nucleophilic Substitution Reactions of Trifluoro-s-triazine	85
5.3	Reactions Between Trichloro-s-triazine (Cyanuric Chloride) and Tertiary, Aromatic Amines Acting as Carbon Nucleophiles	86
5.3.1	With N,N-Dialkylanilines and Toluidines	86
5.3.1.1	Mechanism	89
5.3.2	With 5-Membered Heterocyclic Rings	89
5.4	Reactions of Fluorinated Triazines With Pyrroles	94
5.4.1	Spectroscopy of Pyrrolyl-s-Triazines	95
5.4.1.1	¹ H NMR	95
5.4.1.2	¹⁹ F NMR	98
5.4.1.3	¹³ C NMR	98
5.5	Reactions of Fluorinated Triazines With N-Methylindole	104
5.5.1	Spectroscopy of Indolyl-s-Triazines	105
5.5.1.1	¹ H NMR	105
5.5.1.2	¹⁹ F NMR	105
5.5.1.3	¹³ C NMR	107
5.6	Reaction of Tetrafluoropyrimidine With N-Methylindole	108
5.7	Reactions of Fluorinated s-Triazines With Anilines	109

5.7.1	Reaction of Trifluoro-s-triazine with N,N-Dimethylaniline	109
5.7.2	Reactions of Perfluoroisopropyl-s-triazine with N,N-Dimethyl- and Diethylaniline	109
5.7.3	Reactions of Perfluorodi-isopropyl-s-triazine with N,N-Dimethyl- and Diethylaniline	111
5.7.4	Collected NMR data for Phenyl-s-triazines	112
5.7.5	Potential Use of Compounds (77)-(82A) as Non-Linear Optic Materials	112
5.8	Reaction of Trifluoro-s-Triazine With PS	114
5.9	Reaction of Trifluoro-s-Triazine With N-Ethylaniline	115
5.10	Reaction of Trifluoro-s-Triazine With N,N-Dimethylamino-2-Methoxybenzene	116
5.11	Reaction of Trifluoro-s-Triazine With N,N,N',N'-Tetramethyl-Diamino-benzene	116
5.12	Reaction of Perfluorodiisopropyl-s-Triazine With 2-(N,N-Dimethyl-amino)-pyridine	117
5.13	Summary	118

EXPERIMENTAL SECTION

<u>Instrumentation</u>	120
-------------------------------	-----

Chapter Six

Experimental to Chapter Two

6.1	Preparation of HF/ether solution	121
6.2	Preparation of PS/HF Complex	121
6.3	Reactions of PS/HF as a Catalyst in C-C Bond Forming Reactions	121
6.3.1	Attempted Oligomerisation of Tetrafluoroethylene	121
6.3.2	Dimerisation of Hexafluoropropene	122
6.3.3	Attempted Oligomerisation of Perfluorocyclobutene	122
6.3.4	Attempted Formation of a Stable Perfluorinated Carbanion	122
6.3.5	Perfluoroalkylation of Pentafluoropyridine	123
6.3.6	Perfluoroalkylation of Tetrafluoropyrimidine	123
6.3.7	Perfluoroalkylation of Trifluoro-s-triazine	124
6.4	Reactions of PS/HF in C-F Bond Forming Reactions	124
6.4.1	With Benzoyl Chloride	124
6.4.2	With Hexafluoroacetone	124
6.4.2.1	Preparation of Heptafluoroisopropyl benzoate	125

6.4.2.2	Preparation of Heptafluoroisopropyl benzyl ether	125
6.4.3	With 2,4-Dinitrochlorobenzene	125
6.4.4	With Benzyl Bromide	125
6.4.5	With Octyl Iodide	126
6.4.6	With 1,2-Epoxybutane	126
6.4.7	Preparation of Triflate (21)	126
6.4.8	Attempted Reaction between PS/HF and Triflate (21)	126
6.4.9	Preparation of Tosylate (22)	127
6.4.10	Attempted Reaction of PS/HF with Tosylate (22)	127

Chapter Seven

Experimental to Chapter Three

7.1	Preparation and Reactions of Trialkylamine Hydrogen Fluoride Complexes	128
7.1.1	Preparation of $R_3N.HF$	128
7.1.2	Methodology for Standard Reactions	128
7.1.2.1	Reaction of Base Hydrofluoride Complexes with Benzoyl Chloride	129
7.1.2.2	Reaction of Base Hydrofluoride Complexes with Benzyl Bromide	129
7.1.2.3	Reaction of Base Hydrofluoride Complexes with 2,4-Dinitrochlorobenzene	129
7.1.3	Reactions of Trialkylamine.HF Complexes	130
7.1.3.1	With Benzoyl Chloride	130
7.1.3.2	With Benzyl Bromide	130
7.1.3.3	With 2,4-Dinitrochlorobenzene	131
7.2	Preparation and Reactions of Polyalkylpiperidine HF Complexes	131
7.2.1	Preparation of Pentasubstituted Piperidine Bases	131
7.2.1.1	N-ethyl-2,2,6,6,-Tetramethylpiperidine	131
7.2.1.2	N-allyl-2,2,6,6-Tetramethylpiperidine	132
7.2.1.3	N-benzyl-2,2,6,6-Tetramethylpiperidine	132
7.2.2	Preparation of the Base.HF Complexes (35)-(39)	132
7.2.3	Reactions	133
7.2.3.1	With Benzoyl Chloride	133
7.2.3.2	With Benzyl Bromide	134
7.2.3.3	With 2,4-Dinitrochlorobenzene	134
7.3	Preparation and Reactions of Tetramethylguanidine HF Complex	134
7.3.1	Attempted Preparation of Pentamethylguanidine	135

Chapter Eight**Experimental to Chapter Four**

8.1	Reaction between PS/HF, Perfluorocyclobutene and Hexafluoropropene	136
8.2	Reaction Between PS and Co-dimer (3)	136
8.3	Preparation of Co-dimer (43)	136
8.4	Reaction between PS and Co-dimer (43)	137
8.5	Reaction between PS and Perfluorobicyclopentylidene	137
	8.5.1 At Low Dilution	137
	8.5.2 At High Dilution	137
8.6	Preparation of Perfluorobicyclobutylidene	138
8.7	Reaction between N,N-Dimethylaniline and Perfluorobicyclobutylidene	138
8.8	Reaction between N-Methylindole and Perfluorobicyclobutylidene	138
8.9	Reaction between PS and Perfluorobicyclobutylidene	139

Chapter Nine**Experimental to Chapter Five**

9.1	Preparation of Trifluoro-s-Triazine	140
9.2	Preparation of Perfluoroisopropyl-s-triazine (14) and perfluorodi-isopropyl-s-triazine (15)	140
9.3	Reactions of Pyrroles With Fluorinated Triazines	140
	9.3.1 Pyrrole with trifluoro-s-triazine	140
	9.3.2 Pyrrole with perfluoroisopropyl-s-triazine	141
	9.3.3 Pyrrole with perfluorodi-isopropyl-s-triazine	141
	9.3.4 N-methylpyrrole with trifluoro-s-triazine	141
	9.3.5 N-methylpyrrole with perfluoroisopropyl-s-triazine	141
	9.3.6 N-methylpyrrole with perfluorodi-isopropyl-s-triazine	142
9.4	Reactions of N-Methylindole With Fluorinated triazines	142
	9.4.1 With trifluoro-s-triazine	142
	9.4.2 With perfluoroisopropyl-s-triazine	142
	9.4.3 With perfluorodi-isopropyl-s-triazine	142
9.5	Reaction of N-Methylindole With Tetrafluoropyrimidine	143
9.6	Reactions of Anilines With Fluorinated Triazines	143
	9.6.1 N,N-Dimethylaniline with Trifluoro-s-triazine	143
	9.6.2 N,N-Dimethylaniline with perfluoroisopropyl-s-triazine	143
	9.6.3 N,N-Diethylaniline with perfluoroisopropyl-s-triazine	144

9.6.4	N,N-Dimethylaniline with perfluorodi-isopropyl-s-triazine	144
9.6.5	N,N-Diethylaniline with perfluorodi-isopropyl-s-triazine	144
9.6.6	Proton Sponge with trifluoro-s-triazine	145
9.6.7	N-Ethylaniline with trifluoro-s-triazine	145
9.6.8	N-Ethylaniline with perfluoroisopropyl-s-triazine	145
9.6.9	Preparation of N,N-Dimethylamino-2-methoxybenzene	146
9.6.10	Reaction of N,N-Dimethylamino-2-methoxybenzene with trifluoro-s-triazine	146
9.6.11	Reaction of N,N,N',N'-Tetramethyl-1,4-diaminobenzene with trifluoro-s-triazine	146
9.6.12	Reaction of 2-N,N-Dimethylamino-pyridine with Perfluoro-di-isopropyl-s-triazine	147



Appendices

<u>Appendix One</u>	<u>N.M.R. Spectra</u>	149
<u>Appendix Two</u>	<u>Infra Red Spectra</u>	174
<u>Appendix Three</u>	<u>Mass Spectra</u>	185
<u>Appendix Four</u>	<u>Colloquia and Conferences Attended</u>	228
<u>References</u>		236


CHAPTER ONE**SOURCES OF FLUORIDE ION****1.1 General Introduction**

Since the Second World War research in organofluorine chemistry has increased tremendously, motivated by the discovery of the unusual properties acquired by some molecules on the introduction of fluorine or fluorine containing substituents. Organofluorine compounds have been used in an ever widening range of applications from refrigerants to artificial blood (for some examples see Table 1). As fluorine containing compounds are not generally found in nature the area of organofluorine chemistry is entirely synthetic, and consequently a whole new range of reactions and reaction mechanisms may be studied, adding to our understanding of organic chemistry.

Table 1. Some Applications of Fluorine Containing Compounds

<u>Product</u>	<u>Application</u>
CF ₂ Cl ₂	Refrigerant
CF ₃ CHBrCl	Anaesthetic
- (CF ₂) _n -	Polymer with high thermal and chemical stability. Non-stick properties
Perfluorodecalin	Artificial blood
Fluoro Steroids	Anti-inflammatory Agent
 5-F-Uracil	Anti-Cancer Drug
 Trifluralin	Plant Protection (Weed control in Maize)





Surfactants

Dyes containing CF₃ groups

Good light fastness

Differences in chemical and physical properties between hydrocarbon and fluorocarbon systems are mainly due to a) electronegativity differences, 2) unshared electron pairs on fluorine, 3) the more easy displacement of fluorine as fluoride ion, and 4) the greater bond strength of C-F.

As the number of fluorine containing compounds grows, the variety of fluorinating reagents and methods of synthesis also increase and intense research continues in the development of new classes of fluorinating reagents. This thesis is concerned with the development of new types of fluoride ion reagent and so a brief description of the chemistry and the sources of fluoride ion follows.

1.2 Formation of C-F Bonds using Fluoride Ion

The introduction of fluorine into an organic molecule via the displacement of a leaving group by fluoride ion in a nucleophilic substitution reaction has proved to be extremely efficient in many cases¹⁻⁴. Nucleophilic substitution reactions at saturated, unsaturated and aromatic carbon are possible and general mechanisms are outlined below¹.


1.2.1 Nucleophilic Substitution at Saturated Carbon




1.2.2 Nucleophilic Substitution Involving Unsaturated Carbon


Three modes of nucleophilic substitution involving fluoride ion can occur in unsaturated systems:-

(i) Addition Elimination



(ii) S_N2(iii) Substitution with Rearrangement - S_N2!

1.2.3 Nucleophilic Substitution at Aromatic Carbon



Other aspects of fluoride ion chemistry, such as catalysing the oligomerisation of perfluoroalkenes, are discussed at the relevant sections of the following text.

Hence, fluoride ion is a valuable synthetic reagent and a discussion of modern sources of fluoride ion follows.

1.3 Sources of Fluoride Ion

1.3.1 Alkali Metal Fluorides

A general order of reactivity of the alkali metal fluorides is: CsF > KF > NaF, LiF, which may be attributed to the fact that caesium fluoride has the lowest lattice energy. The easy availability of the alkali metal fluorides, KF and CsF, means that these reagents continue to be widely used. However, their low reactivity, low solubility, hygroscopic nature and the harsh reaction conditions required has limited their use in more complex systems. Hence, new sources of fluoride ion, which may introduce fluorine into molecules in greater yield and greater selectivity, continue to be developed. A brief description of the main classes of modern fluoride ion reagents which have been investigated in an attempt to alleviate these problems will be discussed here for completeness, but are reviewed in more detail elsewhere⁵.



Examples of the use of these modern reagents in synthesis, from the recent literature (1988-1990 where possible), are presented.

The reactivity of the alkali metal fluorides may be improved by either increasing their solubility or by increasing the surface area of the reagent. The following methods have been used to improve the reactivity of the alkali metal fluorides.


1.3.1.1 Crown Ether Activation

The nucleophilic substitution process is enhanced by the use of a chelating crown ether which selectively complexes with the cation leaving the unsolvated fluoride ion ("naked fluoride") strongly nucleophilic⁶. The addition of a crown ether to KF increases its solubility, thus increasing the concentration of fluoride ion in solution⁷. 18-Crown-6 was found to be the most active catalyst⁸ due to its excellent complexing ability with K^+ and Cs^+ .


Selected examples of the use of alkali metal fluorides in conjunction with crown ethers are as follows^{6,8}:-



KF has been used with a combination of 18-Crown-6 and tetraphenylphosphonium bromide in halax reactions⁹.




Primary alcohols are converted to the fluorides by a $CsF/18\text{-Crown-6}/\text{methanesulphonyl fluoride}$ system¹⁰.




Fluorodestannylation reactions have been performed¹¹.

1.3.1.2 Phase Transfer Catalysis

The concentration of fluoride ion in the organic phase of a reaction may be increased by using KF in conjunction with a phase transfer catalyst (PTC). Reactions of the following type were studied¹²:-





The role of the PTC, usually a tetraalkylammonium halide, may be considered thus:-




Hence, fluoride ion is transferred from the aqueous/solid phase into the organic phase enhancing the reactivity of KF.

1.3.1.3 Additives to KF

The addition of alkyipyridinium salts¹³ and tetraalkylammonium chlorides¹⁴ to KF has been shown to be effective in halogen exchange reactions:-




The addition of tetraphenylphosphonium bromide improves the reactivity of KF¹⁵. Fluorodenitration reactions have been performed using this system¹⁶.



Tetraphenyl phosphonium bromide has been used as an additive supported on a cross-linked styrene/*p*-chloro-methyl styrene copolymer in fluorinations of aromatic chlorides¹⁷.

1.3.1.4 Spray-Dried and Freeze-Dried KF

The surface area of "normal" KF (0.1m²/g) is much less than that of "spray-dried" KF (1.3m²/g) and consequently spray-dried KF shows greater reactivity than normal KF¹⁸.




Similarly, "freeze-dried" KF has been investigated, but was originally found to be ineffective for the fluorination of some activated chlorinated compounds¹⁹. However, renewed interest in this reagent showed that fluorination was possible²⁰:-




1.3.1.5 KF on Support Reagents

An alternative to the use of phase transfer catalysis to improve the reactivity of KF is the use of supported reagents which show greater reactivity due to the increased reagent surface area. The reactivity of KF-alumina²¹ was seen to be very low due to surface OH-F hydrogen bonding. However, non-surface hydroxylated support materials, such as KF-CaF₂, have been successfully applied as sources of fluoride ion^{22,23}, as shown below²⁴.




Inorganic support materials may be surface dehydroxylated prior to use to avoid OH-F interactions²⁵. For instance, the surface of alumina is modified by organosilylation of the hydroxyl groups using hexamethyldisilazane. The resultant KF-modified support was used to fluorinate benzyl bromide in 61% yield.

Cross-linked copolymers of divinylbenzene and styrene were used as a support material for KF to good effect. The copolymers contain no surface OH groups and can be synthesized to have specific surface properties²⁶:-.




1.3.1.6 Ion Exchange Resins

Anion exchange resins, such as Amberlite IRA 900 of the form below, have been used as carriers of fluoride ion²⁷.



The resin may be considered as both a phase transfer catalyst and a support material, acting in the following way:-



1.3.2 Alternatives to Alkali Metal Fluorides

Compounds based on elements of Group V (nitrogen, phosphorous) and Group VI (sulphur, selenium) have been studied as sources of fluoride ion and a brief outline of

their use is given here. However, the very hygroscopic nature and high cost of some of these reagents give large grounds for improvement.

1.3.2.1 Tetraalkyl Ammonium and Phosphonium Salts

Tetraalkylammonium fluorides such as tetrabutylammonium fluoride (TBAF) have received considerable attention over the last few years. "Anhydrous" TBAF is prepared by heating TBAF.3H₂O at 40-45°C under high vacuum for several hours. It must be used immediately due to its hygroscopic nature and has been used in a full range of fluoride ion reactions²⁸, as shown in Table 2.

Table 2 Reactions using TBAF as source of Fluoride ion

<u>Substrate</u>	<u>Time/hr</u>	<u>Product</u>	<u>Yield/%</u>
CH ₂ =CH-CH ₂ Br	0.1	CH ₂ =CH-CH ₂ F	85
Ph-CH ₂ Br	8	Ph-CH ₂ F	100
CH ₃ (CH ₂) ₇ Br	<1	CH ₃ (CH ₂) ₇ F	48
		CH ₂ =CH(CH ₂) ₅ CH ₃	12
		CH ₃ (CH ₂) ₇ OH	40
CH ₃ (CH ₂) ₇ OTos	<1	CH ₃ (CH ₂) ₇ F	98
PhCOCl	<1	PhCOF	81


Also, fluorodenitration reactions of aromatic nitro groups have been performed²⁹ and fluorodeoxy sugars have been prepared from pyranosides³⁰.



Recently, a method for the preparation of anhydrous and HF₂⁻ free tetramethylammonium fluoride has been described³¹. Me₄NF.H₂O is first heated under vacuum to remove most of the water and then recrystallised in dry isopropanol as the alcoholate. The solvated alcohol is removed under vacuum at 80°C.

Me₄NF was found to abstract a proton from acetonitrile³² resulting in the slow formation of HF₂⁻ and the dimerisation of the acetonitrile:-





Also, chlorinated solvents such as chloroform undergo halox reactions.




This shows the remarkable reactivity of the fluoride ion when present as a soluble salt free from HF, HF_2^- and water. Their use in organic synthesis is yet to be investigated but their reactions with solvents may be a problem.

Tetraphenylphosphonium hydrogendifluoride has been used successfully in a variety of halogen exchange reactions³³. Similarly, tetraamidophosphonium hydrogendifluoride reacts with decyl tosylate and p-chloronitrobenzene to give the corresponding fluorides in 100% and 93% yield respectively³⁴.

1.3.2.2 Phosphoranes

The salts methyltri-n-butylfluorophosphorane ($(\text{n-C}_4\text{H}_9)_3\text{PFCH}_3$) and phenyl tetrafluorophosphorane ($\text{C}_6\text{H}_5\text{PF}_4$) have been used in tosyl displacement reactions in the preparation of 2-fluoro alkyl compounds³⁵.




1.3.2.3 Amine Hydrofluoride Salts


Pyridine/HF mixtures and Triethylamine Trishydrofluoride have been used in a variety of fluorination reactions and will be discussed in detail in sections 1.5 and 1.6.

1.3.2.4 Diethylaminosulphur Trifluoride (DAST) and Tris(dimethylamino)-sulphonium Difluorotrimethylsilicate (TAS-F)

DAST is prepared by reacting diethylaminotrimethylsilane with sulphur tetrafluoride³⁶:-



The main synthetic use for DAST is converting alcohols to fluorides³⁷ and carbonyl groups to geminal difluorides.




However, DAST is thermally unstable and recently the more stable morpholino sulphur trifluoride has been recommended for fluorinations of alcohols³⁸.

The more reactive fluoride ion source, TAS-F, is prepared by the reaction of dimethylaminosulphurdifluoride with dimethylaminotrimethylsilane:-





TAS-F readily converts even unreactive halides to fluorides under very mild conditions and has been used to form stable perfluorinated carbanion salts³⁹, in a fluoride promoted C-C bond cleavage reaction.



A comprehensive review on the use and reactions of DAST, TAS-F and related reagents has been published recently³⁶.

1.3.2.5 Tris(morpholino)selenonium Fluoride


The reaction of RSiMe_3 ($\text{R}=\text{morpholino}$) with selenium tetrafluoride gives tris(morpholino) selenonium fluoride, which has been used in a few fluorination reactions⁴⁰.



1.4 Hydrogen Fluoride as a Fluorinating Agent

1.4.1 Classical Processes

Anhydrous Hydrogen fluoride (HF) is one of the least expensive fluorinating agents and has been used in a variety of halox and electrophilic addition reactions²:-




However, most reactions using HF are performed under pressure or in the vapour phase due to its low boiling point (19.6°C). The corrosive and harmful nature of HF requires rigorous safety precautions and specialist equipment.


1.4.2 HF used in Conjunction with Lewis Base Co-solvents

To overcome the high volatility of HF several authors have studied complexes of HF with various Lewis bases as co-solvents.

Stable solutions of HF in amides⁴¹ (formation of amide/HF complexes), carbamic acids and esters⁴² (hydrofluorination of epoxide rings in steroids), trialkylphosphines⁴³ (formation of triethyl and triphenyl phosphine/HF complexes) and tetrahydrofuran⁴⁴ (ring opening hydrofluorination of epoxides in steroids) were investigated. There has been renewed interest in THF/HF solutions^{45,46}, the solution formed being a more effective source of fluoride ion than HF and also more acidic than pyridine/HF. Examples of the use of HF/THF solutions are below⁴⁴⁻⁴⁶:-



- -via an umpolung strategy involving the phenoxonium ion as an intermediate



It was not until the introduction of amine/HF complexes, in particular Pyridine/HF, that a general fluorinating agent using HF coupled to a base became widely accepted. These reagents are discussed in detail in the next section.

1.5 Pyridine Poly(Hydrogen Fluoride) - Olah's Reagent

The use of Lewis Base/HF solutions as fluorinating reagents became widely accepted upon the introduction of Pyridine/HF solutions by Olah⁴⁷, although pyridine/HF solutions had been used previously in epoxide ring opening hydrofluorination reactions of steroids by Bergstrom⁴⁸.

Pyridine forms stable solutions with anhydrous HF. The solution generally used consists of about 9 equivalents of HF to 1 of pyridine (70% w/w HF, 30% w/w pyridine) and is stable to 55°C.


Pyridinium Fluoride can be prepared by the reaction of formyl fluoride with pyridine through the decarbonylation of the intermediate N-formylpyridinium fluoride⁴⁹.



However, no spectral data appears in the publication.

1.5.1 Structure of Pyridinium Poly(Hydrogen Fluoride)

The ¹H n.m.r. spectrum for pyridinium poly(hydrogen fluoride) shows a typical pattern for pyridinium ring protons but the ¹⁹F n.m.r. spectrum at -60°C shows a quintet ($J_{\text{HF}}=120$ Hz) at 188.1ppm, indicating the presence of a polyhydrogen fluoride species in which each fluorine atom is surrounded by four protons:-



It is unclear whether any exchange occurs between the pyridinium cation and the HF matrix and we would also expect to see n.m.r. peaks due to species such as HF_2^- . The presence of pyridinium hydrogen difluoride in pyridinium poly(hydrogen fluoride) solutions was shown in low temperature X-ray crystallographic work by Mootz⁵⁰ (see below). Perhaps, as $\text{PyH}^+\text{HF}_2^-$ and PyH^+F^- have melting points of -1°C and -31°C respectively, from differential thermal analysis, these species were not seen in the n.m.r. spectrum as they are solid at -60°C (No n.m.r. solvent is indicated in the publication). Hence, at -60°C the only species present in pyridinium poly(hydrogen


fluoride)solutions are Pyridine.5HF and higher homologues which give rise to the observed quintet.

The pyridine/HF mixture was studied by differential thermal analysis and X-ray crystallography by Mootz⁵⁰, who identified 8 intermediary compounds, $C_5H_5N.nHF$ ($n=1-8$), with melting points between -1 and $-124^\circ C$. X-ray structures of complexes $n=1-4$ were obtained. See Fig. 1 for structures of $n=1,2$.

The following points can be noted from the structures reported:-

1) In complex $C_6H_5N.HF$ ($n=1$) the hydrogen atom in the NHF hydrogen bond is found much closer to the fluorine atom than to the nitrogen atom suggesting that the hydrogen bond is of the type F-H...N rather than F..H-N, i.e. the base is not protonated by the

Fig. 1 X-ray Crystal Structures of Pyridine.nHF Complexes



The structures of $C_5H_5N.nHF$ with $n=1, 2$, and 3 . One formula unit each with interatomic distances (pm) and angles. The N-F and F-F distances in the hydrogen bonds are underlined.

acid. This is the first hydrogen bond of this type reported and the shortest between nitrogen and fluorine.


2) Complex $C_6H_5N.2HF$ ($n=2$), can be reformulated as $C_5H_5NH^+HF_2^-$, pyridinium hydrogendifluoride, with the H atom in an off-centre position.

3) Ionic formulae are also true of the higher pyridine/HF complexes, e.g. $C_5H_5NH^+H_2F_3^-$ ($n=3$), and $C_5H_5NH^+H_3F_4^-$ ($n=4$).


1.5.2 Reactions of Pyridinium poly(hydrogen fluoride)

Pyridinium poly(hydrogen fluoride) was found to be an effective fluorinating agent for various additions to alkenes and alkynes, for deaminative and dediazonative halogenation reactions, for fluorine substitution of hydroxyl groups and halogen exchange reactions^{47, 51}, as summarised in Table 3.

Table 3. Reactions Performed Using Pyridinium Poly(Hydrogen Fluoride)





Dediazonative fluorinations of *p*-aminophenols have been performed recently^{52,53}.




More recently, pyridinium poly(hydrogen fluoride) has been used in ring opening hydrofluorination reactions of aziridines^{54,55}, azabicycloalkane⁵⁶ and cyclopropane⁵⁷ ring systems, as summarised in Table 4.




Table 4. Ring Opening Hydrofluorination Reactions using Pyridine Poly(Hydrogen Fluoride)

Aziridines^{54, 55}



Ring opening hydrofluorination of 3-aza-1,8,8-trimethyl-tricyclo [5.2.1.0^{2,4}] octane leads to a variety of products⁵⁸.



Azabicycloalkanes⁵⁶Cyclopropanes⁵⁷

Poly-4-vinyl pyridinium poly(hydrogen fluoride), a solid pyridine poly(hydrogen fluoride) has been used in hydrofluorinations and bromofluorinations of alkenes⁵⁹.

Most recent publications dealing with Amine/HF systems utilise Et₃N.3HF as the fluorinating agents and it is to these systems that we now turn.

1.6 Amine Hydrofluoride Salts

1.6.1 Introduction

Amine hydrofluoride salts have been known since 1879 when Beamer and Clarke⁶⁰ prepared white, crystalline aniline hydrofluoride. A comprehensive study of amine hydrofluorides was carried out by Berliner and Mann⁶¹, who prepared hydrofluoride salts of aromatic, primary, secondary and tertiary amines. They suggested a structure of base.4HF after titration with sodium hydroxide.

The use of amine hydrofluorides as fluorinating agents remained limited to steroids⁴⁸ until Franz⁶² prepared a series of amine trishydrofluorides, such as Et₃N.3HF. These hydrofluoride complexes were found to be stable, distillable under vacuum, could be handled without hazard and did not corrode borosilicate glass. Since then Et₃N.3HF and other amine trishydrofluorides have been used in a range of fluorination reactions (see Section 1.6.3), similar to those carried out using Olah's reagent (Section 1.5.2). Spectroscopic studies of these systems are discussed here followed by their use in synthesis.

1.6.2 Spectroscopy of Amine.HF Systems

The structure of R₃N.nHF systems (R = alkyl) is not very well understood. Unlike the case of pyridine/HF, no crystallographic data of the complexes present in R₃N.nHF mixtures has been published. N.m.r. spectroscopy has been used to investigate these systems in solution and so a discussion of the n.m.r. and also the i.r. of R₃N.nHF and related systems, such as F⁻, HF and HF₂⁻ containing species, follows.

1.6.2.1 Nuclear Magnetic Resonance Spectroscopy

Before we can discuss the n.m.r. spectra of R₃N.HF salts, we need to review the literature data concerning the species that may be involved in solutions of these salts i.e. F⁻, HF and HF₂⁻

The literature covering the n.m.r. of fluoride ion in solution is very confused; different authors quote different signs for upfield and downfield, references, solvents and concentrations, all of which affect the fluorine shift. The situation was improved on the publication of a review by Hudlicky⁶³, who repeated some of the n.m.r. experiments.

The following table is taken from that review⁶³, and shows the variation of fluorine shift for each species as measured by various authors.

Table 5 N.M.R. shifts of F^- , HF and HF_2^- . Chemical shifts are given in negative values of ppm upfield of $CFCI_3$.

<u>Species</u>	F^-	HF	HF_2^-
KF(aq)	-124.8		
KF(aq)	-120.2		
KF(aq)	-117.5		
Pr_4NF	-114.6		
HF(aq)		-204.0	
HF(anhyd)		-196.0	
Bu_4NHF_2			-144.0
Pr_4NHF_2			-149.4

From this table we can make the general assumption that fluoride ion has a shift between -114-125ppm; HF between -160-200ppm and HF_2^- between -144-149ppm.

However, we must note that the shifts for fluoride ion were measured in either aqueous or aqueous organic solutions, and so are not really due to fluoride ion alone because of hydrogen bonding between F^- and water. This was confirmed by Christe⁶⁴, who has very recently prepared anhydrous tetramethylammonium fluoride (Me_4NF). ^{19}F n.m.r. of this compound in solution (MeCN) showed a singlet at -73.2ppm, significantly shifted from other values for F^- by about 40ppm! However, Me_4NF was found to react with the acetonitrile³² to form HF_2^- (section 1.3.2.1), so the fluoride ion is still not "naked" but must hydrogen bond with acetonitrile as F^- is such a strong base.

Christe concluded that the large upfield shifts noted for fluoride ion in aqueous solution was probably due to the presence of HF_2^- , HF or both; the shift of the singlet, broadened by the rapid exchange between these species, depending on the relative molar amounts of each species.

The lowfield shift for anhydrous Me_4NF is supported by solid state measurements on CsF, by Clark⁶⁵. On reducing the water content in CsF samples a downfield shift occurs to the limiting anhydrous CsF case where the shift is -79ppm.

Hence, the amount of water present seems to be a controlling factor on the F^- chemical shift.

Intuitively, we would predict that on going from HF to F^- , thus increasing the electron density on F, an upfield shift would occur, such as in the case of ^{13}C NMR (e.g. an upfield shift is observed on negatively charged carbon atoms in stable fluorinated carbanions⁶⁶). However in this case, on going from HF to F^- , a downfield shift is seen. The change in electron density on F may be negligible due to the small size of the HF molecule.

The shift of F^- in Me_4NF is also dependent on the solvent (e.g. -136.7 in ethanol, -97 in dichloromethane)⁶⁴. This could be attributed to the strength of the hydrogen bonding between the solvent and fluoride ion, but the work of Symons⁶⁷ showed that the trends in F^- shift (^{19}F NMR) in a series of different solvents was similar to the trends in the shifts of Xenon (^{129}Xe NMR), as compared to ^{35}Cl NMR data. Xenon would not be expected to hydrogen bond with the solvents, so if the trends in chemical shifts of Xe and F^- are analogous then the controlling factor behind the shifts must be the same, but not the strength of the hydrogen bonding with the solvent. However, the factors which effect the shifts in the Xe case are not stated. The ease of solvation of F^- may be an important factor.

However as a general rule we may observe that F^- , HF and HF_2^- come in the following shift ranges:-

F^- : -75ppm (anhydrous, ion-pair, e.g. $Me_4N^+F^-$)

F^- : -114-125ppm (in presence of HF and HF_2^- , broad due to exchange)

HF_2^- : -144-149ppm ($J_{HF}=122$ Hz)

HF : -160-200ppm (broad due to exchange)


1.6.2.1.1 $R_3N.nHF$ Systems

The 1H and ^{19}F n.m.r. of $R_3N.nHF$ ($n=1,1.5,2$) systems were studied by Cousseau⁶⁸ at room temperature and at $-80^\circ C$, (Table 6).

Table 6 ^{19}F N.M.R. Spectra of $\text{R}_3\text{N}\cdot n\text{HF}$ SystemsChemical shifts are quoted as negative values upfield of CFCl_3

	<u>Room Temperature</u>	<u>-80°C</u>	
		$\delta_{\text{F}}(\text{F}^-)$	$\delta_{\text{F}}(\text{HF}_2^-)$
$\text{Et}_3\text{N}\cdot\text{HF}$	-150.5	-123.6	-152.0 ($J_{\text{HF}}=138\text{Hz}$)
$\text{Et}_3\text{N}\cdot 1.5\text{HF}$	-153.0	not observed	-151.7 ($J_{\text{HF}}=138\text{Hz}$)
$\text{Et}_3\text{N}\cdot 2\text{HF}$	-158.0		
$\text{Bu}_3\text{N}\cdot\text{HF}$	-151.2	-125.3	-152.0 ($J_{\text{HF}}=138\text{Hz}$)
$\text{Bu}_3\text{N}\cdot 2\text{HF}$	-159.2		

Hence, we may consider that amine.HF systems are a series of equilibria in solution:-



These assumptions agree with the observed singlet seen in the ^{19}F spectrum at room temperature caused by fast proton and fluorine exchange:-



So, in compounds with no "free HF", i.e. $\text{R}_3\text{N}\cdot\text{HF}$, we see signals due to F^- and HF_2^- as the rates of exchanges (3) and (4) are slowed down due to excess amine. Cousseau concluded that $\text{Bu}_3\text{N}\cdot\text{HF}$ exists in the ionic forms in solution which give rise to both F^- and HF_2^- signals.

Apart from the low temperature n.m.r. of Olah on pyridine/HF systems⁴⁷ (no F^- or HF_2^- observed), this appears to be the only n.m.r. study performed on amine.HF complexes.

1.6.2.2 Infra Red Spectroscopy

A short discussion of the infra red data concerning HF and HF₂⁻ systems appears here, as we need to ascertain later whether Amine.HF salts contain these species.

1.6.2.2.1 Hydrogen Fluoride




The i.r. spectrum of hydrogen fluoride in the gas phase as well as in solution (CCl₄) was measured⁶⁹ and the absorption band found to be at 2.61μ = 3820cm⁻¹.

1.6.2.2.2 Hydrogen Difluoride HF₂⁻

The HF₂⁻ anion, a linear triatomic species, can exist either as a symmetrical (centred H) or an asymmetrical (non-centred H) species, as reviewed by Emsley⁷⁰. Most difluoride species have a centred H, e.g. KHF₂, as shown by X-ray crystallography and i.r. spectroscopy^{71,72}. The very short R(F...F) distance was measured to be 225pm. However, not all crystals have a perfectly centred HF₂⁻ anion. p-Toluidinium hydrogendifluoride forms a secondary hydrogen bond (N-H...F-H-F) which displaces the proton from the central position⁷³.

The i.r. absorptions for both symmetrical and asymmetrical difluoride anions are shown in Table 7, data from ref. 70.

Table 7. Infra Red absorptions for Hydrogen Difluoride Species

ν_1 Symmetric Stretch			
ν_2 Bend			
ν_3 Asymmetric Stretch			
<u>Geometry</u>	<u>ν_1</u>	<u>ν_2</u>	<u>ν_3</u>
KHF ₂	centred	inactive	1225, 1274 1450
Pr ₄ NHF ₂	centred	inactive	1255, 1315 1900
p-Me-C ₆ H ₄ -NH ₃ ⁺ HF ₂ ⁻	non-centred	450	1080, 1230 1740

All absorptions measured in cm⁻¹.

If the difluoride ion is centred ν_1 is i.r. inactive. The doublet observed for ν_2 in KHF₂ is due to lattice energy effects lifting the degeneracy of this mode, not due to the


HF₂⁻ being asymmetric. In asymmetric HF₂⁻, the doublet has a separation of 150cm⁻¹, as in the case of p-toluidinium hydrogendifluoride.

1.6.2.2.3 Trialkylamine Hydrogen Fluoride Systems

The IR spectra of Me₃N.nHF and Et₃N.nHF complexes in solid argon matrices at 10K were measured by Andrews⁷⁴. Fundamental vibration modes for both the 1:1 and 1:2 complexes were measured for both systems.

The vibrational modes for R₃N.HF and R₃N.2HF and the measured absorptions for each system are as follows:-

1:1 Complex




ν_s - HF fundamental stretching vibration

ν_1 - HF librational motion

	ν_s / cm^{-1}	ν_1 / cm^{-1}
Me ₃ N.HF	2589	1030
Et ₃ N.HF	2527	-

The trimethylamine HF complex also shows two distinct perturbed C-N stretching modes.

1:2 Complex



ν_{sa} - H-F stretching mode for H_a-F

ν_{sb} - H-F mode between H_b and fluorine in H_aF

	$\nu_{sa} / \text{cm}^{-1}$	$\nu_{sb} / \text{cm}^{-1}$
Me ₃ N.2HF	1870	2548
Et ₃ N.2HF	1889	-


In general the 1:2 complexes were characterised by a strong band near the 1900cm⁻¹ region of "shared" proton vibrations. The vibrational spectra suggest that H_a is shared between N and the inside F. H_b-F is lengthened but H_b is too close to the terminal F for F..H-F to be considered as an asymmetric difluoride ion.

1.6.3 Reactions of Amine Hydrofluorides as Fluorinating Agents

Amine.HF systems, usually $\text{Et}_3\text{N} \cdot 3\text{HF}$, have been used to carry out a range of fluorination reactions similar to those performed with pyridine/HF. A comprehensive review of the reactions involving $\text{Et}_3\text{N} \cdot 3\text{HF}$, $\text{iPr}_2\text{NH} \cdot 3\text{HF}$ and $\text{Me}_3\text{N} \cdot 2\text{HF}$ as fluorinating agents is discussed here. Halex, halofluorination, sulphenylation fluorination, ring opening hydrofluorination and oxidative fluorination reactions are presented.


1.6.3.1 Halogen Exchange Reactions (Halex)

Franz used $\text{Et}_3\text{N} \cdot 3\text{HF}$ to prepare fluoroacetone, cyanuric fluoride, difluorophosgene, oxaly fluoride, and sulphur tetrafluoride⁶². Other fluorinations of activated, chlorinated heterocycles have been performed⁷⁵.




1.6.3.2 Halofluorination


$\text{Et}_3\text{N} \cdot 3\text{HF}$ used in conjunction with N-halosuccinimides is a useful reagent for the halofluorination of alkenes⁷⁶. The reactions are stereospecifically anti-additions and for unsymmetrical alkenes the orientation of addition follows Markownikoff's rule.




Similarly, bromofluorinations of allylic alcohols have been achieved⁷⁷.




In halofluorinations of cyclic dienes, such as cyclodeca-1,5-diene, transannular π -participation of the second double bond is observed⁷⁸.



The same effect is seen in the halofluorination of norbornadiene where a variety of products are obtained⁷⁹.




Similarly, transannular oxygen participation is seen in the halofluorination of 9-oxabicyclo[6.1.0]non-4-ene⁸⁰.



1.6.3.3 Sulphenylation Fluorination


Stereoselective sulphenylation fluorination can be carried out using $(\text{MeSSMe}_2)^+ \text{BF}_4^-$ with $\text{Et}_3\text{N}\cdot 3\text{HF}$ to form β -fluoro thio(methyl)ethers^{81, 82}.




An alternative method of synthesizing β -fluoro thioethers is by first forming the episulphonium chloride by addition of phenylsulphur chloride to the alkene followed by ring cleavage by the fluorinating agent⁸³.

1.6.3.4 Phenylselenofluorination

β -phenylselenofluorides may be synthesized by the reaction of N-phenylselenophthalimide with alkenes in the presence of $\text{Et}_3\text{N}\cdot 3\text{HF}$ in a one-pot reaction⁸⁴.




Similar reactions are seen with alkynes⁸⁵.




1.6.3.5 Ring Opening Reactions

1.6.3.5.1 Aziridines

α,β -fluoroamines may be prepared by the ring opening of aziridines⁸⁶.




The stereochemical course of the reaction depends on the relative stabilities of the aziridinium ion and the open carbonium ion and the nature of the fluorinating agent (acidity and nucleophilicity). In cyclic systems trans-fluoroamines were obtained by the reaction of N-activated aziridines with $\text{Et}_3\text{N}\cdot 3\text{HF}$, whereas the cis compound is obtained with Olah's reagent⁸⁶.



1.6.3.5.2 Aziridinium Ion

Fluorodeoxyglucopyranosides have been prepared from altropyranosides bearing N,N-diallylamine and mesylate groups in a trans configuration^{87, 88}.




Neighbouring group participation by the diallylamino group produces the aziridinium ion which is ring opened by fluoride ion resulting in a 1,2 shift of the nitrogen atom. The diallylamino group is then reduced using hydrogen on a palladium catalyst. Similar reactions have been carried out using this methodology^{89, 90}.

Triflate groups may be replaced by fluorine in carbohydrate systems using this idea⁹¹.


1.6.3.5.3 Epoxides

Epoxide ring systems are regioselectively opened with amine.HF complexes to give the corresponding trans-addition products^{92, 93}.


28



Optically active α,β -fluoroalcohols may be prepared from the corresponding optically active epoxide⁹⁴⁻⁹⁶.




Fluoroalkyl epoxides give similar products⁹⁷.



1.6.3.6 Oxidative Fluorinations

Phenols may be oxidatively fluorinated at the anode to form difluorocyclohexadienones using $\text{Et}_3\text{N} \cdot 3\text{HF}$ with lead tetra-acetate⁹⁸.



1.6.3.7 Other Systems

Mixtures of melamine/HF have been used in hydrofluorination reactions of alkenes^{99, 100} (e.g. cyclohexene gives fluorocyclohexane in 98% yield). Recently, amine.HF systems have been used as catalysts in conjunction with chromium chloride impregnated support materials in the fluorination of carbon tetrachloride by HF to give a mixture of chlorofluorocarbons¹⁰¹.

Since the writing of this chapter, a review covering a similar subject area has been published¹⁰². This review concentrates on the use of the pyridine/HF system and is not comprehensive with respect to the trialkylamine hydrofluoride systems. Also, no discussion of the structures of hydrofluoride salts is present.

CHAPTER TWO

1.8-Bis(dimethylamino)naphthalene Hydrofluoride (PS/HF) as a Potential Source of Soluble Fluoride Ion

2.1 Introduction

The types of fluoride ion reagent that are currently available have been discussed in Chapter One. However, there remains a need for a source of soluble fluoride ion, as reagents such as the alkali metal fluorides are largely insoluble in organic solvents which limits their use in organic synthesis.

One obvious possibility is the use of amine.HF complexes, as reagents such as $\text{Et}_3\text{N}\cdot 3\text{HF}$ have already been used successfully as fluorinating reagents. Thus, we decided to investigate whether amine.HF systems could act as fluoride ion donors in solution.

However, problems associated with amine.HF complexes in solution are that an equilibrium exists between amine.HF, free HF, HF_2^- and the free amine, and the position of equilibrium will affect the reactivity of the amine.HF complex as fluoride ion donors. Factors which may affect this equilibrium are the base strength and the size of the hydrocarbon moiety of the amines. We therefore wanted to determine whether the proton of HF could be effectively "buried" in the hydrocarbon part of the amine thus leaving the fluoride ion free to react with organic substrates.


Hence, a series of amine.HF complexes were synthesised and their reactivity as fluoride ion sources monitored. We chose sterically hindered, non-nucleophilic, strong bases as the carriers of HF in order to "bury" the proton as much as possible. The amine.HF complexes were then used as the source of fluoride ion in a range of experiments designed to measure their reactivity as fluoride ion donors. Details of the experiments performed are given in the following discussion.

Initially we investigated the HF complex of Proton Sponge as PS is a very strong base (pK_a of the conjugate acid =12.3), sterically hindered around the basic site and is a large molecule which should aid the solubility of the HF complex in organic solvents. A discussion on the preparation, structure, basicity, spectroscopy and reactions of PS follows.

2.2 1,8-Bis(dimethylamino)Naphthalene (Proton Sponge)

2.2.1 Preparation

1,8-Bis(dimethylamino)naphthalene was first obtained as an oil by Brown¹⁰³ who reduced 1,8-dinitronaphthalene by tin chloride/HCl to form the diamine which was then methylated using excess methyl sulphate. Alder obtained the product as a solid (m.p. 47-48°C) by extracting into an aqueous pH 8 solution¹⁰⁴. An improved synthesis was published in 1972 when the diamine was methylated using excess dimethyl sulphate in the presence of sodium hydride¹⁰⁵:-



1,8-Bis(dimethylamino)naphthalene is now marketed by Aldrich under the name "Proton Sponge", and will be referred to as PS in the following discussion.

A series of other N,N,N',N'-tetrasubstituted compounds have been prepared from the parent diamine in a similar manner¹⁰⁶.


2.2.2 Structure

The structure of PS is especially interesting because of the steric effects that are encountered. The 1,8-naphthalene substituents are said to be *peri* to each other and are much closer to one another than to ortho substituents on the aromatic ring.


Steric strain of *peri* substituents may be overcome by (1) a stretching of the bonds; (2) in-plane deflection of substituents; (3) out-of-plane deflection; (4) distortion or buckling of the ring. A stretching of the bonds is ruled out because of the high energy required for such a process and so a compromise situation is reached. Other factors in the case of PS are (1) there is no possibility of bringing even one of the dimethylamino groups into the plane of the ring; (2) if the methyl groups are out of each others way then the nitrogen lone pairs will be brought face-to-face; (3) If the nitrogen lone pairs are favourably situated, one pair of the methyl groups will interfere strongly with a nitrogen atom.

Einspahr and Robert carried out an X-ray crystal structure determination (Fig. 2) to investigate these effects¹⁰⁷.

Fig. 2 X-Ray Crystal Structure of 1,8-Bis(dimethylamino)naphthalene



A representation of the molecule viewed along the C(9)-C(10) bond, including selected intramolecular non-bonded distances. E.s.d.'s are about 0.05 Å for the C-H and N-H distances and about 0.07 Å for the H-H distances.



From the structure we can see that the naphthalene ring is non-planar and the molecule has accommodated the bulky *peri* substituents with surprisingly little strain. The C(ring)-N bonds have retained a significant amount of *p*-character, and two of the N-C(methyl) bonds remain in the plane of the ring. Close interlocking of the hydrogen atoms is the key to the molecular conformation, the atoms of the methyl groups being neatly staggered with respect to the ring hydrogen atoms.

2.2.3 Structure of 1,8-Bis(dimethylamino)naphthalene Salts


Protonation of PS causes the following modifications to the structure¹⁰⁸; (1) the N-N distance is shortened as lone pair repulsions are eliminated (this is one contributing factor to the high basicity of PS); (2) the naphthalene ring becomes more planar; (3) in the N-H-N bridge the proton does not exist in a straight line between the nitrogen atoms.

Errors in X-ray crystallography limits the evidence for the proton existing in a symmetric or unsymmetrical N-H-N bridge. N_{1s} ESCA studies¹⁰⁹ on PSH⁺BF₄⁻ show two inequivalent nitrogen atoms which suggests an unsymmetrical N-H-N bridge. This is supported by positive proton NMR isotope effects¹¹⁰.

2.2.4 Basicity

PS (pK_a of the conjugate acid = 12.3) has a much greater basicity than the parent 1,8-naphthalenediamine (pK_a 4.6) and ten million times greater than that of N,N-dimethylaniline (pK_a 5.1). The reasons for this huge increase in basicity have been of considerable interest. Basicities of some substituted 1,8-diaminonaphthalenes are listed below (Table 8).

Table 8. pK_a Values for Proton Sponge Conjugate Acids and Related Compounds



R	R ₁	pK_a
Me	H	12.1
Et	H	12.7
Me	OMe	16.1
Et	OMe	16.3

The high basicity of PS type compounds is considered to be due to the following reasons; (1) relief of lone pair-lone pair repulsions on protonation resulting in a decrease in the steric strain of the system; (2) protonation gives a monocation with a very stable intramolecular hydrogen bond; (3) the naphthalene ring is forced to be more planar on protonation; (4) recent calculations suggest that the basicity is due to the destabilisation of the neutral form of PS because of its inability to form a hydrogen bond between the methyl groups and the nitrogen atoms¹¹¹.

The increase in basicity on addition of 2,7-methoxy substituents is attributed to the "buttressing" effect, the methoxy groups forcing the naphthalene ring to adopt a more planar configuration in the free base thus giving a greater relief in strain on protonation^{112, 113}.

The high basicity of PS is accompanied by very low rates of proton transfer¹¹⁴ due to the intramolecular hydrogen bond which is stabilised by hydrophobic NMe₂ groups. Proton transfer is considered to be a two step process; the hydrogen bond is broken followed by deprotonation. This low rate of proton transfer means that PS has not found many uses as a base catalyst in organic syntheses (section 2.2.6).

2.2.5 Spectroscopy

2.2.5.1 Ultraviolet Spectroscopy

The long wavelength band of PS in the UV spectrum appears at 335nm (log ϵ_{\max} 3.96) which is shifted to 285nm (log ϵ_{\max} 3.78) on protonation¹⁰⁴.

2.2.5.2 Infra Red Spectroscopy

The infra red spectra of PS salts have been studied by Polish workers¹¹⁵⁻¹¹⁷. The protonic absorption band due to the protonic vibrations in the potential well between the two nitrogen atoms, i.e. N-H⁺-N, is found at low frequencies¹¹⁵, e.g. PS/HBr at 543cm⁻¹. The absorption is dependent on the counter ion which interacts with the N-H-N bridge. A shift to higher frequencies is seen with an increase in the base strength of the anion and the isotopic ratio (ISR), ν_H/ν_D , was found to decrease as the frequency increased. N-H-N stretching absorptions and ISR ratios are listed in Table 9.

Table 9. N-H-N Infra Red Stretching absorptions in Proton Sponge Salts.


$\begin{array}{c} X^- \\ \vdots \\ N-H^+-N \end{array}$	$\begin{array}{c} X^- \\ \vdots \\ N-H^+ \\ \diagup \quad \diagdown \\ N \quad N \end{array}$	
Low absorption High ISR	High absorption Low ISR	
Acid	ν_s / cm^{-1}	ISR
HBF ₄	463	1.85
HPF ₆	479	2.05
HI	509	2.0
HBr	543	1.7
Pentachlorophenol	590	1.0

The rich structure of this band was due to the coupling with other low frequency, internal vibrations of the naphthalene ring. Except for a narrow band at 1200cm⁻¹, there are no absorptions at a higher frequency that may be due to protonic vibrations.

The IR spectra of the salts in acetonitrile solutions have been investigated¹¹⁷. In solution the protonation of PS is seen in the absorptions of the methyl groups. The N-Me bands at $2900\text{-}3000\text{cm}^{-1}$ disappear for PS.HBF₄ as the base is still fully protonated in solution. Absorptions due to N-Me may still be seen if some free base remains in solution, as is the case for PS.Pentachlorophenol. The absorption at 1576cm^{-1} in the free base due to the asymmetry of the buckled naphthalene ring also decreases in intensity when the base is protonated as protonation causes a flattening of the ring removing the asymmetry. The protonic absorption band is a continua extending from $300\text{-}3000\text{cm}^{-1}$. The interaction of polar solvent molecules cause drastic changes in the bridge geometry and hence the potential shape of the proton motion. For all PS salts the broad continua indicates complete dissociation has taken place.

2.2.5.3 Nuclear Magnetic Resonance Spectroscopy

The mode of rotation of the peri NMe₂ groups in the free base is an interesting question because of the steric considerations. At room temperature the methyl protons have a shift of 2.77ppm (singlet) but at -133°C a 1:1 doublet is seen, centred at 2.74ppm ($J = 25.6$ Hz). The symmetry of the spectral changes suggests that conformational changes are taking place, the naphthalene ring flipping from the most stable C₂ conformer through a C_{2v} transition state to the other C₂ conformer¹¹⁸.





On protonation, the proton on the nitrogen is extremely deshielded and has a very high shift, e.g. $\text{PSH}^+\text{BPh}_4^-$ $\delta_{\text{H}^+} = 18.46\text{ppm}$ ¹¹⁰, $\text{PSH}^+\text{CO}_2\text{CF}_3^-$ $\delta_{\text{H}^+} = 19.5\text{ppm}$ ¹⁰⁴.

2.2.6 Reactions

PS is weakly nucleophilic for steric reasons and is recovered unchanged after four days reflux with ethyl iodide¹⁰⁴. PS has been used in organic synthesis as a strong hindered, non-nucleophilic base in base catalysed reactions¹¹⁹⁻¹²⁰, but the slow rate of proton transfer has limited its use.

36



Recently, PS has been shown to react with electrophiles in aromatic substitution reactions but these will be discussed in the introduction to Chapter 4.

2.3 The Hydrofluoride Complex of 1,8-Bis(dimethylamino)naphthalene (Proton Sponge)

2.3.1 Preparation of PS/HF

The PS/HF complex (1) is formed by adding the required 1:1 stoichiometric amount of an HF/ether solution to an ethereal solution of the base. The ether was removed to leave a white solid. Elemental analysis showed that the solid was a 1:1 base:HF complex. The PS/HF complex was completely soluble in acetonitrile and sulpholane. NMR, IR and mass spectrometry were obtained for the complex in an attempt to rationalise its structure and prove whether it exists as a simple salt or as a mixture of base, base.HF, baseH⁺, HF₂⁻ etc. Spectroscopic data is as follows.


2.3.2 Spectroscopical Studies of PS/HF

2.3.2.1 NMR

¹H and ¹⁹F NMR spectra were recorded for the complex (1) dissolved in deuterated acetonitrile at 400MHz. The ¹H spectrum is shown overleaf (Fig 3).

From the ¹H spectrum we can observe the following; (1) two singlets at 2.8 and 3.0ppm; (2) about five peaks in the aromatic region; (3) broad peaks at 13.6 and 18.7ppm. The ¹⁹F NMR spectrum consists of a sharp singlet at -169ppm at room temperature.

From the literature we know that the peak at 18.7ppm is due to the protonated PS (N-H⁺...N) (c.f PSH⁺. CF₃CO₂⁻ at 19.5ppm¹⁰⁴). The origin of the peak at 13.6ppm is uncertain. If it was due to HF₂⁻ being present in solution we would expect to see a doublet in the ¹⁹F spectrum at about -145ppm⁶³ and this is not the case. Hence the peak at 13.6ppm is probably due to free HF. Thus the complex in solution probably exists as a mixture of base, protonated base in equilibrium with HF, as shown below.




The broadened singlets seen at 2.8 and 3.0ppm may be explained as corresponding to PS and to protonated PS. Similarly, five resonances are seen in the aromatic region because of this mixture.

The position of the fluorine resonance at -169ppm is downfield from HF (-180ppm) and upfield from fluoride ion (-112ppm) (from reference 63 and

Figure 3

^1H NMR Spectrum of PS/HF in CD_3CN

**^{19}F NMR Spectrum
-169.0ppm (s)**



established in section 1.6.2.1 for another base.HF system) and indicates an equilibrium between base, HF and fluoride ion.

Low temperature solution state NMR studies were unsuccessful as the solid PS/HF precipitates out of solution at about -20°C .


However we can postulate that the PS/HF complex, a 1:1 complex in the solid state, exists as an equilibrium between the free base, protonated base and HF in solution. These exchanges in solution generate fluoride ion which may be used as a source of soluble fluoride ion as discussed in section 2.4.

2.3.2.2 Infra Red

The literature concerning the infra red spectra of related proton sponge salts both in the solid state and in solution have been discussed in section 2.2.5.2.

The infra red spectra of PS and PS/HF complex were recorded in acetonitrile solution at 0.3 mol l^{-1} concentration. The region between 3000 and 1500cm^{-1} is shown below (Fig. 4).

Figure 4
Solution Infra Red Spectra of PS and PS/HF in MeCN



The Bohlmann bands at 2785, 2840, and 2880 cm^{-1} (A) (lit¹¹⁷. 2780, 2831, and 2869 cm^{-1}), are reduced on protonation but do not completely disappear as for fully protonated molecules¹¹⁷. Hence we can conclude that there is some free base present in solution which is consistent with the ^1H NMR data above. The main difference in the PS/HF spectrum is the appearance of the large broad bands at 1840 and 2050 cm^{-1} (B), which are not seen in other PS salts¹¹⁷. These must be due to N-H-F vibrations. Free HF occurs⁶⁹ at 3820 cm^{-1} and so we are observing the effect of the base complexing with the HF. A broad peak is observed at 580 cm^{-1} which agrees with results for other salts indicating that the N-H-N bridge has a bent geometry as we would expect.

The IR spectrum of PS/HF in a Nujol mull gives different results. A large broad band is seen at 1800 cm^{-1} which must be due to N-H-F vibrations.

2.3.2.3 Mass Spectroscopy

The PS/HF complex is not observed in the mass spectrum, the complex decomposing in the spectrometer even in the FAB mode, the most appropriate method for analysing salts. The peak seen at 215 corresponding to protonated PS is probably due to the free PS being protonated by the FAB matrix (methanol/glycerol). Thus FAB mass spectrometry was unsuccessful in determining the types of species present in the solid state of the PS/HF complex.

2.4 Reactions of PS/HF

A series of experiments were performed using PS/HF as the source of fluoride ion in order to determine whether the complex could act as a fluoride ion donor in solution.

PS/HF was used to catalyse C-C bond forming reactions (oligomerisations, perfluoroalkylations) and in C-F bond forming reactions (nucleophilic substitution reactions with a range of substrates). The results of the reactions performed using PS/HF as the source of fluoride ion are discussed below along with literature results using other sources of fluoride ion for comparison.


2.4.1 PS/HF as a Catalyst in C-C Bond Forming Reactions

2.4.1.1 Oligomerisation Reactions

It is well established that fluoride ion catalyses the oligomerisation reactions of perfluorinated alkenes¹²¹, cyclic alkenes¹²² and alkynes¹²³. The source of fluoride ion is usually KF or CsF as heterogeneous catalysts and so a series of analogous reactions was performed using PS/HF as a homogeneous catalyst.

Tetrafluoroethylene was not oligomerised using PS/HF as the catalyst even when the temperature of the reaction was raised to 110°C and the pressure of the TFE in the reaction vessel was raised to 10 bar. (This reaction was performed at the ICI Experimental Site, Widnes).


Hexafluoropropene was dimerised to its thermodynamic dimer (2), as the only product by GC (72% yield, 100% conversion), at room temperature in two days using acetonitrile as the solvent with the hexafluoropropene in a 7:1 molar excess.




The product dimer separates from the acetonitrile. The solvent layer can then be recharged with hexafluoropropene and the reaction repeated to give the same yield of dimer, demonstrating the catalytic nature of the process. In an analogous reaction hexafluoropropene was dimerised to its thermodynamic dimer (2) in 95% yield under the same conditions when CsF was used as the source of fluoride ion¹²¹.

Oligomers of perfluorocyclobutene¹²² could not be isolated from its attempted oligomerisation using PS/HF as the catalyst in various solvents and at elevated temperatures.

The co-dimer (3) of hexafluoropropene and perfluorocyclobutene is formed using PS/HF as the catalyst. However, the alkene (3) so formed reacts with the PS to form the annelation product (4).



The co-dimerisation process may proceed by the following route:-




The perfluorinated alkene then reacts further with PS and this reaction is discussed in detail in Chapter 4.

The mechanism of these oligomerisations may not be a simple fluoride ion catalysed process as proton transfer from the HF to the intermediate carbanion may take place.

Attempts to form a stable perfluorinated carbanion, by the reaction of hexafluoropropene dimer (2) and PS/HF, failed. Stable perfluorinated carbanions have been synthesised in reactions between perfluoroalkenes and CsF⁶⁶.

2.4.1.2 Perfluoroalkylation Reactions



Fluoride ion catalyses perfluoroalkylation reactions of fluorinated heterocycles the carbanion formed on addition of fluoride ion to the fluoroalkene being trapped by the heterocycle, as outlined in the following mechanism.




Previously, alkali metal fluorides have been used as the source of fluoride ion in perfluoroalkylation reactions between hexafluoropropene and pentafluoropyridine(6)¹²⁴, tetrafluoropyrimidine(9)¹²⁵ and trifluoro-s-triazine(13)¹²⁶.

A series of perfluoroalkylation reactions of fluorinated nitrogen heterocycles by hexafluoropropene using PS/HF as the catalyst were performed and are tabulated below.

(R_F = (CF₃)₂CF)





All products were identified by g.c./m.s. and ^{19}F NMR as compared to the literature data. Yields were determined by g.c. for accuracy.

In an analogous reaction, (6) was perfluoroalkylated¹²⁴ by hexafluoropropene, using KF as the source of fluoride ion, to give (7)(94% yield) and (8)(trace) after heating in sulpholane at 130°C for 12 hr. Similarly, (9) was perfluoroalkylated¹²⁵ by hexafluoropropene, using CsF as catalyst, to give (11)(45% yield) and (12)(43%) after heating in sulpholane at 100°C for 16 hr. Also, (13) gave (14)(39% yield), (15)(51%) and the trisubstituted derivative (16)(5%) using CsF as the catalyst after heating in sulpholane at 70°C for 20 hr.

Comparison of perfluoroalkylation reactions using alkali metal fluorides and those using PS/HF as the source of fluoride ion show that much milder conditions are required when PS/HF is used.

2.4.2 The Use of PS/HF in C-F Bond Forming Reactions

A series of reactions designed to form C-F bonds were carried out using PS/HF as the source of fluoride ion in nucleophilic substitution reactions at unsaturated and saturated carbon.

In all the halogen exchange reactions PS/HX (X = Cl, Br, I) precipitates as the reactions proceed as white solids. Hence the PS may be recovered by heating the PS/HX with base.


2.4.2.1 At Unsaturated Carbon

The reaction between PS/HF and benzoyl chloride gave a 76% yield of benzoyl fluoride at room temperature in acetonitrile after 24 hr. The yield was calculated from NMR integration, referenced to a benzotrifluoride marker.

Hexafluoroacetone reacts with KF to form a carbinolate anion which can then be trapped by electrophiles¹²⁷. The KF-hexafluoroacetone complex was not isolated. PS/HF reacts with hexafluoroacetone to form the same type of species (17) which was observed by ^{19}F NMR.




The shift of the tertiary fluorine was -107.9ppm, deshielded from usual C-F resonances indicating that the fluorine atom is adjacent to an atom bearing a negative charge, as seen in the NMR shifts in perfluorinated carbanions⁶⁶. The complex (17) was trapped by the electrophiles benzoyl chloride and benzyl bromide to give the products (18) and (19) respectively.



2,4-Dinitrochlorobenzene was fluorinated to 2,4-dinitrofluorobenzene by PS/HF in 45% yield after 48 hrs reflux in acetonitrile.

2.4.2.2 At Saturated Carbon


Reactions between PS/HF and octyl iodide, benzyl bromide and 1,2-epoxybutane were performed and the results are shown below:-



Benzyl bromide is the most reactive species towards nucleophilic attack due to resonance stabilisation of the intermediate carbocation.

There is an increasing need for reagents that can introduce fluorine into biologically active molecules. Consequently we attempted to fluorinate the triflate and tosylate derivatives of diacetone-D-glucose (20) which have previously been fluorinated by $\text{Bu}_4\text{NF}\cdot 3\text{H}_2\text{O}$ ¹²⁸ and TAS-F¹²⁹ respectively.

The triflate (21) and the tosylate (22) were prepared by literature methods^{129, 130}, as shown below.



(i) R = Trif, $(\text{CF}_3\text{SO}_2)_2\text{O}$, 0°C , pyridine, CH_2Cl_2

(ii) R = Tos, Tosyl Chloride, r.t., pyridine

There was no fluorination of the tosylate (22) after two days reflux in acetonitrile. The triflate (21) was not fluorinated at room temperature. Refluxing in acetonitrile caused the glucose to decompose and the triflate salt of PS (23) was recovered. No fluorination of the glucose (20) could be detected.

2.5 Summary

We have shown that the PS/HF system is stable, easily prepared, easily handled, does not corrode borosilicate glass and is completely soluble in acetonitrile. and can be used as a source of soluble fluoride ion in a range of reactions. The PS/HF system has been used to catalyse C-C bond forming reactions and in C-F bond forming reactions.

The structure of the PS/HF system in solution is uncertain but probably exists as a mixture of base, base.HF, $\text{baseH}^+\cdot\text{HF}_2^-$ etc. in an analogous way to the $\text{Bu}_3\text{N}\cdot\text{HF}$ system (Chapter 3). Whatever the structure of PS/HF and its mode of action, the ability of PS to bind with HF to produce a source of soluble fluoride ion is clear.

A feature of the PS/HF system is that the PS/HX (X = Cl, Br, I) salts, formed as products in halogen exchange reactions are insoluble in acetonitrile and precipitate out of the reaction mixture.

e.g.



The proton sponge free base can then be regenerated by heating these salts with sodium hydroxide solution.

However, the fact that PS/HF reacted with the fluorinated alkene (3) *in situ* restricts the use of the system. We decided to investigate other base.HF systems as sources of soluble fluoride ion to determine the factors which may affect the ability of base.HF complexes acting as fluoride ion donors, as it is clear from the reactions performed using PS/HF as the source of soluble fluoride ion that amine.HF systems can be used as sources of soluble fluoride ion.

CHAPTER THREE

BASE HYDROFLUORIDE COMPLEXES AS POTENTIAL SOURCES OF SOLUBLE FLUORIDE ION

3.1 Introduction

The success of the PS/HF system as a source of soluble fluoride ion prompted us to synthesize a series of sterically hindered base HF complexes. We hoped to determine the factors which governed the reactivity of these systems as fluorinating agents in order to develop the most efficient reagent. We hoped to distinguish between the effect of increasing the steric hindrance around the basic nitrogen atom and base strength.

After preliminary work in this laboratory⁵ we chose to compare hydrofluoride complexes of trialkylamines, pentamethylpiperidines and tetramethylguanidine.

The literature concerning hydrofluoride salts has been reviewed in Chapter One.

3.2 Trialkylamine Hydrofluoride Complexes

The HF complexes of trialkylamines were prepared in the same way as the PS/HF system (section 2.3.1). i.e adding the required 1:1 stoichiometric amount of an HF/ether solution to an ethereal solution of the base.

The following base.HF complexes were prepared:-

$R_3N.HF$: (24) R=Ethyl, (25) R=Butyl, (26) R=Hexyl, (27) R=Octyl, (28) R=Dodecyl.

3.2.1 NMR Spectroscopy

1H and ^{19}F NMR spectra for each of the $R_3N.HF$ complexes were recorded at room temperature in deuterated acetonitrile. The proton resonance for the H-F proton occurs downfield between 10.8 and 12.7ppm as broad singlets as tabulated below (Table 10). All $R_3N.HF$ complexes give a singlet in the ^{19}F NMR spectrum at room temperature between -154 and -159ppm (Table 10).

Table 10. NMR Chemical Shifts of $R_3N.HF$. 1H shift for HF and ^{19}F Shifts Only

R	δ_H	δ_F
Et	12.7	-154.8
Bu	12.7	-157.1
Hex	12.5	-157.0
Oct	10.8	-156.6
Dodec	11.6	-159.3

No information about the species that are present in solution can be gained from the NMR spectra at room temperature except to say that the HF has bound to the base as the fluorine shift is downfield from free HF. Consequently a study of the low temperature NMR of $Bu_3N.HF$ complex was carried out and is discussed in the next section.


3.2.2 Low Temperature NMR Study of $Bu_3N.HF$

Following the work by Cousseau⁶⁸ we investigated the low temperature NMR spectrum of $Bu_3N.HF$ at $-80^\circ C$ in an attempt to ascertain the species that are present in solutions of Base.HF complexes. The spectrum obtained for $Bu_3N.HF$ complex at $-80^\circ C$ is shown overleaf.


Using the information discussed in Section 1.6.2.1.1 the following assignments may be made:-

<u>Peak</u>		<u>Assignment</u>
-65.38		Ion Pair e.g $Bu_3NH^+F^-$
-111.99		Fluoride Ion F^- hydrogen bonding with the solvent ⁶³
-126.56		Si-F or B-F from etched NMR tube
-131.42		SiF_6^{2-} from etched NMR tube ⁶⁴
-144.97	triplet $J_{HF} = 123$ Hz	$H_2F_3^-$
-147.82	doublet $J_{HF} = 147$ Hz	HF_2^- ⁶³
-148.74		$Bu_3N.HF$ Complex ⁶⁸
-150.68		
-174.12		$H(HF)_n^+$ type species ⁶⁸

Figure 5 ^{19}F NMR Spectrum of $\text{Bu}_3\text{N}\cdot\text{HF}$ Complex at -80°C in d_6 -Acetone



The NMR spectrum at -80°C shows that there are a number of species present in solutions of $\text{Bu}_3\text{N.HF}$ systems. The peaks at -112 indicating F^- , at -174 indicating $\text{H}(\text{HF})_n^+$ species and the smaller triplet and doublet at -144 and -148ppm corresponding to H_2F_3^- and HF_2^- respectively proves that a number of exchange processes must be occurring in $\text{Bu}_3\text{N.HF}$ solutions which may be summarised as follows:-




The peak at -112ppm is direct evidence for the existence of fluoride ion in $\text{Bu}_3\text{N.HF}$ solutions. All the exchange processes listed above must be occurring very rapidly for a singlet to result in the ^{19}F NMR spectrum at room temperature. It seems reasonable to assume the same model for all amine.HF systems in solution.

3.2.3 Reactions

Three standard experiments were carried out using each of the Base.HF complexes as the source of soluble fluoride ion. The reactions chosen and conditions are given below:-


1) Benzoyl Chloride



2) Benzyl Bromide



3) 2,4-Dinitrochlorobenzene



All reaction yields were calculated by ^{19}F NMR integration on the Bruker AC250 spectrometer operating at 235 MHz with reference to a benzo trifluoride marker. This procedure was adopted to avoid loss of products in reaction work-up and hence provide a more accurate picture of the fluorinating ability of the base.HF salts.

Yields for the three standard reactions using trialkylamine.HF complexes (24)-(28) as sources of fluoride ion are collated in Table 11. In all cases the Base.HX (X=Cl, Br) salts produced do not precipitate from solution unlike the PS case (section 2.4.2).

Table 11. Reaction Yields Using Trialkylamine Hydrofluoride Complexes as Sources of Fluoride Ion


<u>Base.HF</u>	<u>PhCOCl</u>	<u>PhCH₂Br</u>	<u>Dinitrochloro- benzene</u>
(24)	91	18	34
(25)	88	12	84
(26)	90	17	77
(27)	68	14	61
(28)	79	11	14

From the table we can make the following general observations:-

- (1) Benzoyl Chloride is easily fluorinated in good yield by all base.HF complexes.
- (2) Benzyl bromide did not fluorinate in good yield indicating that these systems are probably not very good reagents for $\text{S}_{\text{N}}2$ processes.
- (3) $\text{Bu}_3\text{N.HF}$ appears to be the most efficient fluorinating agent. The $\text{Dodec}_3\text{N.HF}$ shows less reactivity probably due to its lower solubility.


3.3 Tetramethylpiperidinium Hydrofluoride Complexes

2,2,6,6-Tetramethylpiperidine is a very hindered, non-nucleophilic base due to the steric constraints around the nitrogen atom caused by the α -methyl groups. N-alkylation increases this effect.



Hence, we synthesized a series of pentaalkylpiperidinium hydrofluoride complexes to determine whether these complexes are convenient sources of soluble fluoride ion.


A large literature exists for tetramethylpiperidine bases and their derivatives as emphasised in a recent review¹³¹. Most of the work conducted in this field was directed towards the development of anti-ganglionic blocking drugs (29) (relieves hypertension) and in the development of iminoxy radicals for use in ESR (30).



The pK_a 's of the conjugate acids of tetramethylpiperidine (31) and pentamethylpiperidine (29) are 11.24 and 11.25 respectively.


3.3.1 Preparation of Penta-alkylpiperidine Bases and Their Hydrofluoride Complexes

Both 2,2,6,6-Tetramethyl- and 1,2,2,6,6-Pentamethyl-piperidine were obtained commercially (Aldrich). The N-ethyl-, allyl- and benzyl-2,2,6,6-tetramethylpiperidines (32-34) were prepared by reacting the parent piperidine (31) with the appropriate tosylate or bromide as shown below¹³².



- (i) Et-OTos, 100°C, 24 hr, 21% yield
 (ii) allyl bromide, 50°C, 3 days, 24% yield
 (iii) benzyl bromide, 100°C, 8 hr, 32% yield

The HF salts of these five tetramethylpiperidine bases were prepared in the same manner as for all other HF complexes. i.e. pipette the required stoichiometric amount of HF/ether solution into an ethereal solution of the base. The following HF complexes (35)-(39) were prepared:-



Each complex was totally soluble in acetonitrile.

3.3.2 NMR Spectroscopy

Both proton and fluorine NMR spectra of each salt (35)-(39) were recorded in deuterated acetonitrile at room temperature. The resonances corresponding to HF in the proton spectra are collated in Table 12 along with the shifts of the singlets seen in the fluorine spectra.

Table 12. NMR Chemical Shifts for Poly-alkylpiperidine.HF Complexes. ¹H Shift for HF in Base.HF and ¹⁹F shifts Only

Base. HF	δ_{H}	δ_{F}
(35)	9.30	-135.0
(36)	12.7	-148.9
(37)	11.1	-143.9
(38)	12.1	-146.0
(39)	11.7	-152.4

Again, no information about the nature of the species in solution can be obtained from the NMR spectral data at room temperature except to say that HF has bound to the base to form a complex.

3.3.3 Reactions

The same three standard reactions were performed as described in section 3.2.3 using the piperidine hydrofluoride salts (35)-(39) as the sources of soluble fluoride ion. In all cases the Piperidine. HX (X = Cl, Br) produced in the nucleophilic substitution reactions remain in solution in contrast to the PS/HF system.

Yields of the reactions are collated in Table 13.


Table 13. Reaction Yields Using Pentaalkylpiperidine Hydrofluoride Complexes (35)-(39) as Sources of Fluoride Ion

<u>Base.HF</u>	<u>PhCOCl</u>	<u>PhCH₂Br</u>	<u>Dinitrochloro- benzene</u>
(35)	83	25	49
(36)	64	84	78
(37)	78	38	94
(38)	55	no reaction	31
(39)	42	no reaction	23

From the table we can make the following general observations:-

(1) Both methyl and ethyl tetramethylpiperidine HF complexes are good sources of fluoride ion showing good reactivity. These bases have about the same pK_a as tetramethylpiperidine and so there appears to be increased reactivity with increased steric hindrance around the nitrogen atom in these systems.

(2) Allyl and benzyl tetramethylpiperidine HF complexes show little reactivity as sources of fluoride ion. This is perhaps due to the following exchange processes occurring in solution decreasing the availability of the fluoride ion :-



3.4 Tetramethylguanidine Hydrofluoride Complex

Tetramethylguanidine is a very strong base (pK_a of the conjugate acid = 13.6)¹³³ and so it seemed a good alternative to PS in forming a hydrofluoride complex to use as a source of soluble fluoride ion. (PS/HF was found to react with certain substrates which limit its use (section 2.4.1.1)).

The HF complex of tetramethylguanidine was prepared in the same way as for other HF complexes and the two reactions were performed using this complex (40) as the source of fluoride ion. Benzoyl chloride gave benzoyl fluoride in 65% yield and dinitrochlorobenzene gave dinitrofluorobenzene in 36% yield.

Attempts to prepare pentamethylguanidine (41) were unsuccessful. The reaction between tetramethylguanidine and methyl iodide gives pentamethylguanidinium hydriodide (42), in agreement with literature results¹³³. Attempts to remove the HI to generate pentamethylguanidine resulted in the formation of a mixture of tetramethylguanidine and pentamethylguanidine which could not be separated.

3.5 Conclusions


We have shown that a series of sterically hindered strong bases may be coupled with HF to produce complexes that may be used as sources of soluble fluoride ion in a variety of reactions.

The most effective fluorinating reagents were the complexes of the strongest bases e.g. PS, ethyl-tetramethylpiperidine.

To ascertain why these are the most effective reagents we need to find out the species involved in solutions of these complexes. Low temperature NMR is the best method for doing this but was unfortunately unsuccessful for these systems. However,


a ^{19}F NMR spectrum of $\text{Bu}_3\text{N.HF}$ complex was obtained at -80°C . From the spectrum we have shown that a series of equilibria are present forming a variety of species.

Thus we can postulate that the success of a reagent as a fluoride ion source depends on the following equilibrium:-



For a good fluoride ion source k_1 must be large and k_{-1} small. The equilibrium is improved by increasing the base strength of the base (increases k_1) and the steric hindrance around the basic nitrogen atom (reduces k_{-1}).

However the situation is complicated by the fact that other equilibria are occurring as seen in low temperature NMR.



Whether the presence of HF_2^- effects the reactivity of these reagents in these types fluorination reactions is unclear, but $\text{Bu}_4\text{N}^+.\text{HF}_2^-$ has been shown to be an effective source of fluoride ion in its own right¹³⁴.

Comparison of the reactivities of the tetramethylpiperidine HF systems suggests that increasing the steric hindrance around the nitrogen atom causes an increase in fluoride ion reactivity.

These preliminary studies give some idea about the nature of the reactivity of amine.HF complexes as sources of fluoride ion in that amine.HF complexes exist in solution as an equilibrium between numerous species which we have identified by low temperature NMR. Increasing the steric hindrance around the basic site in the amine appears to increase reactivity of the amine.HF systems as sources of fluoride ion.


We have shown that amine.HF complexes can act as fluoride ion donors in solution in several nucleophilic substitution reactions, but the amine.HF systems can not be compared with alkali metal fluorides as their mode of reactivity involves an equilibrium situation.

CHAPTER FOUR**ANNELATION REACTIONS BETWEEN 1,8-BIS(DIMETHYLAMINO)-NAPHTHALENE (PROTON SPONGE) AND FLUORINATED ALKENES****4.1 Introduction**

Proton Sponge (PS) was found to react with fluorinated alkenes via the naphthalene ring in electrophilic substitution reactions to form novel products. Reactions of aromatic systems with electrophiles are well known¹³⁵ but only a few reactions between PS and electrophiles have been recorded and are reviewed below. Reactions between soft carbon nucleophiles with fluorinated alkenes have not been reported, to our knowledge.

4.2 Electrophilic Substitution Reactions of 1,8-Bis(dimethylamino)-naphthalene (PS)**4.2.1 Nitration**


PS is nitrated at the 4 position by a mixture of concentrated nitric and sulphuric acids. However a mixture of nitric and acetic acid produces the 1,4,5,8-tetra nitrated product¹³⁶.



4.2.2 Bromination


PS is brominated to 4-Bromo-1,8-bis(dimethylamino)naphthalene by using bromine in conjunction with an iron/iron (III) chloride catalyst in 39% yield¹³⁷. The same product is realised in greater yield (80%) on bromination with bromine in sulphuric acid¹³⁸.

The Grignard reagent of this 4-bromo derivative may be prepared and may undergo coupling reactions with other aromatic bromides to form potential monomers¹³⁷.




4.2.3 Alkylsulphination

PS is alkylsulphinated in the 4 position¹³⁹.




4.2.4 Formylation

PS is formylated using the complex of dimethylformamide and phosphorus oxychloride (the Vilsmeier reagent) as formylating agent¹³⁸. 4,5-diformyl-1,8-bis(dimethylamino)naphthalene is formed which then undergoes an intramolecular Cannizzaro reaction to produce a naphthopyranone derivative in 36% yield. This reaction is an example of an annelation reaction involving PS.



4.2.5 Reaction with Dinitrobenzofurazan (DNBZ)


PS reacts with the very strong electrophiles dinitrobenzofuroxan (DNBF) and dinitrobenzofurazan (DNBZ) to form zwitterionic products¹⁴⁰, which exist as two conformers, proved by low temperature n.m.r, with restricted rotation around the C4-C7' bond¹⁴¹.



4.3 Reactions of 1,8-Bis(dimethylamino)naphthalene with Fluorinated Alkenes (3) and (43)


The success of the PS/HF system in catalysing the dimerisation of hexafluoropropene (section 2.4.1.1) prompted us to try to prepare the co-dimer of perfluorocyclobutene and hexafluoropropene using PS/HF as the catalyst.

PS/HF does indeed catalyse the co-dimerisation but the alkene so formed reacts further with the naphthalene ring of PS in an electrophilic substitution reaction to form two solid products which could be separated by column chromatography. The major product, an orange solid (4) was isolated in 21% yield. A red solid (5)(trace) was also isolated but remains unidentified.



This reaction therefore revealed a range of possible reactions between tertiary aromatic amines and perfluorinated alkenes. Thus, the direct reaction between PS and the co-dimer (3) gave the same product (4) in a comparative yield (22%).

Similarly PS reacts with the fluoroalkene (43) to give the annelation product (44), another orange solid, in 13% yield.



4.3.1 Structure Elucidation

Before any mechanistic considerations (section 4.4) we must prove the structures of (4) and (44), which was done using all available spectroscopic techniques. As (4) and (44) have similar structures we will consider their spectral properties together.


Both (4) and (44) gave satisfactory elemental analyses for $C_{21}H_{16}N_2F_{10}$ and $C_{23}H_{16}N_2F_{12}$ respectively and mass spectra in the Cl^+ and Cl^- modes.

4.3.1.1 NMR Spectra of (4) and (44)

1H NMR - The 1H NMR spectrum of (4) consists of two singlets at 2.81 and 2.88ppm each having a relative intensity of six, corresponding to the NMe_2 groups, and two AX systems in the aromatic region (6.79 - 7.85ppm) each with a relative intensity of two, thus corresponding to four aromatic protons. The two AX systems ($J_{AX} = 8.3$ and 8.8 Hz) indicate that the naphthalene ring must be a 1,4,5,8-tetrasubstituted derivative.

Similarly, the 1H NMR spectrum of (44) consists of two singlets (2.88, 2.94 ppm) and two AX systems (6.86-7.52ppm, $J_{AX} = 8.4$ and 8.8Hz).

These results agree with a similar tetrasubstituted PS system (45)¹³⁸.




(4) is an unsymmetrical molecule and so the NMe_2 groups each give a singlet. The peak which is further upfield is assigned to the NMe_2 group with the more electronegative substituent on the 4-position of the same benzene ring which gives a deshielding effect, i.e we assume that $(CF_3)_2C$ is a more electron withdrawing group than the cyclobutene ring as more fluorine atoms are present in this substituent. Similarly, the aromatic protons are assigned in this way.

^{19}F NMR - ^{19}F NMR was essential in proving the structures for (4) and (44) and hence the orientation of initial nucleophilic attack by PS on the fluoroalkene.

^{19}F NMR for Annulation Product (4)

There are two possible sites of nucleophilic attack by PS on the fluoroalkene which would produce different products according to the mechanism outlined (section 4.4).




(^1H NMR of (4) and (46) would be similar)

(The red solid (5) isolated in small yield shows two singlets in the ^{19}F NMR spectrum at -68.9 and -110.5ppm, and hence could not be (46). The red solid is most probably a hydrolysed product of (4) as the infra red spectrum reveals a peak at 1790cm⁻¹ which may be a carbonyl group. However, attempts to hydrolyse (4) failed.)

We would expect (4) to give a ^{19}F NMR spectrum consisting of three resonances (CF₃, CF₂, CF₂) of relative intensities 6:2:2, and (46) to give a spectrum consisting of four resonances (F, CF₃, CF₂, CF₂) of relative intensities 1:3:4:2. The spectrum obtained consists of three singlets at -67.2 (6F, 2CF₃), -105.1 (2F, CF₂) and -112.5 (2F, CF₂) corresponding to structure (4).

^{19}F NMR for Annulation Product (44)


Again two possible products could be envisaged.




For structure (44) we would expect a ^{19}F NMR spectrum consisting of four resonances (4CF_2) with relative intensities of 2:2:4:4, and for structure (47) a spectrum consisting of five resonances (5CF_2) with relative intensities 2:2:2:4:2. The spectrum obtained consists of four resonances; -104.9 (s, 2F), -114.6 (s, 2F), -135.2 (s, 4F) and -112.8 and -116.0 (AB, $J_{\text{AB}} = 250\text{Hz}$, 4F); the spectrum along with the ^{19}F 2-D COSY spectrum is shown overleaf (Fig 6).

In the AB system, the peaks centred at -112.8ppm are broader than those at -116.0ppm which is probably due to long distance F-H coupling. The fact that this is an AB system is proved by the COSY spectrum, indicating that the peaks at -112.8 and -116.0 are indeed coupled. Thus the COSY spectrum confirms that there are three singlets and an AB system with the required relative intensities for structure (44).

Also, the shifts of the CF_2 groups in the cyclobutene ring substructures are similar for structures (4) and (44) as shown below.

**Figure 6****¹⁹F 2-D COSY NMR Spectrum**




^{13}C NMR - ^{13}C NMR was not very helpful in establishing the structures of (4) and (44) because of the complexity of the spectra between 115 and 120ppm due to C-F coupling. However, peaks may be assigned for the "hydrocarbon half" of the molecules where C-F coupling is absent.

^{13}C spectra were assigned using the chemical shifts for published, unsubstituted compounds as models¹⁰⁶, and are collated in Table 14.

Table 14. ^{13}C NMR Data for 1,8-(bisdimethylamino)naphthalene Compounds (Aromatic Ring only)

The following numbering system has been used for the naphthalene nucleus for both published compounds (48)-(51), (4) and (44). (For (4) and (44) positions 4 and 5 are substituted).



	<u>C-1</u>	<u>C-2</u>	<u>C-3</u>	<u>C-4</u>	<u>C-9</u>	<u>C-10</u>	N(CH ₃) ₂
	<u>C-8</u>	<u>C-7</u>	<u>C-6</u>	<u>C-5</u>			
(48)	150.7	112.6	125.4	121.7	120.5	137.8	44.4
(49)	147.7	108.2	125.4	119.2	119.2	137.4	-
(50)	151.2	113.0	125.3	123.4	120.8	137.7	-
(51)	149.6	113.2	125.6	123.8	120.0	137.8	-
(4)	156.1	111.5	130.4	107.7	121.4	134.1	43.5
	152.9	108.9	126.9	116.7			
(44)	158.2	111.4	135.8	107.4	unass	134.0	43.0
	154.4	109.4	128.6	unass			

The assignment of the peaks is aided by the fact that non-quaternary carbon atoms, i.e. aromatic C-H, have low relaxation times and hence give large peaks. Thus C-2, C-3, C-6 and C-7 are easily assigned. The shifts are similar to those reported but, of course, C-4 and C-5 substitution would alter their shifts slightly.

For structures (4) and (44) C-2 has a greater shift value than C-7 (C-3 > C-6 etc) as C-4 has the more electronegative substituent $((CF_3)_2C)$ causing a deshielding of that part of the naphthalene ring.



C-1 and C-8 all occur at around 150ppm, as seen in table.14.

A full assignment was not possible due to the complexity of the spectra but the four C_{Ar-H} peaks provide further evidence of a tetrasubstituted naphthalene derivative for structures (4) and (44).

4.3.2 Molecular Modelling of Annelation Product (4)

Molecular modelling studies were carried out by Dr. J. Morley, ICI. and predicted structures are shown overleaf (Fig 7).

From the diagrams we see that the naphthalene ring is slightly buckled and the lone pairs do not face each other but are parallel, much the same as in PS itself¹⁰⁷. Hence we would predict that the annelation product (4) to have the same high basicity as PS.


Figure 7**Computer Generated Structure of (4)**

(A) - Side-on View

(B) - Head-on View - looking along the plane of the Naphthalene Ring from above the Dimethylamino Groups


4.4 Mechanism of the Annulation Reaction

The following mechanism has been postulated for the reaction of PS with fluorinated alkenes (shown is formation of product (4)).




4.4.1 Orientation of Initial Nucleophilic Attack

For each fluoroalkene (3) and (43) there are two possible sites for initial nucleophilic attack.




We have proved that pathway (1) is preferred and following is a rationale of these observations.

The potential intermediate species for each pathway are as follows:-



We would expect the angle strain around the spiro carbon atom in intermediates (52A) and (52B) to be much less than in (53A) and (53B). The formation of intermediates via pathway (1) is more energetically favourable and so the orientation of attack may be controlled by the angle strain in the reaction intermediates.

Also, two intermediate carbanions are possible¹⁴².



When the nucleophile is fluoride ion, (54A) is formed exclusively as shown by the formation of stable perfluorinated carbanions¹⁴². The stability of perfluorocarbanions is governed by the substituents on the carbanion centre. Essentially, C⁻-F is destabilising due to electron pair repulsions and C⁻-C-F is stabilising due to the inductive effect¹⁴³. However for fluoroalkenes (3) and (43) both are of the general formula (R_F)₂C=C(R_F)₂ and so the stabilising effects on each carbanion are apparently very similar. Also the Frontier Orbitals at each unsaturated carbon atom will have similar coefficients¹⁴⁴. Therefore we would not expect there to be any selectivity in the direction of nucleophilic attack if the stability of the intermediate carbanions are the controlling factor.

The direction of nucleophilic attack at C₁ rather than C₂ has been attributed to the increased electronegativity of carbon (C₂) in the ring¹⁴³. This may be explained by a consideration of the hybridisation of the orbitals on C₁ and C₂.

sp₂ hybridised orbitals are at 120° angles and sp₃ orbitals are at 109° angles. sp₃ hybridised orbitals have greater p character than sp₂ orbitals, and conversely sp₂ orbitals have greater s character than sp₃ orbitals. Also, s orbitals are more electronegative than p orbitals as s orbitals are closer to the nucleus of the atom.


In the case of (54A) the angle of the orbitals forming the bonds in the ring are constrained to being less than 90°. Hence, these orbitals are more like sp₃ orbitals than sp₂ orbitals and so are rich in p character. Thus, the orbital containing the carbanion must be rich in s character. In the case of (54B) the orbitals forming the bonds with the CF₃ groups are at an angle of greater than 90° and are not as rich in p character as in the case of (54A). So, the orbital containing the carbanion in (54B) is not as rich in s character as in (54A). So, the orbital containing the carbanion in (54A) has more s character than in (54B). Orbitals richer in s character are more electronegative and so the carbanion is more stable in orbitals rich in s character. Thus the most stable carbanion is (54A) and hence the most likely reaction intermediate. Thus the nucleophile initially attacks C₁ for the more stable carbanion (54A) to be formed, in accordance with our findings.

4.5 Reaction of 1,8-Bis(dimethylamino)naphthalene with Perfluorobicyclopentylidene (55)

4.5.1 At Low Dilution


When PS and fluoroalkene (55) were stirred overnight at room temperature in acetonitrile a dark olive green solid precipitated which was purified by recrystallisation to give flat, square, dark green crystals (56).

Mass spectrometry and elemental analysis of this product gave a molecular formula of $C_{24}H_{16}N_2F_{12}$ which is two fluorine atoms less than that of the expected annelation product (57).




No NMR spectra of the product were obtainable and so the structure of (56) is uncertain. Hence single crystal X-ray Crystallography is being attempted for a structure determination.

The molecular formula of the product indicates that defluorination of the starting alkene has taken place. This can occur by reduction of the alkene in a 1-electron transfer process¹⁴⁵.




The diene (58) so formed could then react with PS in the following type of process¹⁴⁵.



In a separate experiment¹⁴⁶ PS reacted with the diene (58) to give the same product as that obtained between the alkene (55) and PS. The reduction of alkene (55) to the diene (58) has previously been performed using a sodium amalgam route¹⁴⁵, and so this reaction shows the possibility of using tertiary amines as reducing agents (electron donors) in reactions of this type.

The most probable structures are shown below.



The structure is most likely to be (56) because of the crystallinity of the product and the formation of molecular ions in the EI^+ , Cl^+ and Cl^- mass spectra. However, an X-ray structure is required.

4.5.2 At High Dilution

Initially we thought that the product arising from the reaction between PS and perfluorobicyclopentylidene (55) gave the polymeric structure (59). Hence we repeated the reaction at high dilution in an attempt to isolate any monomeric products. Three products were isolated by column chromatography, the first compound eluted being (56). The second and third compounds had a remarkable appearance. The second compound eluted was bright green flakes (60) of a metallic lustre which in solution gave a purple colouration and the third compound eluted was purple flakes (61) of a metallic appearance which in solution gave a green colouration! Firstly we must attempt to prove the structures of these fascinating compounds.

4.5.2.1 Green Crystals (60)

Mass spectrometry and elemental analysis suggest a molecular formula of $C_{24}H_{16}N_2F_{12}O$.

1H NMR - The 1H NMR spectrum shows a singlet at 2.16 (12H, NMe_2 groups) and two AX systems in the aromatic region between 6.95 and 7.92ppm.

The broad peak seen between 2.5 and 3.4ppm may be due to complexing between the solvent and the compound. This may explain the change in colour on going from the crystalline form into solution.

Each AX pattern has additional splitting on the more upfield peaks probably due to long-range proton-fluorine coupling.

However, the singlet at 2.16ppm and the two AX patterns in the aromatic region corresponding to four protons indicate the presence of a 1,4,5,8-tetrasubstituted PS system, as before.

^{19}F NMR - The ^{19}F and ^{19}F 2-D COSY NMR spectra are shown overleaf (fig 8).

The spectrum consists of four resonances; an AB system between -111.0 and -113.4ppm (relative intensity 4), a broadened resonance, essentially a singlet or pseudo AB, at -132.3ppm (4F), and two other singlets (pseudo AB) at -117.8 (2F) and -131.4ppm (2F).

The molecular formula suggests that a structure similar to the expected annelation product (57) is present, with a CF_2 group substituted by a carbonyl group. The infra red spectrum reveals a peak at $1720cm^{-1}$ which could correspond to the carbonyl group. Possible structures based on this assumption are shown below.







Figure 8

¹⁹F 2-D COSY NMR Spectrum




We would expect the CF_2 group adjacent to the $\text{C}=\text{C}$ double bond to be more easily hydrolysed and so it is unlikely that the structure is (60A).


The AB system and singlet at -133ppm are similar to the shifts found for the spiro-cyclopentene substructure in compound (44) and are compared below.



The other two singlets are consistent with all three possible structures (60), (60A) and (60B). However, because of the metallic nature of the compound, structure (60) is favoured as conjugation of the π system is spread throughout the molecule. This structure would give the molecule charge-transfer properties as an electron donor group (NMe_2) is connected via a conjugated π system to an electron acceptor group ($\text{C}=\text{O}$). The molecules may then align in a one-dimensional stack, giving the crystalline state a metallic nature¹⁴⁷. These donor-acceptor properties may also explain the change in colour in going from the crystalline state into solution.



A final assignment of the ^{19}F NMR spectrum is as follows:-



The analysis for (60) is slightly in error. This could be due to complexing with water, which is known for α -fluorocarbonyl compounds. If one molecule of water is present for each of (60) the analysis is correct.

4.5.2.2 Purple Crystals (61)

Mass spectrometry and elemental analysis point to a molecular formula of $C_{24}H_{16}N_2F_{10}O$. Hence we would expect a structure similar to (56).

1H NMR - Again we see a singlet at 2.16ppm corresponding to the NMe_2 groups and two AX systems between 6.4 and 7.4ppm, which point to a 1,4,5,8-tetrasubstituted PS system.



The appearance of the broadened quartet between 2.6 and 3.1ppm may again be due to complexing of the compound with the solvent, which may explain the colour of the compound in solution. The proton decoupled ^{19}F NMR spectrum shows that proton-fluorine coupling does occur and this may explain the added complexity of the AX systems.


^{19}F NMR - The ^{19}F NMR spectrum is very complex and ^{19}F 2-D COSY spectroscopy was essential in determining which peaks were coupled, and is shown overleaf (fig 9).

The COSY spectrum clearly shows the presence of five AB systems and the integrations confirm that these correspond to five inequivalent CF_2 groups. The molecular formula suggests a structure similar to (56), so we can suggest the following structures, each having five inequivalent CF_2 groups.


Figure 9

¹⁹F 2-D COSY NMR Spectrum






We would expect the chemical shifts of the CF_2 groups in the carbonyl substituted pentene ring to have similar values to those in the green crystals, structure (60). If we take the mid points of the AB systems and compare the shifts with the singlets in structure (61) this is indeed so.




Again structure (61) is preferred because of the charge transfer properties associated with a fully conjugated π system connecting electron donor and acceptor groups.

A final assignment of the ^{19}F NMR spectrum of (61) is below. Shifts corresponding to the mid-points of the AB systems are given.



4.5.2.3 Mechanism of Hydrolysis to form (60) and (61)

Both green (60) and purple (61) crystals are the product of hydrolysis of the expected products via the following proposed mechanism.




The water may have been present in the solvent or hydroxyl groups on the alumina used in the separation may have caused hydrolysis.

4.6 Reactions of Perfluorobicyclobutylidene (62) with Tertiary Aromatic Amines

It was of interest to know whether fluorinated alkenes would react with single ring systems. Hence, the fluoroalkene (62) was reacted with PS, N,N-dimethylaniline and N-methylindole. Although the reactions with the aniline and indole are not annelation reactions, we feel that it is wise to include them at this point as they are reactions between fluoroalkenes and soft, carbon, aromatic nucleophiles.


4.6.1 Reaction with N,N-Dimethylaniline


N,N-Dimethylaniline reacts with (62) to give the electrophilic substitution product (63) as white crystals in 73% yield.



The ¹H NMR spectrum of (63) shows the familiar AA'XX' splitting pattern between 6.76 and 7.16 ppm ($J_{AX} = 8.7$ Hz) indicative of a *para*-disubstituted benzene ring and a singlet at corresponding to the NMe₂ group.

The ¹⁹F NMR spectrum shows five resonances of relative intensity 1:2:2:4:2 which have been assigned in comparison with a similar unsubstituted fluoroalkene (64), as shown below.






(63) gives satisfactory elemental analysis and mass spectra.


4.6.2 Reaction with N-Methylindole

Similarly, N-methylindole reacts with fluoroalkene (62) to give the product (65) as white crystals in 46% yield. The orientation of electrophilic substitution in indoles will be discussed in section 5.5.



The ^1H NMR spectrum of (65) is similar to those of adducts (73)-(75) as discussed in section 5.5.


The ^{19}F NMR spectrum is similar to that of product (63). Satisfactory analysis and mass spectra were obtained.



4.6.3 Reaction with PS

PS and perfluorobicyclobutylidene were stirred together in acetonitrile. Addition of water to the reaction mixture gave a white solid, elemental analysis giving a formula of $C_{22}H_{18}N_2F_8O_2$. Clearly hydrolysis of the expected annelation product had taken place.

^{19}F NMR of the product (66) gives two resonances; a singlet at -120.1ppm and an AB system at 129.8 and 133.8ppm with the same relative intensities. This suggests a symmetrical structure and we have tentatively suggested structure (66).




(6 6)

The singlet at -120.1ppm is assigned to the CF_2 groups adjacent to the carbonyl groups and the AB system to the asymmetric CF_2 groups.

The 1H spectrum is more complex in the aromatic region compared to (4) and (44) and this may be due to isomeric forms of the product. The product (66) decomposes in the mass spectrometer so no confirmation of the structure is possible. The infra red spectrum shows an absorption at $1770cm^{-1}$ which could be attributable to the carbonyl bonds.

The product (66) may be formed by the hydrolysis of the expected annelation product, as follows:-



4.7 SUMMARY

We have shown that tertiary aromatic amines can act as carbon nucleophiles in reactions with perfluorinated alkenes. In particular, annelation reactions between 1,8-bis(dimethylamino)naphthalene (PS) and alkenes (3) and (43) have been performed. A mechanism for this reaction has been proposed and reasons for the direction of nucleophilic attack on the fluoroalkene discussed.

The general reaction may be extended to anilines and indoles. Hence we have discovered a general route for the nucleophilic substitution of fluorinated alkenes by soft carbon aromatic nucleophiles which, to our knowledge, has not been reported previously.

CHAPTER FIVE

REACTIONS BETWEEN TERTIARY AROMATIC AMINES ACTING AS SOFT CARBON NUCLEOPHILES WITH FLUORINATED HETEROCYCLES


5.1 Introduction

The success of the reaction between tertiary aromatic amines and fluorinated alkenes (Chapter Four) prompted us to try reactions between tertiary aromatic amines and fluorinated heterocyclic systems. The reactivity of perfluoroheterocyclic compounds towards nucleophiles is well known¹⁴⁸.

Reactions between trifluoro-s-triazine and soft, aromatic carbon nucleophiles were performed. Some reactions between trichloro-s-triazine¹⁴⁹ and aromatic carbon nucleophiles have been reported and are reviewed below, but we are unaware of any previous detailed investigations of reactions between trifluoro-s-triazine and aromatic amines acting as carbon nucleophiles. The only comparable example that we have found in the literature is the reaction between trifluoro-s-triazine and N-methylpyrrole¹⁵⁰ (see below).

5.2 Nucleophilic Substitution Reactions of Trifluoro-s-triazine

Trifluoro-s-triazine is very reactive towards nucleophiles and reactions with O, N and perfluorinated carbanion nucleophiles have been well studied¹⁴⁸. Some examples of nucleophilic substitution reactions of trifluoro-s-triazine are given below which proceed via the mechanism outlined previously in section 2.4.1.2.



Conditions and Yields

(i) r.t., THF, 2 hr, 74%

(ii) 0°C, Ether, 1 hr, 90%

(iii) r.t., THF, K_2CO_3 , 2 hr, 77%

(iv) 70°C, sulfolane, 20 hr. $n=1$, 39%; $n=2$, 51%; $n=3$, 5%

5.3 Reactions between Trichloro-s-triazine (Cyanuric Chloride) and Tertiary, Aromatic Amines Acting as Carbon Nucleophiles

5.3.1 With N,N-Dialkylanilines and Toluidines

Reactions between cyanuric chloride and N,N-Dialkylanilines were studied by Shaw¹⁵¹. It was found that the aniline acted as a ambident nucleophile to give a mixture of products of types (IIA), (IIIA) and (IVA), as outlined in the following scheme.







Table 15 is taken from this paper¹⁵¹ and a discussion of the results follows. All reactions were carried out in the absence of solvent at a temperature of 90°C for 8 hours.

Table 15. Reactions Between Cyanuric Chloride and N,N-Dialkyl Anilines and Toluidines.

Amine	Conversion (%)	(III) (%)	(II) (%)	(IV) (%)
	92	-	55	45
	95.5	30	70	-
	74.5	40	60	-
	38.6	-	100	-
	75.0	55	45	-
	77.5	-	100	-

From the table the following points may be rationalised:-

- (1) N,N-Dimethylaniline gives only the nitrogen substituted product (II).
- (2) N,N-Diethylaniline gives a 1:2 mixture of carbon substituted product to nitrogen substituted product (II). There is an increase in the amount of carbon substituted product with increasing alkyl chain length on the nitrogen. This was accounted for by the increased steric hindrance around the nitrogen atom and the increasing inductive effects of the alkyl chain making the p-carbon more activated towards electrophilic substitution.
- (3) Both the 2- and 4-methyl toluidines yielded only the nitrogen substituted products. For the 4- isomer reaction can only occur at the nitrogen; no ortho-substitution was observed. With the 2- isomer, the 2-methyl group prevents the NEt₂ group from conjugating with the ring and hence decreases activation towards electrophilic attack and so only the nitrogen substituted product is obtained.


In reactions between N,N-dialkylnaphthylamines and cyanuric chloride the carbon substitution product forms exclusively¹⁵². This was attributed to the greater activation of the naphthalene ring towards electrophilic attack and steric hindrance of the nitrogen by the peri hydrogen atom.

The carbon substituted compounds (II) have been patented¹⁵³.

The UV, IR and ¹H NMR spectra of the carbon substituted compounds were presented in a separate paper¹⁵⁴, but details will be included later in the discussion for comparison with our data.


5.3.1.1 Mechanism

A mechanism for the formation of the carbon substituted compounds of cyanuric chloride has been postulated¹⁵⁵, and is shown below. The reaction is claimed to proceed via successive π and σ complexes. The charge transfer complexes may be seen by UV spectroscopy.





5.3.2 With 5-Membered Heterocyclic Rings


Pyrroles are electron rich heterocycles that react readily with a wide range of electrophiles. Generally, substitution at the 2-position is favoured over the 3-position as the intermediate carbocation formed is more extensively delocalised¹⁵⁶, as shown below.




Reactions between cyanuric chloride and furans, pyrroles and thiophenes have been studied in a series of publications by Chakrabarti and Todd for the Lilley Chemical Company. Initially, a metallated derivative of the unsubstituted thiophene or pyrrole was used to couple with the triazine to produce the desired dihalogeno-heteroaryl-s-triazine¹⁵⁷. The metallation involved either reaction with *n*-butyl lithium or formation of a Grignard reagent, as shown below.




In a subsequent publication¹⁵⁸, the use of a metallated species was found to be unnecessary and a series of pyrrolyl-s-triazines were produced by electrophilic substitution reactions.



Unusually, the reaction of cyanuric chloride with 2-acetyl-1-methylpyrrole gave an equimolar mixture of two products¹⁵⁹.




This reaction proceeds via the intermediate (V) which is the likely product of the reaction of cyanuric chloride with an acetyl derivative¹⁶⁰.



In similar work, Shaw reported the reaction of cyanuric chloride with N-ethyl pyrrole and indole to give carbon substituted electrophilic products¹⁶¹.

The reaction between trifluoro-s-triazine and N-methylpyrrole was recorded¹⁵⁰.




However, no spectral data appears in the publication and no detailed investigation of the reactions of trifluoro-s-triazine appears in the literature.


The pyrrolyl-dichloro-s-triazines have been patented as anti parasitic agents¹⁶² and fungicides¹⁶³⁻¹⁶⁵, showing biological activity against fungi such as anthracnose (*Collectotrichum lagenarium*), rice leaf spot disease and grey mold of grapes (*Botrytis cinerea*), amongst others.

The hydrolysis of (N-methylpyrrol-2-yl)-dichloro-s-triazine was studied¹⁵⁰. Successive hydrolyses to the triazin-2-one and then to the triazine-2,4-dione by sodium hydroxide in water was found.

Pyrazoles also react nucleophilically with cyanuric chloride via a metallated species¹⁶⁶.




However, reactions with the more basic heterocycles such as imidazoles, thiazoles, benzothiazoles, and pyridines do not give electrophilic substitution products but instead give quaternary salts. For example, pyridine reacts with 2-chloro-4,6-dimethoxy-s-triazine to give the unstable chloride salt which slowly hydrolyses to the hydroxide¹⁶⁶.



RESULTS AND DISCUSSION**5.4 Reactions of Fluorinated Triazines with Pyrroles**

Both pyrrole and N-methylpyrrole react with trifluoro-s-triazine (13), perfluoroisopropyl-s-triazine (14) and perfluorodiisopropyl-s-triazine (15) to form electrophilic substitution products (67)-(72), in good yield.



(67) R=H, R₁=F, R₂=F

(68) R=H, R₁=F, R₂=R_F

(69) R=H, R₁=R_F, R₂=R_F

(70) R=Me, R₁=F, R₂=F

(71) R=Me, R₁=F, R₂=R_F


(72) R=Me, R₁=R_F, R₂=R_F

R_F = (CF₃)₂CF-

The products were precipitated from the reaction mixture by adding water and then purified by vacuum sublimation.

Electrophilic substitution occurs at the 2-position of the pyrrole ring, as expected, and this is proved by the values of the proton-proton coupling constants (section 5.4.1.1).

Yields, melting points and UV spectral data for compounds (67)-(72) are collated in table 16, below. Satisfactory elemental analyses and mass spectra were recorded for each compound (67)-(72).

Table 16. Yields, melting points and UV spectra of Pyrrolyl-s-triazines


Compound No.	R	R ₁	R ₂	Yield (%)	m.p. (°C)	λ _{max} (nm) (log ₁₀ ε)
(67)	H	F	F	65	156-160	310.1 (4.48)
(68)	H	F	R _F	71	112-115	336.0 (4.41)
(69)	H	R _F	R _F	68	64-66	344.8 (4.58)
(70)	Me	F	F	45	117-118	314.6 (4.44)
(71)	Me	F	R _F	54	110-111	324.5 (4.38)
(72)	Me	R _F	R _F	48	88-89	349.4 (4.33)

(R_F = (CF₃)₂CF-)

5.4.1 Spectroscopy of Pyrrolyl-s-Triazines

5.4.1.1 ¹H NMR

All products (67)-(72) gave similar proton NMR spectra and are collated in table 17. The spectra were assigned with reference to shifts of similar pyrroles bearing electron-withdrawing substituents at the 2-position taken from the literature¹⁶⁷; two literature compounds are included in table 17, for comparison.

Table 17. ¹H NMR Spectra of Pyrrolyl-s-triazines. Chemical Shifts (J_H, Hz)

Compound	N-H	N-CH ₃	H-2 (J _{2,3})	H-3 (J _{3,4})	H-4 (J _{2,4})
	-	-	6.91 -	6.03 -	6.64 -
	-	-	6.76 -	5.90 -	6.54 -
(67)	10.30	-	7.23 (2.3)	6.40 (4.0)	7.34 (1.6)
(68)	10.62	-	7.27 (2.4)	6.43 (4.0)	7.40 (1.2)
(69)	9.50	-	7.15 (1.7)	6.31 (3.5)	7.29 (1.5)
(70)	-	4.04	7.14 (2.4)	6.27 (4.0)	7.44 (1.8)
(71)	-	4.12	7.35 (2.3)	6.33 (4.2)	7.52 (1.8)
(72)	-	4.07	7.05 (2.3)	6.32 (4.2)	7.66 (1.9)





The ring proton coupling constants for pyrroles are diagnostic of the position of substitution. Typical values for the coupling constants in pyrrole rings are as follows¹⁶⁷:-

$$J_{2,4} \ 1.35-1.80 < J_{2,5} \ 1.95-2.30 < J_{2,3} \ 2.40-3.10 < J_{3,4} \ 3.40-3.80$$


The ¹H NMR spectrum of (70) is shown overleaf (Fig. 10), and is typical of the spectra obtained for compounds (67)-(72). From the spectrum we can see that the resonance at 6.27ppm has coupling constants 2.39 and 4.00Hz, and the resonance at 7.43ppm has coupling constants 1.80 and 4.2 Hz. From the list of typical values of coupling constants above we can assign the resonance at 6.27ppm to be H-3, and at 7.43ppm to be H-4. Hence, the pyrrole has undergone substitution at the 2-position, as expected from the electrophilic substitution mechanism outlined in section 5.3.2.

Figure 10

¹H NMR Spectrum of 2-(N-methylpyrrol-2-yl)-4,6-difluoro-s-triazine



$J_{2,3} = 2.39$
$J_{2,4} = 1.80$
$J_{3,4} = 4.10$



5.4.1.2 ¹⁹F NMR

The ¹⁹F NMR shifts for compounds (67)-(72) are tabulated below (Table 18) along with the shifts of the unsubstituted triazines¹²⁶ (13)-(15), for comparison.

Table 18. ¹⁹F NMR Shifts for Pyrrolyl-s-Triazines.

<u>Compound</u>	<u>Ring F</u>	<u>CF₃</u>	<u>C-F</u>
(13)	-30.4	-	-
(14)	-30.4	-74.4	-183.8
(15)	-30.5	-74.8	-185.2
(67)	-39.7	-	-
(68)	-39.7	-74.4	-184.5
(69)	-	-74.4	-184.7
(70)	-45.6	-	-
(71)	-38.4	-74.2	-184.3
(72)	-	-75.0	-185.2

The spectra are as expected by comparison with the shifts of the parent triazines (13)-(15).


5.4.1.3 ¹³C NMR

¹³C spectra were recorded for compounds (68)-(72), (67) being too insoluble for a ¹³C spectrum to be obtained.

The ¹³C NMR shifts for the carbon atoms in the pyrrole ring were assigned by comparison with similar literature compounds bearing electron withdrawing groups at the 2-position¹⁶⁷. Assignment of these peaks is helped by the fact that there is not any C-F coupling and so the peaks are singlets. Also, peaks due to aromatic C-H are large due to their short relaxation time.

¹³C NMR shifts for the pyrrole ring carbons in the pyrrolyl-s-triazines are collated in table 19. Shifts for two literature compounds are also included for comparison.

Table 19. ^{13}C NMR shifts for Pyrrole Ring Carbon Atoms in Pyrrolyl-s-Triazines (68)-(72)



<u>Compound</u>	<u>N-CH₃</u>	<u>C-2</u>	<u>C-3</u>	<u>C-4</u>	<u>C-5</u>
	-	133.8	121.8	112.3	129.8
	-	133.4	118.5	111.2	127.0
(68)	-	128.5	121.2	114.5	130.5
(69)	-	128.2	121.3	114.3	130.6
(70)	39.2	136.1	123.3	111.1	128.0
(71)	39.3	136.8	123.9	111.5	128.0
(72)	39.1	137.7	124.7	111.9	128.1

The assignment of the peaks for the triazine ring carbon atoms in (68)-(72) is more complex as C-F coupling is present. Typical coupling values for 1 bond C-F coupling ($^1J_{\text{C-F}}$), 2 bond ($^2J_{\text{C-F}}$), 3 bond ($^3J_{\text{C-F}}$) and 4 bond ($^4J_{\text{C-F}}$) are as follows¹⁶⁸:-


$$^1J_{\text{C-F}} = 158 - 408 \text{ Hz}$$

$$^2J_{\text{C-F}} = 0 - 103 \text{ Hz}$$


$$^3J_{\text{C-F}} = 0 - 43 \text{ Hz}$$

$$^4J_{\text{C-F}} = 0 - 24 \text{ Hz}$$

Examples of C-F coupling in a related system are as follows and may be taken as a guide in assigning the spectra of (68)-(72).




For mono-, di- and tri-substituted triazines we would expect the following peaks to be seen in the ^{13}C NMR spectrum, given that carbon couples with fluorine through at least four bonds.



Using these models and expected $J_{\text{C-F}}$ values as a guide the ^{13}C NMR spectra for (68)-(72) have been assigned as shown in table 20.


Table 20. ^{13}C NMR shifts for Triazine Ring Carbon Atoms in Pyrrolyl-s-Triazines (68)-(72)

Monosubstituted Triazine (70)




<u>Compound</u>	<u>C-2</u>	<u>C-4</u>
	t (^3J)	d d (^1J , ^3J)
(70)	172.3 (14)	172.0 (226, 19)

Disubstituted Triazines (68) and (71)



<u>Compound</u>	<u>C-2</u>	<u>C-4</u>	<u>C-6</u>	<u>C-alpha</u>	<u>C-beta</u>
	dd (^3J , ^4J)	dd (^2J , ^3J)	dd (^1J , ^4J)	d sept (^1J , ^2J)	qd (^1J , ^2J)
(68)	170.4 (13, 3)	169.2 (22, 12)	171.2 (229, 3)	90.9 (211, 33)	121.3 (288, 27)
(71)	170.2 (13, 3)	168.2 (21, 12)	170.3 (228, 3)	90.3 (211, 35)	120.8 (288, 27)




Trisubstituted Triazines (69) and (72)

<u>Compound</u>	<u>C-2</u>	<u>C-4</u>	<u>C-α</u>	<u>C-β</u>
	t (4J)	dd (2J, 3J)	d sept (1J, 2J)	qd (1J, 2J)
(69)	166.5 (3)	165.9 (22, 4)	90.4 (211, 33)	120.8 (288, 27)
(72)	166.6 (3)	165.5 (22, 3)	90.2 (218, 33)	120.7 (288, 27)


The complete ^{13}C NMR spectrum of compound (72) is shown overleaf (Fig. 11).

Figure 11

¹³C NMR Spectrum of 2-(N-methylpyrrol-2-yl)-4,6-perfluoro-1-isopropyl-s-triazine




Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Assignment
39.09	s		a
90.21	d sept	218, 33	h
111.88	s		c
120.73	qd	288, 27	i
124.75	s		d
128.12	s		e
137.67	s		b
165.55	dd	22, 3	g
166.62	t	3	f



5.5 Reactions of Fluorinated Triazines With N-Methylindole

N-methylindole reacted with the triazines (13)-(15) to give electrophilic substitution products (73)-(75), respectively, in good yield.




(73) $R_1=F$, $R_2=F$

(74) $R_1=F$, $R_2=R_F$

(75) $R_1=R_F$, $R_2=R_F$

($R_F = (CF_3)_2CF-$)

Indoles are very nucleophilic heterocycles and react easily with electrophiles. Electrophilic substitution occurs preferentially at the C-3 site rather than at C-2, in contrast to pyrrole systems, because the intermediate cation formed by attack at C-3 is more stable than that formed at C-2 as the positive charge may be delocalised without involving the benzene ring part of the molecule¹⁵⁶.



Yields and melting points for compounds (73)-(75) are collated in table 21, below. Satisfactory elemental analyses and mass spectra were recorded for each compound (73)-(75).

Table 21. Yields and melting points of Indolyl-s-triazines

Compound No.	Yield (%)	m.p. (°C)
(73)	78	244
(74)	90	190-194
(75)	87	205-206

5.5.1 Spectroscopy of Indolyl-s-Triazines**5.5.1.1 ¹H NMR**

The ¹H NMR spectrum of (74) is shown overleaf (Fig 12). From the spectrum we can see five protons in the aromatic region. The four multiplets are due to the indole ring protons and the singlet corresponds to the proton at the 2-position of the indole ring, by comparison with literature data¹⁶⁷. This singlet proves that the electrophilic substitution has taken place at the 3-position as long range coupling would be seen if the 3-proton was present.

5.5.1.2 ¹⁹F NMR

¹⁹F NMR spectra for the indolyl-s-triazines were similar to those found for the pyrrolyl-s-triazines and were assigned in the same way. The ¹⁹F NMR shifts for the indolyl-s-triazines (73)-(75) are collated in table 22.

Table 22. ¹⁹F NMR Shifts for Indolyl-s-Triazines.(73)-(75)

<u>Compound No.</u>	<u>Ring F</u>	<u>CF₃</u>	<u>C-F</u>
(73)	-40.2	-	-
(74)	-39.2	-74.2	-184.2
(75)	-	-74.2	-184.4


5.5.1.3 ^{13}C NMR

A ^{13}C spectrum was recorded for compound (74), but (73) and (75) are too insoluble for ^{13}C spectra to be obtained.

The ^{13}C NMR shifts for the carbon atoms in the indole ring were assigned by comparison with similar literature compounds bearing electron withdrawing groups at the 3-position¹⁶⁷. Assignment of these peaks is helped by the fact that there is not any C-F coupling and so the peaks are singlets. Also, peaks due to aromatic C-H are large due to their short relaxation time.

^{13}C NMR shifts for the indole ring carbons in the indolyl-s-triazines are collated in table 23. Shifts for two literature compounds are also included for comparison.

Table 23. ^{13}C NMR shifts for Indole Ring Carbon Atoms in Indolyl-s-Triazines (74)



B	<u>C-2</u>	<u>C-3</u>	<u>C-4</u>	<u>C-5</u>	<u>C-6</u>	<u>C-7</u>	<u>C-8</u>	<u>C-9</u>
COMe	133.4	116.2	122.0	120.9	120.9	111.4	124.4	135.9
CHD	138.1	118.2	123.3	122.0	120.8	112.3	124.2	137.1
(74)	140.5	111.7	127.2	124.0	123.0	112.0	124.6	139.6

The similar ^{13}C shifts of literature compounds and the product (74) is further proof of the position of substitution at the 3-position.

The ^{13}C NMR shifts for the triazine ring carbon atoms were assigned in the same way as for the pyrrolyl-s-triazines. The shifts for the triazine ring carbons in the pyrrolyl compounds (68), (71) and the indolyl compound (74) are tabulated below (Table 24) for comparison.


Table 24. ¹³C NMR shifts for Triazine Ring Carbon Atoms in Indolyl-s-Triazine (74)

(Same nomenclature as in Table 20)

<u>Compound</u>	<u>C-2</u>	<u>C-4</u>	<u>C-6</u>	<u>C-α</u>	<u>C-β</u>
	dd (³ J, 4J)	dd (² J, ³ J)	dd (¹ J, 4J)	d sept (¹ J, ² J)	qd (¹ J, ² J)
(68)	170.4 (13, 3)	169.2 (22, 12)	171.2 (229, 3)	90.9 (211, 33)	121.3 (288, 27)
(71)	170.2 (13, 3)	168.2 (21, 12)	170.3 (228, 3)	90.3 (211, 35)	120.8 (288, 27)
(74)	175.0 (13, 3)	168.1 (multiplet)	170.3 (229)	90.2 (210, 33)	120.8 (287, 27)

5.6 Reaction of Tetrafluoropyrimidine With N-Methylindole

N-methylindole also reacts with tetrafluoropyrimidine, which is less activated towards nucleophilic attack than trifluoro-s-triazine, to give a similar electrophilic substitution product (76) as yellow crystals (m.p. 231°C) in 36% yield.




Satisfactory elemental analysis and mass spectra were recorded. The product was too insoluble for NMR spectra to be recorded.

5.7 Reactions of Fluorinated-s-triazine with Anilines

5.7.1 Reaction of Trifluoro-s-triazine with N,N-Dimethylaniline

Trifluoro-s-triazine reacts with N,N-dimethylaniline to give the carbon-substituted product (77) exclusively, as brown-red crystals (m.p. 234-237°C) in 28% yield.



The ^1H NMR spectrum proves the carbon-substituted structure (77) as shown. The spectrum shows a singlet at 3.10ppm corresponding to 6 protons and an AA'XX' system in the aromatic region (6.80 and 8.28ppm, $J_{AX} = 9.0$ Hz), corresponding to 4 protons, as is usual for a 1,4-disubstituted benzene ring. This agrees with ^1H NMR data for an analogous compound produced by Shaw in the reaction between cyanuric chloride and N,N-diethylaniline¹⁵⁴.



The reaction between cyanuric chloride and N,N-dimethylaniline gave no carbon substituted product¹⁵¹. This shows that trifluoro-s-triazine is more susceptible to nucleophilic attack than trichloro-s-triazine.

5.7.2 Reaction of Perfluoroisopropyl-s-triazine (14) with N,N-Dimethyl- and Diethyl-aniline


Similarly perfluoroisopropyl-s-triazine (14) reacts with N,N-dimethylaniline to produce the carbon-substituted product (78) in 36% yield. Again, ^1H NMR proves the structure of (78) and is included in table 25.

However the reaction between N,N-diethylaniline and triazine (14) is not as simple. The same work-up procedure was used to obtain a yellow solid which gave elemental analysis and mass spectra consistent with the formula $\text{C}_{16}\text{H}_{14}\text{N}_4\text{F}_8$. ^1H and ^{19}F NMR revealed the presence of two products which must be isomers from the elemental analysis. The ^1H NMR spectrum of this product mixture is shown overleaf.

¹H NMR Spectrum of N,N-(Diethylamino)-perfluoroisopropyl-s-Triazine
Ortho and Para Isomers



The major product seen in the NMR spectra is the expected *para* substituted product (79A) which gives a triplet and a quartet at 1.20 and 3.49ppm respectively and an AA'XX' system at 6.80 and 8.27ppm ($J_{AX}=9.6\text{Hz}$) in the ^1H NMR spectrum. The minor product must be the *ortho* substituted isomer (79) as shown below.



(78) $\text{R}=\text{Me}$, 36% yield, only A formed.


(79) $\text{R}=\text{Et}$, 45% yield, A:B = 69:31.

The nitrogen substituted product (80) was prepared directly from *N*-ethylaniline and triazine (14) as a white solid (section 5.9) for comparison of NMR data.

The formation of *ortho*-substituted products has not been noted before in reactions between cyanuric chloride and anilines¹⁵¹ (section 5.3.1).

5.7.3 Reaction of Perfluorodisopropyl-*s*-triazine (15) with *N,N*-Dimethyl- and Diethyl-aniline

Reactions between *N,N*-dimethyl- and diethyl-aniline and triazine (15) both gave a mixture of isomers (81A)-(82B).



(81) $\text{R}=\text{Me}$, 77% yield, A:B = 44:56

(82) $\text{R}=\text{Et}$, 72% yield, A:B = 63:37

5.7.4 Collected NMR Data For Phenyl-s-Triazines (77)-(82)

Although products (77)-(82) could not be separated from their respective *ortho* isomers, the ^1H and ^{19}F NMR shifts for these compounds could be assigned from the spectra of the mixtures of isomers (Table 25).

Table 25 ^1H NMR Spectra of Phenyl-s-triazines (77)-(82A)

<u>Compound</u>	<u>N-Me</u>	<u>CH₃</u>	<u>CH₂</u>	<u>AA'XX'</u> <u>System</u>	<u>J_{AX} (Hz)</u>
(77)	3.10	-	-	6.80, 8.28	9.0
(78)	3.40	-	-	6.98, 8.65	9.4
(81A)	3.08	-	-	6.67, 8.36	9.2
(79A)	-	1.20	3.49	6.80, 8.27	9.6
(82A)	-	1.18	3.42	6.65, 8.33	9.2

Table 26 ^{19}F NMR Spectra of 4-Phenyl-s-Triazines (77)-(82A)


<u>Compound</u>	<u>Ring F</u>	<u>CF₃</u>	<u>C-F</u>
(77)	-40.1	-	-
(78)	-37.2	-74.3	-184.8
(81A)	-	-73.7	-184.0
(79A)	-43.5	-78.1	-188.2
(82A)	-	-73.1	-183.8

The enhanced reactivity of the triazines (14) and (15) must be the reason for the formation of the *ortho* substituted products. All the *ortho* substituted products give complicated ^1H NMR spectra each having four aromatic protons. The ^{19}F NMR spectra of the *ortho* substituted compounds are similar to the spectra recorded for the *para* substituted compounds, as expected. ^1H and ^{19}F NMR are listed in the appendix.


5.7.5 Potential Use of Compounds (77)-(82A) as Non-Linear Optic Molecules.

Organic molecules with an electron donor group connected to an electron acceptor group via a conjugated π system may show charge transfer properties, i.e. electron density transferred from the donor part of the molecule to the acceptor part thus creating charge separation. There is a great amount of interest in using these types of molecules in non-linear optical devices¹⁶⁹. Their fluorescent properties have


been studied¹⁷⁰. It is beyond the scope of this thesis to consider the physics of these systems but the following molecules, which all have donor-acceptor structures, exhibit physical properties of interest to non-linear optic theory¹⁷¹.




We may consider the phenyl-s-triazines to have the following structure:-



This donor acceptor structure is plausible as fluorinated triazine rings have been shown to be capable of supporting a negative charge in the formation of stable σ complexes¹⁷².



In computer modelling studies (81A) was found to be non-planar, the benzene ring being at an angle to the triazine ring, as shown below (Fig 13).


Figure 13**Computer Generated Structure of (81)**

This twisted geometry may give rise to unusual fluorescent properties as seen in molecules of a similar structure. Indeed, the fluorescent properties of 2-(N,N-diethylamino)-4,6-dichloro-s-triazine have been studied¹⁷⁰.

Computer calculations suggest that molecules of the type (77)-(82A) do exhibit non-linear optical behaviour but are no better than more readily available molecules.

5.8 Reaction of 1,8-Bis(dimethylamino)naphthalene with Trifluoro-s-triazine


1,8-Bis(dimethylamino)naphthalene reacts with trifluoro-s-triazine to produce a tetrasubstituted naphthalene derivative (83) as red crystals in 53% yield.



^1H NMR of the product (83) is similar to those of the annelation products (4) and (44) as discussed in Chapter 4, proving the tetrasubstituted naphthalene structure. Elemental analysis and mass spectra were obtained consistent with the assigned structure.

5.9 Reactions of Fluorinated -s-triazines with N-Ethylaniline

Reactions between trifluoro-s-triazine and secondary amines are well known as outlined in section 5.2. Reactions between N-ethylaniline and the triazines (13) and (14) were performed to ascertain whether any carbon substituted products were formed. We found that only the nitrogen substituted products (84) and (80) were produced in 69 and 89% yield respectively.



(84) R = F

(80) R = $(\text{CF}_3)_2\text{CF}$


The position of substitution of the triazine ring is proved by the ^1H NMR spectra. No N-H resonances are seen indicating that the N-H proton has been substituted. Also, five protons are seen in the aromatic region rather than four protons forming an AB system which we would expect for a carbon-substituted product.

^{19}F NMR, mass spectra, UV spectra, IR spectra and elemental analyses were obtained for (84) and (80) in accordance with the assigned structures.

We may conclude that N-H is a stronger nucleophile than the activated aromatic ring acting as a carbon nucleophile.

5.10 Reaction of Trifluoro-s-triazine with 2-(N,N-Dimethylamino)-Methoxybenzene

Trifluoro-s-triazine reacts with 2-(N,N-dimethylamino)-methoxybenzene to produce the nitrogen substituted product (85) in 31% yield.




The structure is proved by the ^1H NMR spectrum. Four protons are seen in the aromatic region and there are two singlets at 3.47 and 3.84ppm each having a relative intensity of three indicating that one methyl group on the nitrogen atom has been substituted.

^{19}F NMR, mass spectra, UV spectra, IR spectra and elemental analyses were obtained for (85) in accordance with the assigned structure.

This result indicates that the 2-methoxy group prevents the NMe_2 group from conjugating with the ring thus decreasing activation towards electrophilic attack on the ring, in agreement with Shaws' results in reactions between 2-toluidine and cyanuric chloride¹⁵¹ (section 5.3.1).

5.11 Reaction of Trifluoro-s-triazine with N,N,N',N'-Tetramethyl-1,4-Diaminobenzene

Trifluoro-s-triazine reacts with N,N,N',N'-tetramethyl-1,4-diaminobenzene to produce the nitrogen substituted product (86) in 15% yield.




The structure of (86) is proved by the ^1H NMR spectrum. Two singlets at 2.97 and 3.50ppm with relative intensities 6 and 3 respectively indicate that one of the methyl groups has been substituted. Also, the AB system (6.78 and 7.17ppm ($J_{\text{AB}}=8.9\text{Hz}$)) indicative of a *para*-substituted benzene ring remains.

^{19}F NMR, mass spectra, UV spectra, IR spectra and elemental analyses were obtained for (86) in accordance with the assigned structure.

The nitrogen-substituted product is obtained as the 4-position is blocked. No *ortho* substituted products were obtained, in agreement with Shaws' results in the reaction between 2-toluidine and cyanuric chloride¹⁵¹ (section 5.3.1).

5.12 Reaction between 2-(N,N-Dimethylamino)-pyridine and Perfluorodi-isopropyl-s-triazine

2-(N,N-Dimethylamino)-pyridine reacts with perfluorodi-isopropyl-s-triazine to give the pyridinium hydroxide salt (87) after the reaction was washed with water.



No carbon substituted products were obtained in agreement with Chakrabarti's results¹⁶⁶.

The ^1H NMR spectrum shows a singlet at 3.25ppm corresponding to the NMe_2 group and four protons in the aromatic region. Elemental analysis is consistent with a molecular formula of $\text{C}_{16}\text{H}_{11}\text{N}_5\text{F}_{14}\text{O}$.

5.13 Summary

We have shown that pyrroles, indoles and tertiary anilines can act as carbon nucleophiles in reactions with perfluorinated heterocycles in electrophilic substitution reactions.

Pyrroles are substituted at the 2-position, as proved by ^1H NMR coupling constants, whereas indoles undergo substitution at the 3-position, proved by ^1H and ^{13}C NMR, which is in accordance with the literature¹⁶⁷.

Tertiary anilines react via the para carbon atom to form electrophilic substitution products. The reaction is complicated by the formation of the ortho species when the nucleophile is stronger.

EXPERIMENTAL SECTION

INSTRUMENTATION

Gas liquid Chromatography (g.c.) analysis was carried out on a Hewlett Packard 5890A gas chromatograph fitted with a 25m cross-linked methyl silicone capillary column. Preparative g.c. was performed on a Varian Aerograph Model 920 (catharometer detector) gas chromatograph.

Fractional distillation of product mixtures was carried out using a Fischer Spahlrohr MMS 255 small concentric tube apparatus. Boiling points were recorded during distillation. Melting points were carried out at atmospheric pressure and are uncorrected.

Carbon, hydrogen and nitrogen elemental analyses were obtained using a Perkin-Elmer 240 Elemental Analyser or a Carlo Erba 1106 Elemental Analyser. Analysis for halogens were performed as described in the literature.

Ultraviolet spectra were recorded on a Perkin-Elmer Lambda 2 or a Pye-Unicam PU 8720 UV/Vis spectrophotometer.

Infra Red spectra were recorded on either a Perkin-Elmer 457 or 577 Grating Spectrophotometer using conventional techniques.

Proton NMR spectra were recorded on a Hitachi Perkin-Elmer R-24B (60MHz), a Bruker AC250 (250MHz) and a Varian VXR400S(400MHz) NMR spectrometer.

Fluorine NMR spectra were recorded on a Varian EM360I (56.45MHz), a Bruker AC250 (235MHz) and a Varian VXR400S (365MHz) NMR spectrometer.

Carbon NMR were recorded on a Varian VXR400S (100MHz) NMR spectrometer.

Mass Spectra of solid samples were recorded on a VG 7070E spectrometer. G.c. mass spectra were recorded on the VG 7070E spectrometer linked to a Hewlett Packard 5790A gas chromatograph fitted with a 25m cross-linked methyl silicone capillary column.

REAGENTS

In general chemicals were used as received from suppliers (Aldrich, Lancaster, Fluka) and solvents were dried by literature procedures.

CHAPTER SIX

EXPERIMENTAL TO CHAPTER TWO

6.1 Preparation of HF/Ether Solution

Anhydrous Hydrogen Fluoride gas was bubbled through dry diethyl ether (100ml), under anhydrous conditions, which was contained in an ice cooled FEP bottle. Face masks, gloves and an efficient fume hood are essential when handling anhydrous HF. The concentration of HF was determined by titrating an aliquot of the ether solution with a standard solution of sodium hydroxide using phenolphthalein as the indicator. e.g. a 1ml aliquot of HF/ether solution required 5.65ml of 1.192M NaOH solution giving a 0.134g_{HF}/ml solution.

6.2 Preparation of 1,8-Bis(dimethylamino)naphthalene Hydrogen Fluoride Complex (PS/HF)

1,8-Bis(dimethylamino)naphthalene (Proton Sponge) (1.5g, 7mmol) was dissolved in the minimum amount of dry diethyl ether and the required 1:1 stoichiometric amount of the HF/ether solution (0.134g_{HF}/ml) (1.05ml, 7mmol) was added by pipette. A white solid immediately precipitated and the ether was carefully removed under reduced pressure to leave 1,8-Bis(dimethylamino)-naphthalene hydrofluoride (PS/HF) (1) (1.62g, 98%); m.p. 117-118°C; (Found: C, 69.7; H, 8.5; N, 11.5. C₁₄H₁₉N₂F requires C, 71.8; H, 8.1; N, 11.95%); i.r. spectrum 1; n.m.r. spectrum 1; mass spectrum 1.

The preparation was repeated many times as the material was required usually on the 3g scale, but was scaled up to 25g without any problems. In all the following reactions 1,8-Bis(dimethylamino)-naphthalene hydrogen fluoride complex will be referred to as PS/HF.

6.3 Reactions using PS/HF as the Catalyst in C-C Bond Forming Reactions

6.3.1 Attempted Oligomerisation of Tetrafluoroethylene

This experiment was carried out at the ICI Experimental Site, Widnes with Dr. J. Hutchinson and Mr. M. Martin on 30/1/90.

A 0.5l autoclave was charged with PS/HF (23g, 98mmol) in dimethylformamide and α -Pinene (2 drops) was added (to inhibit any free radical reactions). The autoclave was flushed out with dry nitrogen four times and then with

tetrafluoroethylene three times. Then tetrafluoroethylene was added up to a maximum pressure of 10 bar and a maximum temperature of 110°C. No pressure drop was observed on the manometer. Consequently, on cooling, the autoclave was opened to reveal a red solvent layer and no lower fluorocarbon layer.

6.3.2 Dimerisation of Hexafluoropropene

A Carius tube was charged with PS/HF (1.63g, 7mmol) and acetonitrile (35ml), and hexafluoropropene (7.34g, 49mmol) was transferred under vacuum to the tube which was cooled in liquid air. The tube was sealed and allowed to warm to room temperature in a steel casing and then agitated on a rotating arm for 48 hr at room temperature. The tube was opened to reveal a lower fluorocarbon layer which was collected and determined to be the thermodynamic dimer of hexafluoropropene (2)(5.3g, 72%) as the only product by GC; δ_F (60MHz, CFCl_3) -60ppm (m, 3F, CF_3), -63 (m, 3F, CF_3), -86 (m, 3F, CF_3), -100.5 (s(br), 1F, CF), -119 (m, 2F, CF_2); m/z (EI^+) 281 ($\text{M}^+ - \text{F}$, 24%); as compared to the literature data¹²¹ [$\delta_F = -60.2$ ppm (m, 3F, CF_3), -62.8 (m, 3F, CF_3), -86.4 (m, 3F, CF_3), -100.1 (s (br), 1F, CF), -119.6 (m, 2F, CF_2); m/z (EI^+) 281 ($\text{M}^+ - \text{F}$, 27%)].

6.3.3 Attempted Oligomerisation of Perfluorocyclobutene

A Carius tube was charged with PS/HF (1.07g, 5mmol) and acetonitrile (30ml), and perfluorocyclobutene (12.90g, 79mmol) was transferred under vacuum to the tube which was cooled in liquid air. The tube was sealed and allowed to warm to room temperature in a steel casing and then agitated on a rotating arm for 48 hr at room temperature. All volatiles were transferred under vacuum to a trap and analysis by GC/MS and ^{19}F n.m.r. showed no evidence for perfluorocyclobutene oligomers.

The reaction was repeated at 50°C in acetonitrile and using sulpholane as the solvent but no oligomers of perfluorocyclobutene were isolated.

6.3.4 Attempted Formation of a Stable Perfluorinated Carbanion

A flask was charged with hexafluoropropene dimer (2)(1.8g, 6mmol), PS/HF (1.4g, 6mmol) and sulpholane (5ml) under a plume of dry nitrogen and the mixture was stirred for one week at room temperature. ^{19}F NMR of the mixture revealed only start material. The reaction was repeated in tetraglyme and acetonitrile but only start material was observed by ^{19}F NMR.

6.3.5 Perfluoroalkylation of Pentafluoropyridine

A flask was charged with PS/HF (0.75g, 3mmol), pentafluoropyridine (6)(2.20g, 13mmol) and dry sulpholane (20ml) and then cooled in liquid air and evacuated. After warming to room temperature, hexafluoropropene (3.4g, 23mmol) was added via an expandable gas reservoir and the mixture was stirred vigorously for 7 days at room temperature. The volatile products were removed from the reaction mixture by flash distillation (4.8g). GC/MS showed three main products, which were separated using an SE 30 column at 100°C and identified as hexafluoropropene dimer (2)(2%); pentafluoropyridine (6)(49%); perfluoro-4-isopropylpyridine (7)(44%); δ_F (235 MHz, CD₃CN, CFCI₃) -75.4 (6F, s, CF₃), -86.7 (2F, s, 2,6-ring F), -135.3, 137.5 (2F, d, 3,5-ring F), -180.5 (1F, m, C-F); m/z (EI⁺) 319 (M⁺, 41%), 250 (15, M - CF₃), 200 (100, M - C₂F₅), as compared to the literature data¹²⁴ (δ_F = -74.3 (6F, s, 2CF₃), -87.3 (2F, s, 2,6-ring F), -135.1 (2F, s, 3,5-ring F), -178.5 (1F, s, CF); m/e 319 (M)). Trace amounts of perfluorodi-isopropylpyridine (8)(2%); m/z (EI⁺) 469 (M⁺, 18%); and, perfluorotri-isopropylpyridine (12)(1%); m/z (EI⁺) 619 (M⁺, 16%); were also observed by GC/MS.

6.3.6 Perfluoroalkylation of Tetrafluoropyrimidine

A flask was charged with PS/HF (1.40g, 6mmol), tetrafluoropyrimidine (9)(2.30g, 15mmol) and dry sulpholane (30ml) and then cooled in liquid air and evacuated. After warming to room temperature, hexafluoropropene (5.23g, 35mmol) was added via an expandable gas reservoir and the mixture was stirred vigorously for 3 days at room temperature. The volatile products were removed from the reaction mixture by flash distillation (6.1g). The product mixture was analysed by GC/MS and ¹⁹F n.m.r. and found to consist of perfluoro-4-isopropylpyrimidine (10)(27%); δ_F (235MHz, CD₃CN, CFCI₃) -48.7ppm (2-ring F), -72 (6-ring F), -76.1 (CF₃ groups), -154 (5-ring F), -186.9 (CF); m/z (EI⁺) 302 (M⁺, 35%), 283 (20, M - F), 233 (18, M - CF₃), 183 (51, M - C₂F₅); as compared to the literature data¹²⁵ [δ_F = -48.7ppm (2-ring F), -72.5 (6-ring F), -78.1 (CF₃), -154.5 (5-ring F), -188.7 (CF); m/e (EI⁺) 302 (M⁺, 65%), 283 (36, M-F), 233 (20, M-CF₃), 183 (57, M-C₂F₅)]; perfluoro-2,6-di-isopropylpyrimidine (11)(15%); δ_F (235MHz, CD₃CN, CFCI₃) -48.7ppm (2-ring F), -76.2 (CF₃), -133.5 (5-ring F), -186.9 (CF); m/e (EI⁺) 452 (M⁺, 22%), 433 (33, M - F), 383 (16, M - CF₃), 333 (40, M - C₂F₅); as compared to the literature data¹²⁵ [δ_F = -48.6 (2-ring F), -76.5 (CF₃), -132.5 (5-ring F), -186.3 (CF); m/e (EI⁺) 452 (M⁺, 42%), 433 (44, M-F), 383 (22, M-CF₃), 333 (37, M-C₂F₅)]; perfluoro-2,4,6-tri-isopropylpyrimidine (12)(39%); δ_F (235MHz, CD₃CN, CFCI₃) -76.2 (CF₃), -

124.1 (5-ring F), -182 (CF), -186.9 (CF); m/e (EI⁺) 602 (M⁺, 5%), 583 (9, M - F), 533 (4, M - CF₃), 452 (14, M - C₃F₆); as compared to the literature data¹²⁵ [δ_F = -76.5ppm (CF₃), -123.1 (5-ring F), -182.1 (CF), -186.3 (CF); m/e (EI⁺) 602 (M⁺, 46%), 583 (50, M-F), 533 (22, M-CF₃)].

6.3.7 Perfluoroalkylation of Trifluoro-s-triazine

A flask was charged with PS/HF (0.70g, 3mmol), trifluoro-s-triazine (13)(1.80g, 13mmol) and dry sulpholane (20ml) and then cooled in liquid air and evacuated. After warming to room temperature, hexafluoropropene (7.5g, 50mmol) was added via an expandable gas reservoir and the mixture was stirred vigorously for 3 days at room temperature. The volatile products were removed from the reaction mixture by flash distillation (8.2g). The product mixture was analysed by ¹⁹F n.m.r. and GC/MS and found to consist of two main products; perfluoroisopropyl-s-triazine (14)(35% by GC); m/z (EI⁺) 285 (M⁺, 100%), 266 (98, M - F), 197 (88, M - FCF₃), 166 (51, M - C₂F₅); perfluorodi-isopropyl-s-triazine (15)(8%); m/z (EI⁺) 435 (M⁺, 65%), 416 (100, M - F), 347 (38, M - FCF₃), 316 (11, M - C₂F₅). ¹⁹F n.m.r. of the product mixture revealed perfluoroalkylation; δ_F (235 MHz, CD₃CN, CFCl₃) -33ppm (N=C-F), -73 (CF₃), -183.8 (CF); as compared to the literature data¹²⁶ [δ_F = -30.4-30.6ppm (ring F), -74.4-75.4 (CF₃ groups), -183.8-186.5 (CF)].

6.4 Reactions Using PS/HF in C-F Bond Forming Reactions

6.4.1 With Benzoyl Chloride

A mixture containing PS/HF (1.07g, 4.5mmol), benzoyl chloride (0.70g, 5mmol) and acetonitrile was allowed to stand at room temperature for 24 hr. A white solid, 1,8-bis(dimethylamino)naphthalene hydrochloride (PS/HCl), precipitated. The mixture was filtered and benzo trifluoride (0.17g, 1.16mmol) was added as an nmr marker. ¹⁹F n.m.r. revealed benzoyl fluoride (76% yield by integration); δ_F (235 MHz, CH₃CN, CFCl₃) +17.0 (1F, s, CO-F); as compared to the literature data (δ_F = +17.1ppm)¹⁷³.

6.4.2 Reaction between Hexafluoroacetone and PS/HF

A Carius tube was charged with PS/HF (5.1g, 22mmol) in acetonitrile (20ml) and hexafluoroacetone (5.3g, 32mmol) was transferred under vacuum to the trap which was cooled in liquid air. The tube was sealed and allowed to stand at room temperature overnight. The solution went pale yellow. The tube was opened and ¹⁹F

n.m.r. revealed the desired carbinolate species (17); δ_F (235 MHz, CH_3CN , CFCl_3) -80.33 (6F, s, CF_3), -107.88 (1F, s(br), CF). There was no peak at -164ppm indicating that all the PS/HF had reacted, hence the solution contained 22mmol of the carbinolate species. The reaction solution was used in the two following trapping reactions.

6.4.2.1 Preparation of Heptafluoroisopropyl benzoate (18)

Benzoyl chloride (1.7g, 12mmol) was added dropwise to the carbinolate solution prepared above (10ml, 11mmol) and the mixture was stirred overnight. The volatiles were transferred under vacuum and analysed by GC/MS and ^{19}F n.m.r. There were two products; benzoyl fluoride (51%); and, heptafluoroisopropyl benzoate (18) (49%) which was isolated by preparative GC using a 30% SE 30 column at 150°C; δ_F (235 MHz, CH_3CN , CFCl_3) -78.1 (6F, s, CF_3), -141.1 (1F, s, CF); m/z (EI⁺) 290 (M^+ , 22%), 105 (100, Ar-C=O), 77 (60, C_6H_5), 69 (11, CF_3).

6.4.2.2 Preparation of Heptafluoroisopropyl benzyl ether (19)

Benzyl bromide (2.3g, 13mmol) was added dropwise to the carbinolate solution prepared above (10ml, 11mmol) and the mixture was stirred overnight at room temperature. All volatile materials were transferred under vacuum and analysed by GC/MS and ^{19}F n.m.r. There were two components; benzyl bromide (39%) and, heptafluoroisopropyl benzyl ether (19)(61%) which was isolated by preparative scale GC using a 30% SE 30 column at 150°C; ν_{max} 1230 cm^{-1} (C-O-C stretch); δ_H (250 MHz, CDCl_3 , TMS) 5.01 (2H, s, CH_2), 7.42 (5H, m, Ar-H); δ_F (235 MHz, CDCl_3 , CFCl_3) -79.5 (6F, s, CF_3), -142.6 (1F, s, CF); m/z (EI⁺) 276 (M^+ , 33%), 91 (100, Ar- CH_2); m/z (EI⁺) 276 (M^+ , 33%), 91 (100, Ar- CH_2).

6.4.3 With 2,4-Dinitrochlorobenzene

A mixture containing PS/HF (3.06g, 13mmol) and 2,4-dinitrochlorobenzene (2.69g, 13mmol) was refluxed in acetonitrile for 2 days. Benzotrifluoride was added as an nmr marker and ^{19}F nmr revealed 2,4-dinitrofluorobenzene (45% yield by integration) at -109ppm, as compared to the literature data (δ_F = -107.7ppm)¹⁷³.

6.4.4 With Benzyl Bromide

A mixture containing PS/HF (1.87g, 8mmol), benzyl bromide (1.30g, 8mmol) and acetonitrile ((15ml) was refluxed for 24 hr. A white solid, 1,8-bis(dimethylamino)naphthalene hydrobromide (PS/HBr), precipitated (i.r.

spectrum). The mixture was filtered and benzo trifluoride (0.19g, 1.30mmol) was added as an nmr marker. ^{19}F n.m.r. revealed benzyl fluoride (72% by integration); δ_{F} (235 MHz, CH_3CN , CFCl_3) -206.4 (1F, t, $J_{\text{HF}}=49$ Hz, $-\text{CH}_2\text{F}$); as compared to the literature data ($\delta_{\text{F}}=-207\text{ppm}$)¹⁷³.

6.4.5 With Octyl Iodide

A mixture containing PS/HF (1.10g, 4.7mmol), octyl iodide (0.91g, 3.8mmol) and acetonitrile was heated at reflux for 24 hr. On cooling white crystals, 1,8-bis(dimethylamino)naphthalene hydriodide, precipitated (i.r.spectrum). The mixture was filtered and benzo trifluoride (0.23g, 1.57mmol) was added as an nmr marker. ^{19}F n.m.r. revealed octyl fluoride (65% yield by integration); δ_{F} (235 MHz, CH_3CN , CFCl_3) 218.0 (1F, t, $J_{\text{HF}}=42$ Hz, $-\text{CH}_2\text{F}$); as compared to the literature data ($\delta_{\text{F}}= -219\text{ppm}$)¹⁷³.

6.4.6 Attempted Reaction Between PS/HF and 1,2-Epoxybutane

A mixture containing PS/HF (1.54g, 6.6mmol), 1,2-epoxybutane (0.43g, 6.0mmol) and acetonitrile (15ml) was refluxed for 48 hr. ^{19}F n.m.r. of the mixture showed that no reaction had taken place. The reaction was repeated in sulpholane at 130°C for 3 days with no hydrofluorination taking place, by ^{19}F n.m.r.

6.4.7 Preparation of 1,2:5,6-Di-O-isopropylidene-3-O-triflic- α -D-gulofuranose (21)

Diacetone-D-glucose (20)(1.83g, 7mmol) and pyridine (2.5g, 32mmol) were dissolved in dichloromethane (50ml) and the solution, under dry nitrogen, was cooled down to 0°C with an ice/salt bath. Triflic anhydride (5.1g, 18mmol) was added dropwise, with the temperature of the reaction maintained under 5°C. The solution was stirred at 5°C for 30 mins. A white solid was deposited and the solution went pale yellow. The solution was washed sequentially with ice-cold dilute hydrochloric acid and water. The organic layer was dried (MgSO_4) and evaporated to leave a solid, the desired triflate (21)(2.17g, 79%); δ_{F} (60 MHz, CDCl_3 , CFCl_3) -76.0 (3F, s, CF_3); as compared to the literature data¹²⁹; and was used immediately in the next reaction.

6.4.8 Attempted Reaction Between PS/HF and Triflate (21)

A mixture containing triflate (21)(2.17g, 5.5mmol), PS/HF (1.80g, 7.7mmol) and acetonitrile (20ml) was refluxed overnight. The solution went dark brown and a solid precipitated. This solid was collected by filtration and recrystallised

from water as white needles and found to be the salt, 1,8-bis(dimethylamino)-naphthalene hydrotriflate (23); m.p. 215-218°C; (Found: C, 49.7; H, 5.3; N, 7.6. Calc for C₁₅H₁₉N₂F₃SO₃: C, 49.5; H, 5.2; N, 7.6%); i.r. spectrum recorded; m/z (EI⁺) 214 (naphthalene ion, 40%). ¹⁹F n.m.r. of the remaining reaction solution showed no evidence for a fluorinated glucose. The triflate salt may have been produced after decomposition of the triflated glucose.

The reaction was repeated at room temperature but no fluorinated glucose was observed, by ¹⁹F n.m.r.

6.4.9 Preparation of 1,2:5,6-Di-O-isopropylidene-3-O-toluene-p-sulphonyl- α -D-allofuranose (22)

Diacetone-D-glucose (20)(3.00g, 11.5mmol) was dissolved in pyridine (40ml) and cooled to 0°C. A solution of tosyl chloride (7.00g, 36.7mmol) in pyridine (20ml) was added dropwise. The solution was stirred at room temperature for two days. Water (4ml) was added and the solution was left to stand for a further 20 mins. It was then poured onto ice/water (300ml) and the crude sulphonate was filtered off and recrystallised from aqueous ethanol to yield the desired tosylate (22)(2.50g, 52%); m.p. 120-122°C (lit.¹³⁰, 122-123°C); (Found: C, 54.5; H, 6.2. Calc for C₁₉H₂₆SO₈: C, 55.1; H, 6.3%); IR spectrum recorded; δ_{H} (235 MHz, CDCl₃, TMS) 1.15, 1.19, 1.31 and 1.48 (3H, s, acetal groups), 2.46 (3H, s, Ar-CH₃), 3.89-4.06 (4H, m, unassigned 2CH, CH₂), 4.78 (1H, m, H-1), 4.83 (1H, d, J_{3,4}=3.6 Hz, H-4), 5.93 (1H, d, J_{1,2}=3.6 Hz, H-1), 7.34 and 7.83 (4H, AB, J_{AB}=8.3 Hz, Ar-H); as compared to the literature data¹³⁰.

6.4.10 Attempted Reaction Between PS/HF and Tosylate (22)

A mixture containing tosylate (22)(0.80g, 1.9mmol), PS/HF (0.70g, 3mmol) and acetonitrile (5ml) was refluxed overnight. ¹⁹F n.m.r. of the reaction mixture showed unreacted PS/HF and no evidence for a fluorinated glucose.

CHAPTER SEVEN

Experimental to Chapter Three

7.1 Preparation and Reactions of Trialkylamine Hydrofluoride Complexes

7.1.1 Preparation of Trialkylamine Hydrofluoride Complexes (24)-(28)

All bases were used as supplied (Aldrich). The hydrofluoride salts were prepared by adding a stoichiometric amount of a calibrated HF/ether solution to an ethereal solution of the base, followed by evaporation of the solvent to leave the salt, as described previously (Section 6.1).

The following salts were prepared:-

1) Triethylamine Hydrofluoride (24); hygroscopic solid at r.t; i.r. spectrum 2; n.m.r. spectrum 2; mass spectrum 2.

2) Tributylamine Hydrofluoride (25); liquid; i.r. spectrum 3; n.m.r. spectrum 3; mass spectrum 3.

3) Trihexylamine Hydrofluoride (26); liquid; i.r. spectrum 4; n.m.r. spectrum 4; mass spectrum 4.

4) Trioctylamine Hydrofluoride (27); liquid; i.r. spectrum 5; n.m.r. spectrum 5; mass spectrum 5.

5) Tridodecylamine Hydrofluoride (28); liquid; i.r. spectrum 6; n.m.r. spectrum 6; mass spectrum 6.

7.1.2 Methodology for Standard Reactions

Three standard experiments were performed using each amine.HF complex as the source of soluble fluoride ion. The experiments were chosen to provide a range of fluoride ion reactions. i.e. nucleophilic substitution reactions at unsaturated, saturated and aromatic carbon positions. The same methodology was used for each hydrofluoride salt, so general procedures for the three standard reactions appear in this section. The quantities of each reagent used can be found in the tables listed under the appropriate

section. Any deviation from these procedures are listed under the appropriate sections; e.g. solubility of HCl salt formed in the reactions using the piperidine HF salts. All yields quoted are n.m.r. yields of the crude reaction mixtures, not isolated yields. We chose not to work-up the reactions due to their small scale and the fact that crude n.m.r. yields give the true maximum yield as there are no handling losses associated with work-up.

7.1.2.1 Reaction of Base Hydrofluoride Complexes with Benzoyl Chloride

A mixture containing Base/HF, benzoyl chloride and acetonitrile (10ml) was allowed to stand at room temperature for 24 hr. A white solid, the corresponding Base/HCl salt, was precipitated. Benzotrifluoride was added to the reaction mixture which was shaken to ensure homogeneity. The mass, and hence the number of moles of benzotrifluoride provided a marker for n.m.r. integration. ^{19}F n.m.r. of the reaction mixture was recorded; δ_{F} (235 MHz, CH_3CN , CFCl_3); to reveal a peak at +16.7 ppm due to benzoyl fluoride, as compared to the literature data¹⁷³; at -63ppm due to the benzotrifluoride marker and between -150 and -170 ppm due to unreacted Base/HF. The yield of benzoyl fluoride was calculated by comparing the integration of the peak due to the benzotrifluoride marker and that due to benzoyl fluoride.

7.1.2.2 Reaction of Base Hydrofluoride Complexes with Benzyl Bromide

A mixture containing Base/HF, benzyl bromide and acetonitrile (15ml) was refluxed for 24 hr. On cooling, a white solid, the corresponding Base/HBr salt precipitated. Benzotrifluoride was added to the reaction mixture which was shaken to ensure homogeneity. The mass, and hence the number of moles of benzotrifluoride provided a marker for n.m.r. integration. ^{19}F n.m.r. of the reaction mixture was recorded; δ_{F} (235 MHz, CH_3CN , CFCl_3); to reveal a peak at -63 ppm due to the benzotrifluoride marker; a peak between -150 and -170 ppm due to unreacted Base/HF and a peak at -206.4 (t, $J_{\text{HF}}=49\text{Hz}$) due to benzyl fluoride, as compared to the literature data¹⁷³. The yield of benzyl fluoride was calculated by comparing the integration of the peak due to the benzotrifluoride marker and that due to benzyl fluoride.

7.1.2.3 Reaction of Base Hydrofluoride Complexes and 2,4-Dinitrochlorobenzene

A mixture containing Base/HF, 2,4-dinitrochlorobenzene and acetonitrile was refluxed for 48 hr. A white solid, the corresponding Base/HCl salt, was precipitated. Benzotrifluoride was added to the reaction mixture which was shaken to ensure

homogeneity. The mass, and hence the number of moles of benzotrifluoride provided a marker for n.m.r. integration. ^{19}F n.m.r. of the reaction mixture was recorded; δ_{F} (235 MHz, CH_3CN , CFCl_3); to reveal a peak at -63 ppm due to the benzotrifluoride marker; at -108.8 ppm due to 2,4-dinitrofluorobenzene, as compared to the literature data¹⁷³ and a peak between -150 and -170 ppm due to unreacted Base/HF. The yield of 2,4-dinitrofluorobenzene was calculated by comparing the integration of the peak due to the benzotrifluoride marker and that due to dinitrofluorobenzene.

7.1.3 Reactions of Trialkylamine.HF Complexes

The three standard reactions were performed using the trialkylamine.HF complexes as the source of Fluoride ion. See above for details of the methodology. Hence, in the following tables the HF salt used was $\text{R}_3\text{N.HF}$, where R = alkyl.

7.1.3.1 Reaction with Benzoyl Chloride

<u>R</u>	<u>HF salt</u> g, mmol	<u>Benzoyl Chloride</u> g, mmol	<u>Benzotrifluoride</u> g, mmol	<u>Yield</u> %
Et	0.89, 7.35	0.86, 6.10	0.20, 1.37	91
But	1.01, 4.93	0.62, 4.41	0.17, 1.16	88
Hex	1.29, 4.46	0.54, 3.84	0.29, 1.99	90
Oct	2.32, 6.22	0.80, 5.69	0.28, 1.92	68
Dodec	0.62, 1.15	0.28, 1.98	0.19, 1.35	79

7.1.3.2 Reaction with Benzyl Bromide

<u>R</u>	<u>HF salt</u> g, mmol	<u>Benzyl Bromide</u> g, mmol	<u>Benzotrifluoride</u> g, mmol	<u>Yield</u> %
Et	0.75, 6.20	0.86, 5.03	0.25, 1.71	18
But	0.90, 4.39	0.72, 4.21	0.25, 1.71	12
Hex	1.91, 6.61	1.01, 5.91	0.29, 1.98	17
Oct	2.22, 5.94	0.95, 5.55	0.23, 1.57	14
Dodec	1.90, 3.50	0.53, 3.09	0.24, 1.64	11

7.1.3.3 Reaction with 2,4-Dinitrochlorobenzene

B	HF salt	DNCBenzene	Benzotrifluoride	Yield
	g, mmol	g, mmol	g, mmol	%
Et	0.39, 3.22	0.52, 2.57	0.22, 1.50	34
But	0.72, 3.51	0.63, 3.11	0.33, 2.26	84
Hex	1.81, 6.26	1.06, 5.23	0.20, 1.37	77
Oct	1.54, 4.13	0.75, 3.70	0.33, 2.26	61
Dodec	1.02, 1.88	0.36, 1.78	0.21, 1.44	14

7.2 Preparation and Reactions of Polysubstituted Piperidine Hydrofluoride Complexes**7.2.1 Preparation of Pentasubstituted Piperidine Bases**

Tetramethylpiperidine (31) and pentamethylpiperidine (29) were used as supplied (Aldrich). The three other piperidine bases were prepared as follows:-

7.2.1.1 Preparation of N-Ethyl-2,2,6,6-Tetramethylpiperidine (32)^{1 3 2}

A mixture containing tetramethylpiperidine (31)(24.7g, 0.17mol) and ethyl p-toluene sulphonate (18.0g, 0.09mol) was heated at 100°C for 24 hr in a flask fitted with an air condenser. The mixture solidified into a partly browned cake. On cooling, the product mixture was washed thoroughly with diethyl ether to precipitate a white solid which was collected by filtration and identified as the salt, tetramethylpiperidinium tosylate; m.p. 218-222°C (lit.,¹³² 219-221°C); IR spectrum recorded; m/z (EI⁺)142 (M⁺ piperidinium cation, 6.6%). The ether layer was dried and evaporated and the residue was distilled on the Fischer Spahlrohr to yield N-Ethyl-2,2,6,6-tetramethylpiperidine (32)(6.38g, 21%); b.p. 95.6-96°C/24mmHg; (Found: C, 78.1; H, 14.2; N, 8.4. Calc for C₁₁H₂₃N: C, 78.1; H, 13.6; N, 8.4%); IR spectrum recorded; δ_{H} (235MHz, CDCl₃, TMS) 1.28 (15H, m, CH₃), 1.62 (4H, m, C-CH₂-N), 1.75 (2H, m, C-CH₂-C), 2.72 (2H, q, J=7.1 Hz, CH₂); m/z (EI⁺) 168 (M⁺, 3.8%), 154 (M⁺-Me group).

7.2.1.2 Preparation of N-Allyl-2,2,6,6-Tetramethylpiperidine (33)^{1 3 2}

A mixture containing tetramethylpiperidine (31)(30.7g, 0.22mol) and allyl bromide (13.2g, 0.11mol) was heated at 50°C for 3 days. The mixture turned pale yellow. The mixture was washed thoroughly with diethyl ether and filtered. The ether layer was dried and evaporated and the residue was distilled on the Fischer Spahlrohr to yield N-allyl-2,2,6,6-tetramethylpiperidine (33)(9.3g, 24%); b.p. 103-109°C/20mm Hg (pure by GC); (Found: C, 79.5; H, 13.1; N, 7.9. Calc for C₁₂H₂₃N: C, 79.5; H, 12.7; N, 7.7%); IR spectrum recorded; δ_{H} (60MHz, CDCl₃, TMS) 1.1 (12H, s, CH₃), 1.5 (6H, s, ring CH₂), 3.2 (2H, m, N-CH₂-Ar), 5.1 (2H, m, C=CH₂), 5.9 (1H, m, -CH=); m/z (EI⁺) 181 (M⁺, 9.9%), 166 (100%, M⁺-Me group).

7.2.1.3 Preparation of N-Benzyl-2,2,6,6-Tetramethylpiperidine (34)^{1 3 2}

Benzyl bromide (12.2g, 71mmol) was added to tetramethylpiperidine (31)(26.6g, 0.19mol) over 20 mins at 60°C with stirring. The solution was heated at 100°C for 8 hr. On cooling, the reaction mixture was washed thoroughly with diethyl ether and filtered. The ether layer was dried and evaporated and the residue was distilled on the Fischer Spahlrohr to yield N-benzyl-2,2,6,6-tetramethylpiperidine (34)(5.4g, 32%); b.p. 135°C/6mm Hg; m.p. 30-32°C; (Found: C, 83.7; H, 11.1; N, 6.1. Calc for C₁₆H₂₅N: C, 83.1; H, 10.8; N, 6.1%); IR spectrum recorded; δ_{H} (60MHz, CDCl₃, TMS) 0.97 (12H, s, CH₃), 1.5 (6H, m, ring CH₂), 3.75 (2H, s, N-CH₂-Ar), 7.25 (5H, m, Ar-H); m/z (EI⁺) 231 (M⁺, 3.3%), 216 (100%, M⁺-Me group).

7.2.2 Preparation of the Hydrofluoride Complexes (35)-(39)

The hydrofluoride salts were prepared by adding a stoichiometric amount of a calibrated HF/ether solution to an ethereal solution of the base, followed by evaporation of the solvent to leave the salt, as described previously (Section 6.1).

The following salts were prepared:-

1) 2,2,6,6-tetramethylpiperidine Hydrofluoride (35); m.p. 76-80°C; (Found: C, 66.5; H, 12.4; N, 7.6. C₉H₂₀NF requires C, 67.1; H, 12.4; N, 8.7%); i.r. spectrum 7; n.m.r. spectrum 7; mass spectrum 7.

- 2) 1,2,2,6,6-pentamethylpiperidine Hydrofluoride (36); m.p. 122-124°C; i.r. spectrum 8; n.m.r. spectrum 8; mass spectrum 8.
- 3) N-ethyl-2,2,6,6-tetramethylpiperidine Hydrofluoride (37); m.p. 74-76°C; i.r. spectrum 9; n.m.r. spectrum 9; mass spectrum 9.
- 4) N-allyl-2,2,6,6-tetramethylpiperidine Hydrofluoride (38); m.p. 72-74°C; i.r. spectrum 10; n.m.r. spectrum 10; mass spectrum 10.
- 5) N-benzyl-2,2,6,6-tetramethylpiperidine Hydrofluoride (39); m.p. 83-86°C; i.r. spectrum 11; n.m.r. spectrum 11; mass spectrum 11.

7.2.3 Reactions of Piperidine.HF Complexes (35)-(39)

The same standard reactions were performed using the piperidine.HF salts as the source of Fluoride ion, under the same conditions as those carried out previously with other salts . See Section 7.1.2 for the methodology. However, in these reactions the hydrochloride/hydrobromide salt produced as a side product does not precipitate but remains in solution. In the following tables R refers to the group attached to the nitrogen atom in the piperidine.HF complex.

7.2.3.1 Reaction with Benzoyl Chloride

<u>R</u>	<u>HF salt</u> g, mmol	<u>Benzoylchloride</u> g, mmol	<u>Benzotrifluoride</u> g, mmol	<u>Yield</u> %
H	0.81, 5.0	0.57, 4.0	0.14, 0.9	83
Me	0.43, 2.46	0.32, 2.28	0.25, 1.71	64
Et	1.05, 5.55	0.70, 4.99	0.22, 1.51	78
Allyl	0.06, 0.29	0.27, 1.95	0.25, 1.71	55
Benzyl	0.49, 1.95	0.27, 1.95	0.17, 1.16	42

7.2.3.2 Reaction with Benzyl Bromide

R	HF salt g, mmol	Benzylbromide g, mmol	Benzotrifluoride g, mmol	Yield %
H	1.35, 8.40	1.40, 8.20	0.14, 0.8	25
Me	0.95, 5.43	0.84, 4.91	0.31, 2.12	84
Et	0.58, 3.07	0.49, 2.86	0.27, 1.85	38
Allyl	0.45, 2.24	0.27, 1.58	0.21, 1.44	n.r.
Benzyl	0.27, 2.08	0.18, 1.05	0.11, 0.64	n.r.

7.2.3.3 Reaction with 2,4-Dinitrochlorobenzene

R	HF salt g, mmol	DNCBenzene g, mmol	Benzotrifluoride g, mmol	Yield %
H	1.12, 6.90	1.31, 6.40	0.44, 3.0	49
Me	0.57, 3.26	0.63, 3.11	0.25, 1.71	78
Et	1.08, 5.71	0.80, 3.95	0.42, 2.88	94
Allyl	0.37, 1.84	0.36, 1.78	0.15, 1.03	31
Benzyl	0.31, 1.23	0.28, 1.38	0.18, 1.23	23

7.3 Preparation and Reactions of Tetramethylguanidine HF Complex

The tetramethylguanidine hydrogen fluoride complex (40) was prepared in the same manner as for the preparation of PS/HF (section 6.1).

Tetramethylguanidine Hydrogen Fluoride Complex (40); m.p. 66-68°C; (Found: C, 41.8; H, 11.1; N, 29.4. C₅H₁₄N₃F requires C, 44.4; H, 10.4; N, 29.4%); i.r. spectrum 12; n.m.r. spectrum 12; mass spectrum 12.

Two reactions were performed as described above (section 7.2.3); quantities used and yields are tabulated below:-

<u>Substrate</u>	<u>Substrate</u> g/mmol	<u>HF complex</u> g/mmol	<u>PhCF₃</u> g/mmol	<u>Yield</u> %
PhCOCl	1.40, 9.9	1.15, 9.0	0.21, 1.4	65
Dinitrochloro- benzene	1.37, 7.0	1.07, 8.0	0.24, 1.6	36

7.3.1 Attempted Preparation of Pentamethylguanidine (41)¹³³

Tetramethylguanidine (4.6g, 53mmol) and methyl iodide (7.6g, 54mmol) were stirred in toluene (50ml) at room temperature overnight. A white solid is slowly deposited and this was collected by filtration and identified as pentamethylguanidinium hydriodide (42)(3.2g, 29%); m.p. 137°C (from aq. EtOH) (lit¹³³, 137°C); (Found: C, 30.0; H, 6.5; N, 14.9. Calc for C₆H₁₆N₃l.0.5C₂H₅OH: C, 30.0; H, 6.8; N, 15.0%); δ_{H} (60MHz, D₂O, TMS) 2.9ppm (s, 4H, N-Me), 4.7 (s, 1H, =N-Me).

Heating the pentamethylguanidinium hydriodide salt with NaOH caused a mixture of tetramethylguanidine and pentamethylguanidine to be produced (gc/ms) which could not be separated by distillation.

CHAPTER 8**EXPERIMENTAL TO CHAPTER 4****8.1 Reaction between PS/HF, Perfluorocyclobutene and Hexafluoropropene**

A Carius tube was charged with 1,8-bis(dimethylamino)naphthalene hydrofluoride (1)(2.5g, 10.6mmol) in acetonitrile (20ml) and perfluorocyclobutene (6.7g, 41mmol) and hexafluoropropene (5.0g, 33mmol) were transferred under vacuum to the tube which was cooled in liquid air. After agitating on a rotating arm at room temperature for two days, the tube was opened to reveal a red solvent layer. On adding water (20ml), an orange solid precipitated and was collected by filtration. TLC showed that the solid contained two components. The solid was evaporated onto chromatographic alumina and light petroleum eluted 1,1-Bistrifluoromethyl-(6,7-bisdimethylamino)-2,3-tetrafluoroethano-[1H]-phenalene (4) (1.1g, 21%) as orange crystals; $R_F=0.5$; m.p. 128°C (from aqueous ethanol); λ_{max} (CH₃CN) 273.6nm (log₁₀ ϵ 3.78), 367.6 (3.45), 451.2 (3.78); (Found: C, 52.2; H, 3.4; N, 5.6; F, 38.0. C₂₁H₁₆N₂F₁₀ requires C, 51.9; H, 3.3; N, 5.8; F, 39.0%). n.m.r. spectrum 13; i.r. spectrum 13; mass spectrum 13.

A red solid (5) was also isolated (0.05g) as yet unidentified; δ_F (CFCl₃, CD₃CN, 235MHz) -68.9ppm (s, 6F), -110.5 (s, 2F); i.r. spectrum 14; mass spectrum 14.

8.2 Reaction Between PS and Co-dimer (3)

A mixture containing PS (0.7g, 3.2mmol), co-dimer (3)(1.0g, 3.2mmol) and acetonitrile (10ml) was stirred at room temperature for two days. Water was added to the mixture to precipitate an orange solid which was collected by filtration. The same procedure was then carried out as above (section 8.1) to yield pure (4)(0.35g, 22%) as orange crystals; m.p. 128°C; spectral data as above.

8.3 Preparation of Codimer (43)

A Carius tube was charged with pyridine (2.5g, 32mmol) and perfluorocyclobutene (20g, 0.12mol) and perfluorocyclopentene (25g, 0.12mol) were transferred to the tube which was cooled in liquid air. After rotating on a rotating arm for two days the tube was opened and all volatile products were transferred under vacuum to a trap. These were washed with water, dried (P₂O₅) and distilled on the Fischer Spahlrohr to yield perfluorobicyclobutylidene (62)(2.9g, 8%); b.p. 74-

85°C; and, codimer (43) (2.3g, 5%); b.p. 98-100°C; as compared to the literature data¹²¹.

8.4 Reaction between PS and Codimer (43)

A mixture containing PS (0.6g, 2.8mmol) and codimer (43) (1.0g, 2.7mmol) was refluxed overnight in acetonitrile (5ml). The solvent was removed to leave an orange solid which was washed with water, collected and dried. The solid was evaporated onto chromatographic alumina and 40/60 petroleum ether eluted Spiro[octafluorocyclopentane-1,1'-(6,7-bisdimethylamino)-2',3'-tetrafluoroethano-[1H]-phenalene (44)(0.2g, 13%) as orange crystals; m.p. 137-139°C; $R_F=0.5$; λ_{max} (CH₃CN) 273.6nm (log₁₀ ϵ 4.06), 365.6 (3.74), 452.8 (4.10); (Found: C, 50.4; H, 3.9; N, 4.4. C₂₃H₁₆N₂F₁₂ requires C, 50.3; H, 2.9; N, 5.1%). Mass required for C₂₃H₁₆N₂F₁₂: 548.11219. Found: 548.11762 a.m.u.; n.m.r. spectrum 14; i.r. spectrum 15; mass spectrum 15.

8.5 Reaction between PS and Perfluoro-bicyclopentylidene (55)

8.5.1 At Low Dilution

A mixture containing PS (1.1g, 5.1mmol) and perfluorobicyclopentylidene (55)(1.0g, 2.3mmol) was stirred overnight at room temperature in acetonitrile (5ml). A dark olive green precipitate formed. Water was added to the mixture and the solid was collected by filtration. The solid was absorbed onto chromatographic alumina and light petroleum/dichloromethane (4:1) eluted (7,8)-(9,10)-dihexafluoro-propano-(3,4-bisdimethylamino)-cyclohepta[d,e]-naphthalene (56)(0.54g, 42%); m.p. 243-45°C (decomp) (from acetonitrile); $R_F=0.65$; (Found: C, 51.3; H, 2.8; N, 4.9; F, 40.0. C₂₄H₁₆N₂F₁₂ requires C, 51.4; H, 2.8; N, 5.0; F, 40.7%). no n.m.r. could be recorded; i.r. spectrum 16; mass spectrum 16.

8.5.2 At High Dilution

A mixture containing PS (0.5g, 2.5mmol) and perfluorobicyclopentylidene (55)(1.0g, 2.3mmol) was stirred overnight at room temperature in acetonitrile (120ml). The solvent was removed under vacuum to leave a solid residue, which was absorbed onto chromatographic alumina and petroleum ether/dichloromethane (4:1) eluted (56)(0.23g, 18%), as above; Spiro-[octafluoro-cyclopentane-1,1'-(6,7-bisdimethylamino)-2',3'-tetrafluoropropan-1''-one-[1H]-phenalene (60)(0.14g, 10%) as bright green metallic looking flakes; m.p. >280°C; $R_F=0.45$; (Found: C, 48.4; H, 2.75; N, 4.50. C₂₄H₁₆N₂F₁₂O requires C, 50.0; H, 2.75; N, 4.5%).

$C_{24}H_{16}N_2F_{12}O \cdot H_2O$ requires C, 48.5; H, 3.0; N, 4.7%; n.m.r. spectrum 15; i.r. spectrum 17; mass spectrum 17; and, 7,8-propano-9,10-propan-1'-one-cyclohepta-[d,e]-naphthalene (61)(0.12g, 9%) as bright purple metallic looking flakes; m.p. $>280^\circ C$; $R_F=0.3$; (Found: C, 53.2; H, 3.05; N, 4.75. $C_{24}H_{16}N_2F_{10}O$ requires C, 53.5; H, 2.95; N, 5.2%); $C_{24}H_{16}N_2F_{10}O$ requires 538.110295 a.m.u. Found 538.1100900 a.m.u; n.m.r. spectrum 16; i.r. spectrum 18; mass spectrum 18.

8.6 Preparation of Perfluorobicyclobutylidene (62)

A Carius tube was charged with pyridine (1.04g, 13mmol) and perfluorocyclobutene (28.3g, 0.15mol) was transferred to the tube which was cooled in liquid air. After agitating on a rotating arm for two days the tube was opened and all volatiles were transferred under vacuum. These were washed with water and the lower fluorocarbon layer was separated, dried (P_2O_5) and distilled on the Fischer Spahlrohr to yield perfluorobicyclobutylidene (62)(5.2g, 10%); b.p. $74-85^\circ C$; as compared to the literature data¹²¹.

8.7 Reaction between N,N-Dimethylaniline and Perfluorobicyclobutylidene

A mixture containing N,N-Dimethylaniline (0.5g, 4.1mmol) and perfluorobicyclobutylidene (62)(1.1g, 3.4mmol) was stirred at room temperature overnight in acetonitrile (5ml). Water (15ml) was added to the mixture to precipitate the solid product which was collected by filtration. Recrystallisation from aqueous ethanol and vacuum sublimation yielded Spiro[hexafluorocyclobutane-3,1'-1,2-tetrafluoroethano-1-fluoro-3-(4''-N,N-dimethylaminophenyl)-propene] (63) (1.05g, 73%) as white needles; m.p. $84-85^\circ C$; (Found: C, 45.0; H, 2.3; N, 3.2. $C_{16}H_{10}NF_{11}$ requires C, 45.2; H, 2.3; N, 3.3%). n.m.r. spectrum 17; i.r. spectrum 19; mass spectrum 19.

8.8 Reaction of N-methylindole with Perfluorobicyclobutylidene (62)

A mixture containing N-methylindole (0.4g, 3mmol) and perfluorobicyclobutylidene (62)(1.0g, 3mmol) was refluxed in acetonitrile (5ml) for 1 hr. On cooling, water (15ml) was added to the reaction mixture to precipitate the solid product which was collected by filtration, dried and purified by vacuum sublimation (oil bath temperature $100^\circ C$, <0.1 mm Hg) to white crystals and identified as Spiro[hexafluorocyclobutane-3,1'-1,2-tetrafluoroethano-1-fluoro-3-(N-methyl-indol-3''-yl)-propene] (65)(0.60g, 46%); m.p. $59-60^\circ C$; (Found: C, 47.25; H,

1.8; N, 3.1. $C_{17}H_8NF_{11}$ requires C, 46.9; H, 1.85; N, 3.2%). n.m.r. spectrum 18; i.r. spectrum 20; mass spectrum 20.

8.9 Reaction between 1,8-Bis(dimethylamino)naphthalene and Perfluoro-bicyclobutylidene

A mixture containing 1,8-bis(dimethylamino)naphthalene (1.16g, 5.4mmol) and perfluorobicyclobutylidene (62)(1.56g, 4.8mmol) was stirred overnight at room temperature in acetonitrile (5ml). An orange solid precipitated which was collected by filtration and washed with water. Recrystallisation from aqueous acetonitrile yielded a white solid which was identified as the substitution product (66) (0.83g, 35%); m.p. 210-215°C (decomp); (Found: C, 53.2; H, 3.6; N, 5.6; F, 29.7. $C_{22}H_{18}N_2F_8O_2$ requires C, 53.4; H, 3.6; N, 5.6; F, 30.7%). n.m.r. spectrum 19; i.r. spectrum 21.

CHAPTER NINE

EXPERIMENTAL TO CHAPTER FIVE

9.1 Preparation of Trifluoro-s-triazine (13)

An autoclave (460ml, No. 14) was charged with trichloro-s-triazine (37g, 0.2mol) and flame dried potassium fluoride (125g, 2mol), evacuated and then heated at 310°C for 16hrs (Furnace No. 3). The volatile products were transferred from the hot autoclave to a trap which was cooled in liquid air. The product, trifluoro-s-triazine (13)(20g, 74%) did not require any further purification and was stored in a rotaflo tube.

9.2 Preparation of Perfluoroisopropyl-s-triazine (14) and Perfluoro-diisopropyl-s-triazine (15)

A flask was charged with trifluoro-s-triazine (13)(15.0g, 0.11mol), potassium fluoride (5.0g, 86mmol) and dry-sulpholane (100ml). The flask was frozen down in liquid air and evacuated. Hexafluoropropene (17g, 0.11mol) was added via a bladder and the reaction mixture was heated at 70°C for 16hrs with vigorous stirring. All volatile products were removed by transfer under vacuum and then distilled on the Fischer Spahlrohr to yield perfluoroisopropyl-s-triazine (14)(5.8g, 18%); b.p. 105-106°C; m/z (EI⁺) 285 (M⁺, 43%) and perfluorodi-isopropyl-s-triazine (15)(4.9g, 10%); b.p. 133.7-135.2°C; m/z (EI⁺) 435 (M⁺, 24.5%); as compared to the literature data¹²⁶.

9.3 Reactions of Pyrroles with Fluorinated Triazines

General Procedure - A mixture containing a pyrrole and the corresponding fluorinated triazine was refluxed in acetonitrile (5ml) for two hours. On cooling, water (15ml) was added to the reaction mixture to precipitate the solid product, which was collected by filtration, dried in a desiccator and purified by vacuum sublimation (Oil bath temperature 130°C, <0.1mm Hg). All yields are quoted for pure, isolated products.

9.3.1 Reaction of Pyrrole with perfluoro-s-triazine (13)

Pyrrole (0.5g, 7.4mmol) and perfluoro-s-triazine (13)(1.0g, 7.4mmol) gave 2-(pyrrol-2-yl)-4,6-difluoro-s-triazine (67)(0.87g, 65%) as white crystals; m.p. 156-160°C; (Found: C, 46.0; H, 2.1; N, 30.8. C₇H₄N₄F₂ requires C,

46.15; H, 2.2; N, 30.75%); λ_{\max} (CH₃CN) 310.1nm (log₁₀ε 4.48); n.m.r. spectrum 20; i.r. spectrum 22; mass spectrum 21.

9.3.2 Reaction of Pyrrole with perfluoroisopropyl-s-triazine (14)

Pyrrole (0.80g, 12mmol) and perfluoroisopropyl-s-triazine (14)(2.5g, 9mmol) gave 2-(pyrrol-2-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine (68)(2.05g, 71%) as pale yellow needles; m.p. 112-115°C; (Found: C, 35.7; H, 1.2; N, 16.5; C₁₀H₄N₄F₈ requires C, 36.1; H, 1.2; N, 16.85%); λ_{\max} (CH₃CN) 336.0nm (log₁₀ε 4.41). n.m.r. spectrum 21; i.r. spectrum 23; mass spectrum 22.

9.3.3 Reaction of Pyrrole with Perfluorodi-isopropyl-s-triazine (15)

Pyrrole (0.27g, 4.0mmol) and perfluorodi-isopropyl-s-triazine (15)(1.46g, 3.3mmol) gave 2-(pyrrol-2-yl)-4,6-perfluorodi-isopropyl-s-triazine (69)(1.1g, 68%) as pale yellow needles; m.p. 64-66°C; (Found: C, 32.2; H, 0.75; N, 11.5. C₁₃H₄N₄F₁₄ requires C, 32.4; H, 0.8; N, 11.6%); λ_{\max} (CH₃CN) 344.8nm (log₁₀ε 4.58). n.m.r. spectrum 22; i.r. spectrum 24; mass spectrum 23.

9.3.4 Reaction of N-methylpyrrole with perfluoro-s-triazine (13)

N-methylpyrrole (1.0g, 12mmol) and perfluoro-s-triazine (13)(1.6g, 12mmol) gave 2-(N-methylpyrrol-2-yl)-4,6-difluoro-s-triazine (70)(1.06g, 45%) as pale yellow needles; m.p. 117-118°C; (Found: C, 48.5; H, 3.0; N, 28.4. C₈H₆N₄F₂ requires C, 49.0; H, 3.1; N, 28.6%); λ_{\max} (CH₃CN) 314.6nm (log₁₀ε 4.44) n.m.r. spectrum 23; i.r. spectrum 25; mass spectrum 24.

9.3.5 Reaction of N-methylpyrrole with perfluoroisopropyl-s-triazine (14)

N-methylpyrrole (0.30g, 3.7mmol) and perfluoroisopropyl-s-triazine (14)(1.05g, 3.7mmol) gave 2-(N-methylpyrrol-2-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine (71)(0.69g, 54%) as pale yellow needles; m.p. 110-111°C; (Found: C, 38.2; H, 1.5; N, 15.9; F, 43.4. C₁₁H₆N₄F₈ requires C, 38.15; H, 1.7; N, 16.2; F, 43.95%); λ_{\max} (CH₃CN) 324.5nm (log₁₀ε 4.38). n.m.r. spectrum 24; i.r. spectrum 26; mass spectrum 25.

9.3.6 Reaction of N-methylpyrrole with Perfluorodi-isopropyl-s-triazine (15)

N-methylpyrrole (0.3g, 3.7mmol) and perfluorodi-isopropyl-s-triazine (15)(1.5g, 3.5mol) gave 2-(N-methylpyrrol-2-yl)-4,6-perfluorodi-isopropyl-s-triazine (72)(0.73g, 48%) as pale yellow needles; m.p. 88-89°C; (Found: C, 33.9; H, 1.1; N, 11.3; F, 53.9. C₁₄H₆N₄F₁₄ requires C, 33.9; H, 1.2; N, 11.3; F, 53.6%); λ_{\max} (CH₃CN) 349.4nm (log₁₀ε 4.33). n.m.r. spectrum 25; i.r. spectrum 27; mass spectrum 26.

9.4 Reactions of N-methylindole with Fluorinated Triazines

General Procedure - A mixture containing N-methylindole and the corresponding fluorinated triazine was refluxed in acetonitrile (5ml) for 30mins. On cooling, water (15ml) was added to precipitate the solid product which was dried and purified by vacuum sublimation (Oil bath temperature 150°C, <0.1mm Hg).

9.4.1 Reaction of N-methylindole with perfluoro-s-triazine (13)

N-methylindole (1.0g,7.6mmol) and perfluoro-s-triazine (13)(1.0g, 7.4mmol) gave 2-(N-methylindol-3-yl)-4,6-difluoro-s-triazine (73)(1.43g, 78%) as white crystals (from acetone); m.p. 244°C; (Found: C, 58.25; H, 3.25; N, 22.6. C₁₂H₈N₄F₂ requires C, 58.5; H, 3.25; N, 22.75%). n.m.r. spectrum 26; i.r. spectrum 28; mass spectrum 27.

9.4.2 Reaction of N-methylindole with Perfluoroisopropyl-s-triazine (14)

N-methylindole (0.7g, 5.3mmol) and perfluoroisopropyl-s-triazine (14)(1.5g, 5.3mmol) gave 2-(N-methylindol-3-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine (74)(1.76g, 90%) as pale yellow needles; m.p. 190-194°C; (Found: C, 45.7; H, 2.0; N, 14.2; F, 38.1. C₁₅H₈N₄F₈ requires C, 45.45; H, 2.0; N, 14.1; F, 38.4%); λ_{\max} (CH₃CN) 262.4nm (log₁₀ε 4.12), 355.3 (4.31);. n.m.r. spectrum 27; i.r. spectrum 29; mass spectrum 28.

9.4.3 Reaction of N-methylindole with Perfluorodi-isopropyl-s-triazine (15)

N-methylindole (0.3g, 2.3mmol) and perfluorodi-isopropyl-s-triazine (15)(1.0g, 2.3mmol) gave 2-(N-methylindol-3-yl)-4,6-perfluorodi-isopropyl-

s-triazine (75)(1.1g, 87%) as yellow crystals; m.p. 205-206°C; (Found: C, 39.25; H, 1.45; N, 9.9; F, 48.9. C₁₈H₈N₄F₁₄ requires C, 39.55; H, 1.45; N, 10.25; F, 48.7%); λ_{\max} (CH₃CN) 213.0nm (log₁₀ε 4.56), 246.0 (4.02), 265.0 (4.23), 276.0 (4.14), 365.0 (4.53); n.m.r. spectrum 28; i.r. spectrum 30; mass spectrum 29.

9.5 Reaction of N-methylindole with Tetrafluoropyrimidine

A mixture containing N-methylindole (0.8g, 6mmol) and tetrafluoropyrimidine (1.0g, 6.5mmol) was refluxed overnight in acetonitrile (5ml). On cooling, water (15ml) was added to the reaction mixture to precipitate the solid product which was collected by filtration, dried, recrystallised from acetone as yellow plates and identified as **6-(N-methylindol-3-yl)-2,4,5-trifluoropyrimidine (76)**(0.58g, 36%); m.p. 231°C; (Found: C, 59.35; H, 3.0; N, 15.9; F, 22.0. C₁₃H₈N₃F₃ requires C, 59.3; H, 3.05; N, 15.95; F, 21.7%); λ_{\max} (CH₃CN) 214.0nm (log₁₀ε 4.75), 263.0 (4.35), 344.0 (4.78); i.r. spectrum 31; mass spectrum 30.

9.6 Reactions of Anilines With Fluorinated s-Triazines (13)-(15)

9.6.1 Reaction of N,N-Dimethylaniline with Trifluoro-s-triazine (13)

A mixture containing N,N-Dimethylaniline (1.75g, 14mmol) and trifluoro-s-triazine (13)(2.0g, 15mmol) was refluxed overnight in acetonitrile (5ml). On cooling a red/brown solid precipitated which was collected by filtration, washed with water and recrystallised from acetonitrile to yield pure **2-(4-N,N-dimethylaminophenyl)-4,6-difluoro-s-triazine (77)**(0.96g, 28%); m.p. 234-237°C; (Found: C, 55.6; H, 4.05; N, 23.5. C₁₁H₁₀N₄F₂ requires C, 55.9; H, 4.25; N, 23.7%); λ_{\max} (CH₃CN) 364.0nm (log₁₀ε 4.51); C₁₁H₁₀N₄F₂ requires 236.08735amu. Found 236.08416amu; n.m.r. spectrum 29; i.r. spectrum 32; mass spectrum 31.

9.6.2 Reaction of N,N-Dimethylaniline with Perfluoroisopropyl-s-triazine (14)

A mixture containing N,N-Dimethylaniline (0.5g, 4.1mmol) and perfluoroisopropyl-s-triazine (1.2g, 4.2mmol) was refluxed in acetonitrile (5ml) for 2 hr. On cooling, water (15ml) was added to precipitate the solid product which was collected by filtration and dried. Vacuum sublimation yielded pure **2-(4-N,N-dimethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazine (78)**(0.62g,

36%) as a yellow solid; m.p. 168-170°C; (Found: C, 42.8; H, 2.5; N, 14.2. C₁₄H₁₀N₄F₈ requires C, 43.5; H, 2.6; N, 14.5%); λ_{\max} (CH₃CN) 368.8nm (log₁₀ε 4.40). n.m.r. spectrum 30; i.r. spectrum 33; mass spectrum 32.

9.6.3 Reaction of N,N-Diethylaniline with Perfluoroisopropyl-s-triazine (14)

A mixture containing N,N-Diethylaniline (0.5g, 3.4mmol) and perfluoroisopropyl-s-triazine (14)(1.0g, 3.5mmol) was refluxed in acetonitrile (5ml) for 2 hr. On cooling, water (15ml) was added to precipitate the solid product which was collected by filtration and dried. Vacuum sublimation yielded a mixture of the two isomers 2-(4-N,N-diethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazine (79A) and 2-(2-N,N-diethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazine (79B)(0.63g, 45%) as a yellow solid; (Found: C, 46.0; H, 3.5; N, 13.6. C₁₆H₁₄N₄F₈ requires C, 46.35; H, 3.4; N, 13.5%); λ_{\max} (CH₃CN) 228.0nm (log₁₀ε 3.86), 260.9 (3.93), 405.3 (4.39). n.m.r. spectrum 32; i.r. spectrum 35; mass spectrum 34.

9.6.4 Reaction of N,N-Dimethylaniline with Perfluorodi-isopropyl-s-triazine (15)

A mixture containing N,N-Dimethylaniline (0.5g, 4.1mmol) and perfluorodi-isopropyl-s-triazine (15)(1.5g, 3.4mmol) was refluxed in acetonitrile (5ml) for 3 hr. On cooling, water (15ml) was added to the reaction mixture to precipitate the solid product which was collected by filtration and dried. Vacuum sublimation yielded a mixture of the two isomers 2-(4-N,N-dimethylaminophenyl)-4.6-perfluorodiisopropyl-s-triazine (81A) and 2-(2-N,N-dimethylaminophenyl)-4.6-perfluorodiisopropyl-s-triazine (81B)(0.95g, 77%) as a yellow solid; (Found: C, 37.9; H, 2.0; N, 10.3. C₁₇H₁₀N₄F₁₄ requires C, 38.05; H, 1.85; N, 10.45%); λ_{\max} (CH₃CN) 412.0nm (log₁₀ε 4.28); n.m.r. spectrum 31; i.r. spectrum 34; mass spectrum 33.

9.6.5 Reaction of N,N-Diethylaniline with Perfluorodi-isopropyl-s-triazine (15)

A mixture containing N,N-Diethylaniline (0.6g, 4mmol) and perfluorodi-isopropyl-s-triazine (15)(1.0g, 2.3mmol) was refluxed in acetonitrile (5ml) for 2 hr. On cooling, water (15ml) was added to the reaction mixture to precipitate an orange oil which solidified on standing. This solid was washed repeatedly with water and analysis confirmed the solid to be a mixture of the two isomers 2-(4-N,N-

diethylaminophenyl)-4,6-perfluorodiisopropyl-s-triazine (82A) and 2-(2-N,N-diethylaminophenyl)-4,6-perfluorodiisopropyl-s-triazine (82B)(0.93g, 72%); (Found: C, 40.3; H, 2.7; N, 9.6. $C_{19}H_{14}N_4F_{14}$ requires C, 40.4; H, 2.5; N, 9.9%); λ_{max} (CH₃CN) 419.6nm (log₁₀ε 4.47); n.m.r. spectrum 33; i.r. spectrum 36; mass spectrum 35.

9.6.6 Reaction between 1,8-(Bisdimethylamino)-naphthalene and Trifluoro-s-triazine (13)

A mixture containing 1,8-(bisdimethylamino)-naphthalene (2.1g, 10mmol) and trifluoro-s-triazine (13)(1.5g, 11mmol) was stirred at room temperature overnight in acetonitrile (5ml). The solution turned orange immediately and gradually red crystals precipitated which were collected by filtration and recrystallised from acetonitrile to yield pure 1,8-(bisdimethylamino)-4,5-(bisdifluoro-s-triaz-1-yl)-naphthalene (83)(2.3g, 53%); m.p. 258-260°C; (Found: C, 53.7; H, 3.6; N, 24.9. $C_{20}H_{16}N_8F_4$ requires C, 54.0; H, 3.6; N, 25.2%). $C_{20}H_{16}N_8F_4$ requires 444.1434amu. Found 444.1272amu; n.m.r. spectrum 34; i.r. spectrum 37; mass spectrum 36.

9.6.7 Reaction of N-ethylaniline with Trifluoro-s-triazine (13)

A mixture containing N-ethylaniline (0.9g, 7.4mmol) and trifluoro-s-triazine (13)(1.0g, 7.4mmol) was refluxed in acetonitrile (5ml) for 3 hr. The solvent was removed under reduced pressure to leave an off-white solid which was washed with water and collected by filtration. Vacuum sublimation yielded pure 2-(ethylphenylamino)-4,6-difluoro-s-triazine (84)(1.2g, 69%) as white needles; m.p. 63.5-64°C; (Found: C, 55.6; H, 4.0; N, 24.0. $C_{11}H_{10}N_4F_2$ requires C, 55.9; H, 4.2; N, 23.7%); λ_{max} (CH₃CN) 236.0nm (log₁₀ε 4.26). n.m.r. spectrum 35; i.r. spectrum 38; mass spectrum 37.

9.6.8 Reaction of N-ethylaniline with Perfluoroisopropyl-s-triazine (14)

A mixture containing N-ethylaniline (0.63g, 5.2mmol) and perfluoroisopropyl-s-triazine (14)(1.5g, 5.2mmol) was refluxed overnight in acetonitrile (5ml). The solvent was removed under reduced pressure to leave an off-white solid which was washed with water and collected by filtration. Vacuum sublimation yielded pure 2-(ethylphenylamino)-4-perfluoroisopropyl-s-triazine (80)(1.8g, 89%) as white crystals; m.p. 70-72°C; (Found: C, 43.8; H, 2.65; N, 14.8; F, 40.0. $C_{14}H_{10}N_4F_8$ requires C, 43.5; H, 2.6; N, 14.5; F, 39.4%); λ_{max}

(CH₃CN) 239.0nm (log₁₀ε 4.30); n.m.r. spectrum 36; i.r. spectrum 39; mass spectrum 38.

9.6.9 Preparation of N,N-Dimethylamino-2-methoxybenzene

A flask was charged with 2-methoxyaniline (20g, 0.16mol) and trimethylphosphite (22g, 0.16mol). The mixture was heated until a fine mist appeared after which the heat source was removed and the reaction allowed to subside. The reaction was then heated at reflux for a further 2 hrs. After cooling to 90°C, sodium hydroxide solution (22g in 170ml water) was added and the aqueous mixture was left to stand for 1.5hrs. The amines were extracted with ether and distilled on the Fischer Spahlrohr to yield N,N-Dimethylamino-2-methoxybenzene (13.0g, 54%); pure by GC; b.p. 81.7-82°C/5mm Hg; IR spectrum recorded; m/z (EI⁺) 151 (M⁺, 100%).

9.6.10 Reaction of N,N-Dimethylamino-2-methoxybenzene with Trifluoro-s-triazine (13)

A mixture containing N,N-Dimethylamino-2-methoxybenzene (1.0g, 6.6mmol) and trifluoro-s-triazine (13)(1.2g, 8.8mmol) was refluxed overnight in acetonitrile (5ml). Water (15ml) was added to the mixture to precipitate the solid product which was collected by filtration. Vacuum sublimation yielded pure 2-methyl(2-methoxyphenyl)-amino-4,6-difluoro-s-triazine (85)(0.51g, 31%) as white crystals; m.p. 139°C; (Found: C, 52.25; H, 3.85; N, 22.35. C₁₁H₁₀N₄OF₂ requires C, 52.4; H, 3.95; N, 22.2%). n.m.r. spectrum 37; i.r. spectrum 40; mass spectrum 39.

9.6.11 Reaction of N,N,N',N'-tetramethyl-1,4-diaminobenzene with trifluoro-s-triazine (13)

A mixture containing N,N,N',N'-tetramethyl-1,4-diaminobenzene (0.6g, 3.6mmol) and trifluoro-s-triazine (13)(0.4g, 3.0mmol) was refluxed overnight in acetonitrile (5ml). Water was added to the mixture to precipitate the solid product which was collected by filtration. Vacuum sublimation yielded pure 2-methyl(4-N,N-dimethylaminophenyl)amino-4,6-difluoro-s-triazine (86) (0.12g, 15%) as white crystals; (Found: C, 54.5; H, 5.1; N, 26.7. C₁₂H₁₃N₅F₂ requires C, 54.3; H, 4.9; N, 26.4%). n.m.r. spectrum 38; i.r. spectrum 41; mass spectrum 40.

9.6.12 Reaction between 2-N,N-dimethylamino-pyridine and Perfluorodi-isopropyl-s-triazine (15)

A mixture containing 2-N,N-dimethylaminopyridine (0.5g, 3.5mmol) and perfluorodi-isopropyl-s-triazine (1.5g, 3.4mmol) was refluxed in acetonitrile (5ml) for 6 hr. On cooling, water (15ml) was added to precipitate the solid product which was collected by filtration. Vacuum sublimation yielded the pyridinium salt (87)(0.9g, 47%) as a pale yellow solid; (Found: C, 34.65; H, 1.7; N, 12.8. $C_{16}H_{11}N_5OF_{14}$ requires C, 34.6; H, 2.0; N, 12.6%). n.m.r. spectrum 39; i.r. spectrum 42; mass spectrum 41.

APPENDICES

APPENDIX ONE**NUCLEAR MAGNETIC RESONANCE SPECTRA**

1. 1,8-(Bisdimethylamino)-naphthalene Hydrogen Fluoride Complex (PS/HF) (1)
2. Triethylamine Hydrogen Fluoride Complex (24)
3. Tributylamine Hydrogen Fluoride Complex (25)
4. Trihexylamine Hydrogen Fluoride Complex (26)
5. Trioctylamine Hydrogen Fluoride Complex (27)
6. Tridodecylamine Hydrogen Fluoride Complex (28)
7. 2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (35)
8. N-Methyl-2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (36)
9. N-Ethyl-2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (37)
10. N-Allyl-2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (38)
11. N-Benzyl-2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (39)
12. Tetramethylguanidine Hydrogen Fluoride Complex (40)
13. 1,1-Bistrifluoromethyl-6,7-bisdimethylamino-2,3-tetrafluoro-ethano-[1H]-phenalene (4)
14. Spiro[octafluorocyclopentane-1,1'-(6,7-bisdimethylamino)-2',3'-tetrafluoroethano-[1H]-phenalene] (44)
15. Spiro[octafluorocyclopentane-1,1'-(6,7-bisdimethylamino)-2',3'-tetrafluoro-propan-1''-one-[1H]-phenalene] (60)
16. 7,8-propano-9,10-propan-1''-one-cyclohepta-[d,e]-naphthalene (61)
17. Spiro[hexafluorocyclobutane-3,1'-1,2-tetrafluoroethano-1-fluoro-3-(4''-N,N-dimethylaminophenyl)-propene] (63)
18. Spiro[hexafluorocyclobutane-3,1'-1,2-tetrafluoroethano-1-fluoro-3-(N-methylindol-3''-yl)-propene] (65)
19. White Solid (66)
20. 2-(pyrrol-2-yl)-4,6-difluoro-s-triazine (67)
21. 2-(pyrrol-2-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine (68)
22. 2-(pyrrol-2-yl)-4,6-perfluorodi-isopropyl-s-triazine (69)
23. 2-(N-methylpyrrol-2-yl)-4,6-difluoro-s-triazine (70)
24. 2-(N-methylpyrrol-2-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine (71)
25. 2-(N-methylpyrrol-2-yl)-4,6-perfluorodi-isopropyl-s-triazine (72)
26. 2-(N-methylindol-3-yl)-4,6-difluoro-s-triazine (73)
27. 2-(N-methylindol-3-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine (74)
28. 2-(N-methylindol-3-yl)-4,6-perfluorodi-isopropyl-s-triazine (75)
29. 2-(4-N,N-dimethylaminophenyl)-4,6-difluoro-s-triazine (77)
30. 2-(4-N,N-dimethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazine (78)

31. 2-(4-N,N-dimethylaminophenyl)-4,6-perfluorodi-isopropyl-s-triazine (81A) and 2-(2-N,N-dimethylaminophenyl)-4,6-perfluorodi-isopropyl-s-triazine (81B)
32. 2-(4-N,N-diethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazine (79A) and 2-(2-N,N-diethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazine (79B)
33. 2-(4-N,N-diethylaminophenyl)-4,6-perfluorodi-isopropyl-s-triazine (82A) and 2-(2-N,N-diethylaminophenyl)-4,6-perfluorodi-isopropyl-s-triazine (82B)
- 34 1,8-(Bisdimethylamino)-4,5-(bisdifluoro-s-triaz-2-yl)-naphthalene (83)
35. 2-(ethylphenylamino)-4,6-difluoro-s-triazine (84)
36. 2-(ethylphenylamino)-4-perfluoroisopropyl-6-fluoro-s-triazine (80)
37. 2-methyl-(2-methoxyphenyl)-amino-4,6-difluoro-s-triazine (85)
38. 2-methyl-(4-N,N-dimethylaminophenyl)-amino-4,6-difluoro-s-triazine (86)
39. Pyridinium Salt (87)


NMR spectra were recorded in d_3 -acetonitrile solutions unless otherwise stated. Reference compounds (^1H and ^{13}C - Me_4Si , ^{19}F - CFCl_3) were used internally.

^1H NMR Spectrum of PS/HF in CD_3CN


Assignment of ^1H NMR spectrum discussed in section 2.3.2.1

^{19}F NMR Spectrum

-169.0ppm (s)




No. 2 Triethylamine HF Complex




Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
¹H				
1.21	t	J _{a, b} = 7.3	9H	a
2.96	q		6H	b
12.68	s (br)		1H	c
¹⁹F				
-154.81	s			d

No. 3 Tri-n-butylamine HF Complex




Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
¹H				
1.07	m		9H	a
1.51	m		6H	b
1.78	m		6H	c
3.04	m		6H	d
12.67	s (br)		1H	e
¹⁹F				
-157.09	s			f

No. 4 Tri-n-hexylamine HF Complex




Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
¹H				
0.99	t	J _{a, b} = 6.6	9H	a
1.41	m		18H	b, c, d
1.69	m		6H	e
2.87	t	J _{e, f} = 7.0	6H	f
12.50	s (br)		1H	g
¹⁹F				
-157.04	s			h

No. 5 Tri-n-octylamine HF Complex



Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
¹H				
1.07	t	J _{a, b} = 6.1	9H	a
1.49-1.78	m		36H	b, c, d, e, f, g
2.96	m		6H	h
10.76	s (br)		1H	i
¹⁹F				
-156.6	s			j

No. 6 Tri-n-dodecamine HF Complex




Chemical Shift (ppm)	Multiplicity Coupling Constants (Hz)	Relative Intensity	Assignment
----------------------	---	--------------------	------------

¹H			
0.4-2.90	m (overlapping peaks)		CH ₃ , CH ₂
11.61	s (br)		a

¹⁹F			
-159.30	s		b

No. 7 2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex




Chemical Shift (ppm)	Multiplicity Coupling Constants (Hz)	Relative Intensity	Assignment
----------------------	---	--------------------	------------

¹H			
1.21	s	12H	a
1.52	m	4H	b
1.66	m	2H	c
4.90	s (br)	1H	d
9.30	s (br)	1H	e

¹⁹F			
-134.96	s (br)		f

No. 8 1,2,2,6,6-Pentaméthylpiperidine HF Complex




Chemical Shift (ppm)	Multiplicity Coupling Constants (Hz)	Relative Intensity	Assignment
----------------------	---	--------------------	------------

¹ H			
1.27	s	12H	a
1.68	m	6H	b, c
2.48	s	3H	d
12.74	s (br)	1H	e

¹⁹ F			
-148.9	s (br)		f

No. 9 N-Ethyl-2,2,6,6-Tetraméthylpiperidine HF Complex




Chemical Shift (ppm)	Multiplicity Coupling Constants (Hz)	Relative Intensity	Assignment
----------------------	---	--------------------	------------

¹ H			
1.32	s	12H	a
1.32	t	3H	e
1.71	m	2H	c
1.81	m	4H	b
3.00	q J _{d,e} =7.4	2H	d
11.10	s (br)	1H	f


¹⁹ F			
-143.9	s (br)		g

No. 10 **N-Allyl-2,2,6,6-Tetramethylpiperidine HF Complex**



Chemical Shift (ppm)	Multiplicity Coupling Constants (Hz)	Relative Intensity	Assignment
¹ H			
1.32	s	12H	a
1.68	m	2H	c
1.73	m	4H	b
3.62	d J _{d, e} =6.0	2H	d
5.26	dd J _{e, f} =17.0	2H	f, g
6.11	ddt J _{e, g} =10.0	1H	e
12.1	s (br)	1H	h
¹⁹ F			
-146.0			i

No. 11 **N-Benzyl-2,2,6,6-Tetramethylpiperidine HF Complex**



Chemical Shift (ppm)	Multiplicity Coupling Constants (Hz)	Relative Intensity	Assignment
¹ H			
1.23	s	12H	a
1.71	m	6H	b, c
4.11	s	2H	d
7.25	t J _{f, g} =7.3	1H	g
7.30	t J _{e, f} =7.6	2H	f
7.54	d	2H	e
11.7	s (br)	1H	h
¹⁹ F			
-152.4	s (br)		


No. 12 **Tetramethylguanidine Hydrogen Fluoride Complex**



Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
----------------------	--------------	-------------------------	--------------------	------------

¹ H				
2.71	s		12H	a
3.63	s		1H	b
8.63	s (br)		1H	c
¹⁹ F				
-142.97				d

No. 13 **1,1-Bistrifluoromethyl-6,7-bisdimethylamino-2,3-tetrafluoro-ethano-[1H]-phenalene**



123.7	s	f
126.0	s	i
126.9	s	d
130.4	s	m
134.1	s	r
152.9	s	b
153.0	m	j
156.1	s	o

The complexity of the ^{13}C spectrum prevents a full assignment, especially for peaks in the region of 115-125ppm.

Chemical Shift (ppm)	Multiplicity Coupling Constants (Hz)	Relative Intensity	Assignment
----------------------	---	--------------------	------------

^1H

2.81	s	6H	a
2.88	s	6H	p
6.79	AX $J_{AX}=8.3$	1H	c
7.45		1H	d
6.98	AX $J_{AX}=8.8$	1H	n
7.85		1H	m


^{19}F

-67.23	s	6F	k
-105.10	s	2F	g
-112.55	s	2F	h

^{13}C

43.5	s (br)		a, p
107.7	s		l
108.9	s		c
111.5	s		n
115-120	many overlapping peaks		CF_2 and CF_3 g,h,k
116.7	s		e
121.4	s		q

No. 14 **Spiro[octafluorocyclopentane-1,1']-(6,7-bisdimethylamino)-2,3'-tetrafluoroethano -[1H]-phenalene]**



109.4	s	c
111.4	s	o
115.4	s	r
115-120	many overlapping peaks	g, h, k, l
128.6	s	d
134.0	s	s
135.8	s	n
151.9	m	j
154.4	s	b
158.2	s	p

<u>Chemical Shift</u> <u>(ppm)</u>	<u>Multiplicity</u>	<u>Relative Intensity</u>	<u>Assignment</u>
	<u>Coupling Constants (Hz)</u>		

¹H

2.88	s	6H	a
2.94	s	6H	q
6.86	} AX J _{AX} = 8.4	1H	c
7.33		1H	d
7.04	} AX J _{AX} = 8.8	1H	o
7.52		1H	n

¹⁹F


-104.9	s	2F	g
-112.8	} AB J _{AB} = 249.1	4F	k
-116.0			
-114.6	s	2F	h
-135.2	s	4F	l

¹⁹F 2-D COSY spectrum also recorded.

¹³C

43.0	s (br)	a, q
105.6	s	
107.4		


No. 15 Spiro[octafluorocyclopentane-1,1'-(6,7-bisdimethylamino)-2',3'-tetrafluoropropan-2''-one-[1H]-phenalene]



Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
2.16	s	12H	a, j
6.95	AX J _{AX} = 8.8	1H	b
7.25		1H	c
7.05		1H	i
7.92	AX J _{AX} = 8.8	1H	j
-111.0	AB J _{AB} = 246.7	4F	f
-113.4			
-117.8	s (pseudo AB)	2F	d
-131.4	s (pseudo AB)	2F	e
-132.3	s (pseudo AB)	4F	g

¹⁹F 2-D COSY spectrum also recorded.


No. 17 Spiro-[hexafluorocyclobutane-3,1'-1,2-tetrafluoroethano-1-fluoro-3-(4''-N,N-Dimethylaminophenyl)-propene]



Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
2.96	s	6H	a
6.76	AB J _{AB} =8.7	2H	b
7.16		2H	c
-101.22	s	1F	h
-114.04	s	2F	f
-119.04	s	2F	g
-117.05	AB J _{AB} =214	4F	d
-120.93			
-127.99	AB J _{AB} =222	2F	e
-133.89			

No. 16

7,8-Propano-9,10-propan-2''-one-cyclo-hepta-[d,e]-naphthalene



AB $J_{AB} = 240$ 2F g

-143.4

^{19}F 2-D COSY spectrum also recorded. The midpoints of the AB systems are -117.6, -121.2, -123.7, -132.4 and -138.8ppm.

Chemical Shift (ppm)	Multiplicity Coupling Constants (Hz)	Relative Intensity	Assignment
----------------------	---	--------------------	------------

^1H


2.16	s	12H	a, k
6.49	-AX $J_{AX} = 8.8$	1H	j
6.92		1H	i
6.51	-AX $J_{AX} = 8.8$	1H	b
7.33		1H	c

^{19}F

-106.5	-AB $J_{AB} = 278$	2F	d
-128.8			
-109.3	-AB $J_{AB} = 262$	2F	h
-133.1			
-119.4	-AB $J_{AB} = 260$	2F	f
-128.0			
-128.8	-AB $J_{AB} = 283$	2F	e
-136.0			
-133.4			

No. 18

Spiro-[hexafluorocyclobutane-3,1'-1,2-tetrafluoroethano-1-fluoro-3-(N-Methylindol-3-yl)-propene



Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
	Coupling Constants (Hz)		

¹H


3.85	s	3H	a	
7.20	t	7.6	1H	f or g
7.33	t	7.6	1H	f or g
7.39	d	8.0	1H	h
7.51	d	8.3	1H	e
7.56	s (br)	1H	b	

¹⁹F

-100.63	s	1F	p	
-114.70	s	2F	n	
-119.30	s	2F	o	
-117.22	} AB	J _{AB} =215	4F	k
-120.45				
-128.22	} AB	J _{AB} =221	2F	l
-130.37				

No. 19

White Solid (*)



Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
	Coupling Constants (Hz)		

¹H


2.19	s	2H	f
3.12	s	12H	a
7.82-8.40		4H	b, c

¹⁹F

-120.14	s	4F	e	
-130.40	} AB	J _{AB} =215	4F	d
-133.89				

No. 20

2-(pyrrol-2-yl)-4,6-difluoro-s-triazine



Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
----------------------	--------------	-------------------------	--------------------	------------

¹H


6.40	dd	J _{c,d} =4.0	1H	c
7.23	m	J _{b,c} =2.3	1H	b
7.34	dd	J _{b,d} =1.6	1H	d
10.30	s (broad)		1H	a

¹⁹F

-39.68	s			e
--------	---	--	--	---

No. 21

2-(pyrrol-2-yl)-4-fluoro-6-perfluoroisopropyl-s-triazine



Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
----------------------	--------------	-------------------------	--------------------	------------

¹H

6.43	m	J _{c,d} =4.0	1H	c
7.27	m	J _{b,c} =2.4	1H	b
7.40	m	J _{b,d} =1.2	1H	d
10.62	s (br)		1H	a


¹⁹F

-39.68	s		1F	j
-74.43	s		6F	i
-184.47	s		1F	h

¹³C


90.9	d sept	211, 33		h
114.5	s			c
121.2	s			d
121.3	qd	288, 27		i
128.5	s			e
130.5	s			b
169.2	dd	22, 12		g
170.4	dd	13, 3		f
171.2	dd	229, 3		j

No. 22 **2-(pyrrol-2-yl)-4,6-perfluorodi-isopropyl-s-triazine**



Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
¹H				
6.31	dd	J _{c,d} =3.5	1H	c
7.15	m	J _{b,c} =1.7	1H	b
7.29	dd	J _{b,d} =1.5	1H	d
9.50	s (br)		1H	a
¹⁹F				
-74.42	s		6F	i
-184.72	s		1F	h
¹³C				
90.4	d sept	211, 33		h
114.3	s			c
120.8	qd	288, 27		i
121.3	s			d
128.2	s			e
130.6	s			b
165.9	dd	22, 4		g
166.5	t	3		f


No. 23 **2-(N-methylpyrrol-2-yl)-4,6-difluoro-s-triazine**



Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
¹H				
4.04	s		3H	a
6.27	dd	J _{c,d} =4.0	1H	c
7.15	m	J _{b,c} =2.4	1H	b
7.44	dd	J _{b,d} =1.8	1H	d
¹⁹F				
-45.60	s (broad)			g,h
¹³C				
39.21	s			a
111.09	s			c
123.34	s			d
128.0	s			e
136.13	s			b
172.03	dd	226, 19		g, h
172.35	t	14		f

For ¹³C spectra, it is important to have the proton decoupler on during acquisition but off during delay. This prevents artificial nuclear overhauser enhancement of peaks with low relaxation times (e.g. non-substituted aromatic carbons), hence giving greater resolution for peaks with a longer relaxation time (e.g. C-F carbons). However, a much longer acquisition time is necessary.


No. 24 **2-(N-methylpyrrol-2-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine**



Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
¹H				
4.12	s		3H	a
6.33	dd	J _{c,d} =4.2	1H	c
7.35	m	J _{b,c} =2.3	1H	b
7.52	dd	J _{b,d} =1.8	1H	d
¹⁹F				
-38.4	s		1F	j
-74.2	s		6F	i
-184.3	s		1F	h
¹³C				
39.28	s			a
90.28	d sept	211, 35		h
111.55	s			c
120.76	qd	288, 27		i
123.95	s			d
128.06	s			e
136.85	s			b
168.18	dd	21, 12		g
170.21	dd	13, 3		f
170.27	dd	228, 3		j

¹³C acquisition time 17.5 hr

No. 25 **2-(N-methylpyrrol-2-yl)-4,6-perfluorodi-isopropyl-s-triazine**




Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
¹H				
4.07	s		3H	a
6.32	dd	J _{c,d} =4.2	1H	c
7.05	m	J _{b,c} =2.3	1H	b
7.66	dd	J _{b,d} =1.9	1H	d
¹⁹F				
-74.96	s		6F	i
-185.19	s		1F	h
¹³C				
39.09	s			a
90.21	d sept	218, 33		h
111.88	s			c
120.73	qd	288, 27		i
124.75	s			d
128.12	s			e
137.67	s			b
165.55	dd	22, 3		g
166.62	t	3		f

Spectra recorded in d₆-acetone

No. 26

2-(N-methylindol-3-yl)-4,6-difluoro-s-triazine



Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
----------------------	--------------	--------------------	------------

¹H

4.1	s	3H	a
7.4	m	2H	d, e
7.7	m	1H	f
8.6	m	1H	c
8.7	s	1H	b


¹⁹F

-40.2	s		g
-------	---	--	---

Resolution of ¹H spectrum is poor due to the low solubility of the product

No. 28

2-(N-methylindol-3-yl)-4,6-perfluorodi-isopropyl-s-triazine



Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
----------------------	--------------	--------------------	------------

¹H


3.97	s	3H	a
7.43	m	2H	d, e
7.59	m	1H	f
8.44	m	1H	c
8.56	s	1H	b

¹⁹F

-74.17	s	6F	h
-184.44	s	1F	g

Resolution on ¹H spectrum poor due to the low solubility of the product

No. 27 **2-(N-méthylindol-3-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine**



140.53	s		b
168.14	m (br)		l
170.29	dm	229	k
175.02	dd	13, 3	j

<u>Chemical Shift</u> <u>(ppm)</u>	<u>Multiplicity</u> <u>Coupling Constants (Hz)</u>	<u>Relative Intensity</u>	<u>Assignment</u>
---------------------------------------	---	---------------------------	-------------------

Spectra recorded in d_6 -acetone

^1H

4.04	s		3H	a
7.36	m		2H	f, g
7.67	dd	$J_{g,h}=6$ $J_{f,h}=2$	1H	h
8.45	dd	$J_{e,f}=6.5$ $J_{e,g}=3$	1H	e
8.59	s		1H	b

^{19}F


-39.2	s		1F	k
-74.2	s		6F	n
-184.2	s		1F	m

^{13}C

34.23	s			a
90.17	d sept	210, 33		m
111.72	d	1.9		c
111.98	s			g
120.77	dq	287, 27		n
122.98	s			f
123.98	s			e
124.63	s			h
127.16	s			d
139.65	s			i

No. 29

2-(N,N-dimethylaminophenyl)-4,6-difluoro-s-triazine




Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
----------------------	--------------	--------------------	------------

^1H			
Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
3.10	s	6H	a
6.80	} AX $J_{AX}=9.0$	2H	b
8.28		2H	c
^{19}F			
Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
-40.13	s		d

No. 30

2-(4-N,N-Dimethylaminophenyl)-4-perfluoro-isopropyl-6-fluoro-s-triazine



Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
----------------------	--------------	--------------------	------------


Spectra recorded in CDCl_3

^1H			
Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
3.40	s	6H	a
6.98	} AX $J_{AX}=9.4$	2H	b
8.65		2H	c
^{19}F			
Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
-37.25	s	1F	d
-74.27	d	6F	f
-184.82	sept	1F	e

No 32


2-(N,N-diethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazine

Para isomer



Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
¹H			
1.20	t 7	6H	a
3.49	q 7	4H	b
6.80	-AX J _{AX} =9.6	2H	c
8.27		2H	d
¹⁹F			
-43.50	s	1F	e
-78.09	d 6.8	6F	g
-188.17	sept 6.8	1F	f
¹³C			
12.5	s		a
44.8	s		b
89.3	d sept 213. 33		i
111.16	s		d
119.74	qdq 288. 27		j
130.30	s		f
132.74	s		e
153.14	s		c
167.95	dd 21.9, 11.5		h
169.79	d 234		k
176.00	d 12.5		g

Ortho Isomer




Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
¹H			
1.07	t 7.1	6H	a
3.51	q 7.1	4H	b
7.16-7.48		4H	c, d, e, f
¹⁹F			
-43.50	s	1F	g
-78.03	d 6.4	6F	i
-187.09	sept 6.8	1F	h

The resonances corresponding to the ortho isomer in the ¹³C spectrum cannot be resolved due to the small concentration of this isomer. However, line broadening of the spectrum indicates the presence of the ortho isomer.

No. 33


2-(4-N,N-Diethylaminophenyl)-4,6-perfluorodiisopropyls-triazine

Para Isomer




Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
¹H				
1.18	t	7.2	6H	a
3.42	q	7.2	4H	b
6.65	} AX	J _{AX} =9.2	2H	c
8.33			2H	d
¹⁹F				
-73.15	d	7.1	12F	f
-183.78	sept	7.1	2F	e

Ortho Isomer




Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
¹H				
1.05	t	7.2	6H	a
3.42	q	7.2	4H	b
7.34-7.60			4H	c, d, e, f
¹⁹F				
-73.33	d	7.5	6F	i
-183.60	sept	7.2	1F	h

No. 34 1,8-(Bisdimethylamino)-4,5-(bisdifluorotriaz-1-yl)-naphthalene



Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
3.09	s	6H	a
7.02	AX J _{AX} =8.6		b
8.36			c
¹⁹ F			
-40.2			d


No. 35 2-(ethylphenylamino)-4,6-difluoro-s-triazine



Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
1.26	t	3H	a
4.03	q	2H	b
7.21	d	2H	c
7.45	m	3H	d, e
¹⁹ F			
-37.32	s	1F	f
-37.88	s	1F	f

No. 36

2-(ethylphenyl)amino-4-perfluoroisopropyl-s-triazine



Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
----------------------	--------------	-------------------------	--------------------	------------

¹H


1.24	t	7.2	3H	a
4.76	q	7.2	2H	b
7.29	t	7.2	2H	d
7.48	m		3H	c, e

¹⁹F

-39.9	s		1F	f
-74.45	s		6F	h
-184.65	s		1F	g

No. 37

2-(methyl-(2-methoxy-phenyl)amino)-4,6-difluoro-s-triazine



Chemical Shift (ppm)	Multiplicity	Coupling Constants (Hz)	Relative Intensity	Assignment
----------------------	--------------	-------------------------	--------------------	------------

Spectra recorded in d₆-acetone

¹H


3.47	s		3H	a or f
3.84	s		3H	a or f
7.06	t	7.0	1H	c or d
7.18	d	8.1	1H	b or e
7.33	d	7.5	1H	b or e
7.39	t	7.6	1H	c or d

¹⁹F

-40.33	s		1F	g
-41.14	s		1F	g

No. 38

2-(methyl(N,N-dimethylamino)phenyl)-4,6-difluoro-s-triazine



Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
	Coupling Constants (Hz)		

Spectra recorded in d_6 -acetone

^1H


2.97	s	6H	d
3.50	s	3H	a
6.78	} AB $J_{AB}=8.9$	2H	b
7.17		2H	c

^{19}F

-40.6	s	1F	e
-41.1	s	1F	e

No. 39

Pyridine Salt



Chemical Shift (ppm)	Multiplicity	Relative Intensity	Assignment
	Coupling Constants (Hz)		

^1H

3.25	s	6H	e
6.84	ddd $J_{c,d}=6.4$	1H	c
7.06	ddd $J_{a,b}=9$ $J_{a,c}=0.8$ $J_{a,d}=0.8$	1H	a
7.91	ddd $J_{b,c}=7$ $J_{b,d}=1.9$	1H	b
8.06	ddd	1H	d

^{19}F


-74.95	s	6F	g
-185.30	s	1F	f


APPENDIX TWO**INFRA RED SPECTRA**




1. 1,8-(Bisdimethylamino)-naphthalene Hydrogen Fluoride Complex (PS/HF) (1)(Nujol mull)
2. Triethylamine Hydrogen Fluoride Complex (24)
3. Tributylamine Hydrogen Fluoride Complex (25)
4. Trihexylamine Hydrogen Fluoride Complex (26)
5. Trioctylamine Hydrogen Fluoride Complex (27)
6. Tridodecylamine Hydrogen Fluoride Complex (28)
7. 2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (35)
8. N-Methyl-2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (36)
9. N-Ethyl-2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (37)
10. N-Allyl-2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (38)
11. N-Benzyl-2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (39)
12. Tetramethylguanidine Hydrogen Fluoride Complex (40)
13. 1,1-Bistrifluoromethyl-6,7-bisdimethylamino-2,3-tetrafluoro-ethano-[1H]-phenalene (4)
14. Unidentified Red Solid (5)
15. Spiro[octafluorocyclopentane-1,1'-(6,7-bisdimethylamino)-2',3'-tetrafluoroethano-[1H]-phenalene] (44)
16. (7,8)-(9,10)-dihexafluoropropano-3,4-bisdimethylamino-cyclohepta-[d,e]-naphthalene (56)
17. Spiro[octafluorocyclopentane-1,1'-(6,7-bisdimethylamino)-2',3'-tetrafluoropropan-2-one-[1H]-phenalene] (60)
18. 7,8-propano-9,10-propan-2-one-cyclohepta-[d,e]-naphthalene (61)
19. Spiro[hexafluorocyclobutane-3,1'-1,2-tetrafluoroethano-1-fluoro-3-(4''-N,N-dimethylaminophenyl)-propene] (63)
20. Spiro[hexafluorocyclobutane-3,1'-1,2-tetrafluoroethano-1-fluoro-3-(N-methylindol-3''-yl)-propene] (65)
21. White Solid (66)
22. 2-(pyrrol-2-yl)-4,6-difluoro-s-triazine (67)
23. 2-(pyrrol-2-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine (68)
24. 2-(pyrrol-2-yl)-4,6-perfluorodi-isopropyl-s-triazine (69)
25. 2-(N-methylpyrrol-2-yl)-4,6-difluoro-s-triazine (70)
26. 2-(N-methylpyrrol-2-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine (71)
27. 2-(N-methylpyrrol-2-yl)-4,6-perfluorodi-isopropyl-s-triazine (72)
28. 2-(N-methylindol-3-yl)-4,6-difluoro-s-triazine (73)
29. 2-(N-methylindol-3-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine (74)

30. 2-(N-methylindol-3-yl)-4,6-perfluorodi-isopropyl-s-triazine (75)
31. 6-(N-methylindol-3-yl)-2,4,5-trifluoro-pyrimidine (76)
32. 2-(4-N,N-dimethylaminophenyl)-4,6-difluoro-s-triazine (77)
33. 2-(4-N,N-dimethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazine (78)
34. 2-(4-N,N-dimethylaminophenyl)-4,6-perfluorodi-isopropyl-s-triazine (81A) and 2-(2-N,N-dimethylaminophenyl)-4,6-perfluorodi-isopropyl-s-triazine (81B)
35. 2-(4-N,N-diethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazine (79A) and 2-(2-N,N-diethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazine (79B)
36. 2-(4-N,N-diethylaminophenyl)-4,6-perfluorodi-isopropyl-s-triazine (82A) and 2-(2-N,N-diethylaminophenyl)-4,6-perfluorodi-isopropyl-s-triazine (82B)
37. 1,8-(Bisdimethylamino)-4,5-(bisdifluoro-s-triaz-2-yl)-naphthalene (83)
38. 2-(ethylphenylamino)-4,6-difluoro-s-triazine (84)
39. 2-(ethylphenylamino)-4-perfluoroisopropyl-6-fluoro-s-triazine (80)
40. 2-methyl-(2-methoxyphenyl)-amino-4,6-difluoro-s-triazine (85)
41. 2-methyl-(4-N,N-dimethylaminophenyl)-amino-4,6-difluoro-s-triazine (86)
42. Pyridinium Salt (87)

All solids were recorded as KBr discs unless otherwise stated. All liquids were run as thin films between KBr plates









25 3 4 5 Microns 6 7 10 14 20 40


No. 13




No. 14



No. 15



No. 16



No. 17



4000 3500 3000 2500 2000 1800 1600 1400 1200 1000 800 600 400 250

25 3 4 5 Microns 6 7 10 14 20 40


No. 18

No. 19

No. 20

No. 21

4000 3500 3000 2500 2000 1800 1600 1400 1200 1000 800 600 400 250
Wavenumber (cm⁻¹)



25 3 4 5 Microns 6 7 10 14 20 40


No. 22




No. 23



No. 24




No. 25



No. 26




4000 3500 3000 2500 2000 1800 1600 1400 1200 1000 800 600 400 250
Wavenumber (cm⁻¹)




25 3 4 5 Microns 6 7 10 14 20 40




No. 32



No. 33



No. 34




No. 35



No. 36

4000 3500 3000 2500 2000 1800 1600 1400 1200 1000 800 600 400 250
Wavenumber (cm⁻¹)



APPENDIX THREE**MASS SPECTRA**



1. 1,8-(Bisdimethylamino)-naphthalene Hydrogen Fluoride Complex (PS/HF) (1)
2. Triethylamine Hydrogen Fluoride Complex (24)
3. Tributylamine Hydrogen Fluoride Complex (25)
4. Trihexylamine Hydrogen Fluoride Complex (26)
5. Trioctylamine Hydrogen Fluoride Complex (27)
6. Tridodecylamine Hydrogen Fluoride Complex (28)
7. 2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (35)
8. N-Methyl-2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (36)
9. N-Ethyl-2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (37)
10. N-Allyl-2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (38)
11. N-Benzyl-2,2,6,6-Tetramethylpiperidine Hydrogen Fluoride Complex (39)
12. Tetramethylguanidine Hydrogen Fluoride Complex (40)
13. 1,1-Bistrifluoromethyl-6,7-bisdimethylamino-2,3-tetrafluoro-ethano-[1H]-phenalene (4)
14. Unidentified Red Solid (5)
15. Spiro[octafluorocyclopentane-1,1'-(6,7-bisdimethylamino)-2',3'-tetrafluoroethano-[1H]-phenalene] (44)
16. (7,8)-(9,10)-dihexafluoropropano-3,4-bisdimethylamino-cyclohepta-[d,e]-naphthalene (56)
17. Spiro[octafluorocyclopentane-1,1'-(6,7-bisdimethylamino)-2',3'-tetrafluoropropan-2-one-[1H]-phenalene] (60)
18. 7,8-propano-9,10-propan-2-one-cyclohepta-[d,e]-naphthalene (61)
19. Spiro[hexafluorocyclobutane-3,1'-1,2-tetrafluoroethano-1-fluoro-3-(4''-N,N-dimethylaminophenyl)-propene] (63)
20. Spiro[hexafluorocyclobutane-3,1'-1,2-tetrafluoroethano-1-fluoro-3-(N-methylindol-3''-yl)-propene] (65)
21. 2-(pyrrol-2-yl)-4,6-difluoro-s-triazine (67)
22. 2-(pyrrol-2-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine (68)
23. 2-(pyrrol-2-yl)-4,6-perfluorodi-isopropyl-s-triazine (69)
24. 2-(N-methylpyrrol-2-yl)-4,6-difluoro-s-triazine (70)
25. 2-(N-methylpyrrol-2-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine (71)
26. 2-(N-methylpyrrol-2-yl)-4,6-perfluorodi-isopropyl-s-triazine (72)
27. 2-(N-methylindol-3-yl)-4,6-difluoro-s-triazine (73)
28. 2-(N-methylindol-3-yl)-4-perfluoroisopropyl-6-fluoro-s-triazine (74)
29. 2-(N-methylindol-3-yl)-4,6-perfluorodi-isopropyl-s-triazine (75)
30. 6-(N-methylindol-3-yl)-2,4,5-trifluoro-pyrimidine (76)

31. 2-(4-N,N-dimethylaminophenyl)-4,6-difluoro-s-triazine (77)
32. 2-(4-N,N-dimethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazin(78)
33. 2-(4-N,N-dimethylaminophenyl)-4,6-perfluorodi-isopropyl-s-triazine (81A) and 2-(2-N,N-dimethylaminophenyl)-4,6-perfluorodi-isopropyl-s-triazine (81B)
34. 2-(4-N,N-diethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazine (79A) and 2-(2-N,N-diethylaminophenyl)-4-perfluoroisopropyl-6-fluoro-s-triazine (79B)
35. 2-(4-N,N-diethylaminophenyl)-4,6-perfluorodi-isopropyl-s-triazine (82A) and 2-(2-N,N-diethylaminophenyl)-4,6-perfluorodi-isopropyl-s-triazine (82B)
36. 1,8-(Bisdimethylamino)-4,5-(bisdifluoro-s-triaz-2-yl)-naphthalene (83)
37. 2-(ethylphenylamino)-4,6-difluoro-s-triazine (84)
38. 2-(ethylphenylamino)-4-perfluoroisopropyl-6-fluoro-s-triazine (80)
39. 2-methyl-(2-methoxyphenyl)-amino-4,6-difluoro-s-triazine (85)
40. 2-methyl-(4-N,N-dimethylaminophenyl)-amino-4,6-difluoro-s-triazine (86)
41. Pyridinium Salt (87)

No. 1

FAB⁺

GS1#4 x1 Bgd=1 12-OCT-89 15 3-0 02 04 70E FB+
 BpM=8 1=3.0v Hm=964 TIC=107052000 Acnt: Sys: FAB
 G.SANDFORD *x1.0 PT= 0⁰ Cal: GLYC120CT




Mass	% Base	Mass	% Base
167.09	4.11	196.61	0.24
167.21	0.29	197.13	0.19
167.58	0.31	197.33	0.53
167.61	0.28	197.51	0.31
168.09	30.07	198.11	4.65
169.09	12.20	198.50	0.23
169.68	0.40	199.12	9.53
170.10	11.82	199.36	0.33
170.45	0.34	200.12	15.61
170.88	0.25	200.27	0.49
170.93	0.28	201.13	7.74
171.11	6.37	202.11	1.73
171.45	0.36	203.03	0.34
172.12	2.79	208.07	0.99
173.13	2.15	209.08	0.34
174.09	0.44	210.09	0.49
174.92	0.41	211.10	1.49
178.98	0.39	211.33	0.33
180.06	1.32	212.05	0.44
181.09	5.05	212.14	0.51
182.09	3.28	213.13	4.16
182.31	1.29	213.59	0.69
183.10	10.11	214.15	7.57
183.65	0.23	215.15	100.00
184.10	6.26	216.16	17.55
185.11	3.38	217.15	3.80
185.56	0.29	218.08	1.89
185.82	0.66	219.02	0.39
186.11	2.06	221.10	0.75
		222.08	0.74
		223.15	0.87
		225.04	0.63

No. 2

EI⁺

GS6385* x1 Bgd=1 20-NOV-89 14:20:00:40 70E EI+
 BpM=0 I=2.3v Ha=215 TIC=62364800 Acnt: Sys: ACE
 G. SARNOFORD PT= 0° Cal: PFK269
 *x1.0


EI⁺ Data

Mass	% Base	Mass	% Base
23.41	0.31	82.82	1.60
25.94	3.08	83.93	0.60
26.95	16.50	84.81	81.17
27.90	2.92	84.93	0.23
27.94	11.18 F	85.82	4.21
27.95	3.56 F	85.94	100.00
28.96	16.44	86.81	2.52
29.95	28.87	86.94	5.85
30.96	0.23	90.89	0.23
32.90	0.94	99.94	6.56
38.93	1.30	100.95	23.58
39.93	0.77	101.95	2.69
40.94	2.77	103.80	0.90
41.94	14.30	104.80	0.35
42.39	0.22	148.81	0.83
42.92	0.44		
42.95	1.76		
43.89	0.40		
43.95	10.73		
46.87	4.18		
51.91	0.21		
53.92	1.54		
54.93	1.03		
55.93	7.65		
56.94	2.07		
57.94	25.66		
58.94	0.78		
65.83	0.98		
67.91	0.31		
68.92	0.32		
69.92	3.62		
70.92	1.37		
71.93	3.55		
81.84	2.50		

No. 3

EI⁺

GS58X05* x1 8gd=1 7-AUG-89 14:20:01:43 78E EI*
 BpM=0 I=396uv Hw=166 TIC=10252000 Acnt: Sys:ACE
 G. SANDFORD PT= 8° CaL: PFK7AUG
 *X1.0




Mass	% Base
40.88	24.03
41.88	15.07
42.90	5.42
43.90	18.49
52.94	0.69
53.95	0.62
54.97	4.31
55.97	3.38
56.99	7.92
57.99	15.46
70.05	2.50
84.09	3.00
84.98	2.58
86.10	3.19
98.11	4.50
100.12	50.37
101.13	2.96
128.12	0.65
142.17	100.00
*143.18	8.73
*184.20	7.77
185.21	6.81
186.22	8.04

No. 4

EI⁺

GS4BX#5+ x1 Bgd=1 7-AUG-89 14:20:00:30 78E EI+
 BpM=0 I=7.2v Hn=312 TIC=127377000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK7AUG

*X3.0
 X3.0*




EI⁺ Data

Mass	% Base		
40.87	9.19	140.12	0.24
41.87	4.06	142.14	0.23
42.89	17.32	154.13	0.78
43.89	8.62	156.14	0.21
44.90	0.23	168.12	0.32
52.93	0.47	184.16	1.08
53.94	0.29	185.17	0.27
54.96	4.43	186.18	0.30
55.96	1.59	196.18	0.70
56.97	2.47	197.18	0.26
57.98	8.24	198.19	100.00
58.99	0.27	199.20	14.95
67.01	0.35	200.20	1.05
68.01	0.28	212.20	0.31
69.04	0.61	224.22	0.42
70.03	1.75	240.24	0.49
71.04	0.50	254.26	0.39
72.05	0.26	268.28	3.22
81.06	0.24	269.28	2.83
82.06	0.37	270.29	4.45
83.07	0.49	271.29	0.78
84.07	3.53		
84.96	2.53		
85.08	0.55		
96.07	0.23		
98.09	4.59		
99.08	0.35		
100.10	0.36		
112.10	1.05		
114.10	3.47		
115.10	0.28		
126.08	2.52		
127.09	0.26		
128.09	23.13		
129.10	2.10		

No. 5

EI+

GSN105+ xl 0gd=1 26-SEP-91 09:34:00:50 70E EI+
 BpM=43 I=2.2v Hw=494 TIC=206291008 Rcnt: Sys: ACE
 G. SANDFORD PT= 0° Cal: PFK25SEP
 *X1.0




Mass	% Base	Mass	% Base
119.03	14.66	166.18	1.34
120.05	54.90	167.14	0.58
121.06	16.40	168.14	3.56
122.08	1.78	169.04	10.30
123.11	0.53	170.14	0.60
124.10	0.61	171.16	0.39
125.14	0.57	243.05	2.03
126.16	1.78	247.17	0.38
127.12	1.34	252.33	1.94
128.12	2.05	253.34	0.41
129.11	1.20	254.35	28.16
130.12	0.38	255.31	5.93
131.03	14.69	256.36	0.63
132.04	0.56	260.15	1.04
135.11	0.36	267.06	0.54
137.08	0.51	268.36	0.41
138.14	0.42	269.05	1.81
139.11	2.03	278.16	0.71
140.16	4.99	280.37	1.10
141.12	3.58	281.06	3.56
142.19	16.20		
143.13	2.05		
145.16	0.46		
149.07	1.23		
150.04	0.74		
151.05	0.96		
152.08	23.17		
153.10	3.16		
154.18	1.78		
155.10	1.91		
156.20	4.49		
157.16	0.70		
162.03	1.78		
163.05	0.74		
165.15	0.54		

No. 6

EI⁺

GS6409* x1 Bgd=1 20-NOV-89 14:50:00:56 70E EI⁺
 ApM=0 I=100 Hw=684 TIC=764024000 Rcnt: Sys:ACE
 G. SANDFORD PT= 0° Cal: PFK269
 *X1.0



EI⁺ Data

Mass	% Base	Mass	% Base
353.14	2.04	478.31	1.64
354.15	2.19	479.31	0.61
355.16	0.55	492.33	1.00
362.12	0.51	493.33	0.38
363.13	0.28	506.35	0.61
364.13	10.32 F	507.35	0.24
365.09	3.60 F	518.35	1.07
366.14	100.00 FO	519.36	1.17
367.16	73.26 F	520.36	18.16
368.16	11.24	521.37	8.44
369.17	1.26	522.39	22.54
378.16	0.33	523.39	8.95
380.17	1.08	524.40	1.85
381.18	0.36	525.40	0.21
382.17	0.54		
392.18	2.94		
393.17	1.00		
394.19	0.90		
395.19	0.25		
406.20	0.29		
408.21	0.72		
409.22	0.26		
420.22	0.25		
422.23	0.79		
423.24	0.26		
434.23	0.28		
436.25	0.82		
437.25	0.27		
448.26	0.31		
450.27	1.05		
451.28	0.30		
462.27	0.24		
464.29	1.57		
465.30	0.58		
476.30	0.28		

No.7

Cl⁺

GS3648+ xl 8gd=6 27-APR-89 14:30:00:51 78E C1+
 SpM=0 l=10v Mw=539 TIC=183458888 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cat: PFK26APR





Mass	% Base	Mass	% Base
44.04	1.10	140.14	0.27
58.06	12.94	140.47	0.03
70.06	4.95	140.51	0.03
98.09	1.36	140.55	0.04
113.11	1.01	140.60	0.05
120.08	0.05	140.67	0.03
121.11	0.04	142.24	100.00 FO
122.07	0.19	143.17	31.60 F
123.10	0.11	143.69	0.04
124.01	0.03	143.74	0.04
124.11	0.27	143.76	0.04
125.12	0.23	143.83	0.05
125.61	0.03	143.90	0.04
125.68	0.04	143.93	0.05
125.73	0.04	144.17	1.62
126.13	21.79		
126.42	0.04		
127.14	2.03		
128.14	0.18		
129.10	0.04		
130.08	0.05		
130.12	0.04		
131.08	0.07		
132.07	0.02		
132.11	0.04		
133.09	0.20		
134.07	0.06		
134.12	0.04		
135.09	0.07		
136.08	0.04		
136.11	0.04		
137.06	0.05		
137.08	0.04		
137.14	0.06		
138.13	0.79		
139.03	0.07		
139.13	0.19		

No.8

Cl⁺

GS41023• x1 Bgd=6 0-JUN-89 16:30:01:52 70E Cl•
 BpM=0 I=10v Hm=158 TIC=103024000 Rcnt: Sys:RCE
 G.SANDFORD PT= 0° Cal: PFK19MAY
 *x1.0





Mass	% Base
56 03	1 96
58 05	0 92
70 04	0 43
72 05	2 91
84 05	3 39
124 06	0 31
126 07	0 52
140 09	22 20
141 08	2 32
142 09	1 26
154 08	0 66
156 13	100 00 FO
157 10	13 02 F
158 11	0 69

No. 9

G552826 x1 Bgd=6 15-AUG-89 11:50:02:04 70E Cl+
 Bpm=8 I=2.0v Nm=440 TIC=73872800 Acnt: Sys:RCE
 G.SANDFORD *x1.0 PT= 0° Cal:PFK7RUG

Cl+





Mass	% Base	Mass	% Base
121.14	0.43	162.21	0.48
122.15	0.65	163.22	0.52
123.15	0.71	164.21	0.53
124.15	0.83	165.20	0.27
125.17	0.44	165.27	0.20
126.18	5.59	166.23	0.47
127.18	0.60	167.21	0.22
128.19	0.48	168.25	0.57
132.15	0.31	169.26	0.84
133.15	0.41	170.28	100.00
134.16	0.41	171.28	12.32
135.17	0.35	172.26	0.97
136.17	0.42	173.20	0.28
137.19	0.38	174.21	0.24
138.21	0.98	175.22	0.33
139.20	0.39	176.23	0.28
140.24	0.59	177.20	0.29
141.18	0.22	178.24	0.31
142.26	16.61	179.25	0.24
143.27	1.47	180.26	0.28
144.21	0.30	182.26	0.32
145.17	0.28	183.23	0.22
146.20	0.29	184.25	0.23
147.19	0.54		
148.20	0.36		
149.21	0.40		
150.22	0.36		
151.21	0.28		
152.22	0.52		
153.23	0.46		
154.26	33.71		
155.26	3.88		
156.26	0.69		
158.20	0.26		
159.20	0.23		
160.20	0.24		
161.20	0.34		

No. 10

EI⁺

GSG6814 x1 Bgd=1 28-NOV-89 16:20:01:15 78E EI⁺
 BpM=0 I=10v Hm=367 TIC=452894816 Acnt: Sys:ACE
 6. SANDFORD PT= 0° Cal: PFK269
 *x1-u





Mass	% Base				
20.95	1.09	77.90	0.25	180.97	16.41
24.94	0.32	78.91	1.77	181.98	20.27
25.94	2.70	79.91	2.27	182.98	2.67
26.95	12.93	80.92	4.79	366.19	0.36
27.95	4.73 F	81.92	17.97		
27.96	4.15 F	82.93	2.42		
28.97	11.69	83.94	2.47		
29.96	8.57	92.92	1.36		
36.92	0.65	93.91	2.62		
37.93	1.37	94.92	1.81		
38.94	24.74	95.93	6.97		
39.94	4.43	96.93	5.26		
40.95	71.83	97.94	32.55		
41.95	22.71	98.94	2.20		
42.96	4.47	107.92	3.23		
43.96	1.14	108.93	5.30		
44.94	0.47	109.93	72.91		
49.91	0.49	110.93	6.55		
50.92	0.96	111.94	1.05		
51.92	0.57	119.90	0.23		
52.93	5.72	121.91	1.73		
53.93	4.40	122.92	0.68		
54.94	13.23	123.93	2.40		
55.94	13.73	124.93	2.79		
56.95	2.31	125.94	4.20		
57.95	11.95	149.93	7.55		
58.95	0.37	150.93	0.90		
64.91	1.10	151.95	1.16		
65.91	0.67	153.96	2.27		
66.92	6.54	154.96	0.33		
67.91	3.77	163.94	1.23		
68.93	32.98	164.94	5.69 F		
69.93	6.47	165.97	100.00 F0		
70.93	1.11	166.96	40.71 F		
71.94	1.00	167.96	2.55		
76.90	0.86	179.96	3.98		

No.11

Cl⁺

GS6510* x1 Bgd=6 20-NOV-89 11:40:00:53 78E Cl⁺
 SpM=0 I=10v Hm=323 TIC=193914000 Acnt: Sys:ACE
 G. SANDFORD PT= 0° Cal: PFK269
 *x1.0


Cl⁺ Data

Mass	% Base	Mass	% Base
29.98	0.24	159.99	0.53
31.98	1.16	162.85	0.23
32.99	0.45	198.01	0.27
35.01	100.00 0	203.04	0.21
36.00	2.53	216.05	11.14
43.98	1.12	217.06	1.86
44.98	0.25	230.07	0.47
46.00	0.24	232.12	100.00 0
52.01	1.12	233.10	35.66
57.98	1.68	234.10	3.23
59.96	0.45	279.05	0.60
60.95	0.75	322.17	0.25
69.97	0.29		
71.98	0.30		
73.96	0.43		
76.98	0.95		
77.97	1.02		
79.95	0.20		
81.97	0.32		
87.98	0.56		
90.96	1.62		
98.98	0.21		
101.99	0.56		
105.00	0.37		
105.96	1.11		
107.98	5.75		
108.98	0.51		
112.99	0.42		
113.98	0.27		
115.99	0.31		
119.00	0.26		
126.01	1.51		
138.98	0.21		
142.04	2.75		
145.98	0.28		
147.99	0.70		

No.12

Cl⁺

GSAG010 x1 Bgd=6 25-SEP-91 12:19:01:13 78E Cl⁺
 BpM=0 I=10v Ha=448 TIC=315950016 Rcnt: Sys:ACE
 G.SANDFORD PT= 0° Cat:PFK25SEP





Mass	% Base		
43.06	0.47	94.11	0.39
44.07	26.97	96.12	0.45
45.08	5.00	97.13	0.38
46.09	93.00	98.14	0.50
47.09	2.56	99.13	0.39
49.10	0.39	100.14	1.36
52.12	1.64	102.15	6.35
56.09	0.40	103.16	0.47
57.08	2.34	106.14	0.38
58.10	2.11	108.13	0.39
59.10	1.86	109.14	0.39
60.08	1.13 F	110.14	0.41
60.12	1.19 F	112.15	0.39
61.07	1.02 F	114.17	0.45
61.12	0.54 F	115.11	3.88 F
63.13	0.98	116.20	100.00 FO
69.09	0.80	117.18	34.42 F
70.10	0.43	118.18	1.17
71.10	5.48	119.17	0.32
72.11	9.07	126.17	0.34
73.12	16.54	131.19	0.39
74.11	1.84		
77.11	0.97		
78.11	0.52		
82.11	0.43		
83.10	0.43		
84.12	0.49		
85.12	0.40		
86.13	1.54		
87.14	0.41		
88.13	12.58		
89.12	5.73		
90.12	0.37		
91.13	0.39		
93.10	0.45		

No.13

M.Wt. 486

CI⁺/CI⁻


GSBSA#4* x1 Bgd=1 2-MAY-98 15:15:00:36 78E EI+
 BpM=0 I=447mv Hn=488 TIC=41488888 Acnt: Sys:ACE
 G.SANDYFORD PT= 0° Cal: PFK24APR

HMR
MASS

GSBSA#23* x1 Bgd=6 2-MAY-98 15:15:00:51 78E CI+
 BpM=0 I=10v Hn=674 TIC=388712888 Acnt: Sys:ACE
 G.SANDYFORD PT= 0° Cal: PFK24APR

HMR
MASS

GSBSA#45* x1 Bgd=41 2-MAY-98 15:15:00:19 78E CI-
 BpM=0 I=2.7v Hn=502 TIC=35388888 Acnt: Sys:ACE
 G.SANDYFORD PT= 0° Cal: PFK24APR

HMR
MASSCI⁺ Data

Mass	% Base
397.20	1.42
398.17	3.70 F
399.18	10.05 F
400.19	2.18
401.20	0.57
402.21	0.80
403.17	0.79
404.17	0.82
408.21	0.73
411.25	0.73
412.25	0.73
413.17	0.72
414.21	0.72
417.18	1.59
418.19	1.26
419.19	3.85
420.20	1.09
464.84	1.19 F
467.14	78.65 F
468.16	21.04 F
469.17	3.24 F
470.15	1.04
474.21	0.72
477.20	0.72
485.17	0.76
486.14	1.17 F
487.14	20.32 F
488.16	4.46 F
489.16	1.12
498.21	0.70


No. 14

M.Wt. 464

EI⁺/CI⁺/CI⁻

GS858033* x1 Bgd=32 9-MAY-90 12:10:00:32 70E EI⁺
 BpM=0 I=3.5v Hm=862 TIC=125970000 Acnt: Sys: ACE
 G.SANDFORD PT= 0° Cal: PFK24RPR

Unidentified Red Solid



GS858038* x1 Bgd=36 9-MAY-90 12:10:00:52 70E CI⁻
 BpM=0 I=10v Hm=675 TIC=113343000 Acnt: Sys: ACE
 G.SANDFORD PT= 0° Cal: PFK24RPR



GS858042* x1 Bgd=41 9-MAY-90 12:10:00:00 70E CI⁻
 BpM=0 I=2.7v Hm=613 TIC=29417000 Acnt: Sys: ACE
 G.SANDFORD PT= 0° Cal: PFK24RPR

EI⁺ Data

Mass	% Base
41.99	5.09
44.01	4.95
58.08	22.98
59.09	2.72
108.24	2.47
121.24	6.03
122.25	9.69
123.26	2.90
124.27	7.88
135.25	5.48
136.26	4.96
137.27	6.07
150.24	4.96
151.26	5.00
152.26	4.69
164.27	3.47
165.27	3.13
167.32	4.19
176.31	6.78
177.32	2.35
179.28	2.78
195.32	100.00
196.32	22.62
197.32	2.67
145.33	2.59
150.21	0.33
155.35	0.07
156.37	0.07
157.34	0.22
158.34	0.23
159.35	0.08
163.31	2.32
163.74	0.05
164.32	82.73
164.86	0.06
165.33	46.51
166.33	9.10
167.34	0.96
168.34	0.08


No. 15

M.Wt. 548

Cl⁺/Cl⁻


GSEG#25* x1 Bgd=22 23-JAN-91 14:47:08:02:01 70E Cl-
 BpM=0 I=10v Hw=565 TIC=160509000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK23JAN

HMR:
 MASS:



GSEG#20* x1 Bgd=11 23-JAN-91 14:47:08:02:13 70E Cl-
 BpM=0 I=2.6v Hw=551 TIC=42472000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK23JAN

HMR:
 MASS:

Cl⁺ Data

Mass	% Base		
43.95	3.69	530.23	100.00 F0
44.96	0.67	531.12	30.69 F
45.98	3.52	532.28	5.15 F
58.06	10.89	534.09	3.22 F
59.06	0.47	535.38	0.80 F
71.14	1.26	549.22	7.85 F
72.15	0.39	550.20	36.89 F
352.82	0.60	551.20	9.39 F
353.84	0.44	552.29	1.27 F
367.81	0.44		
368.62	0.39		
391.67	0.41		
403.72	0.39		
404.73	0.39		
417.74	0.39		
426.77	0.35		
433.84	0.40		
471.93	0.39		
484.92	0.39		
485.92	2.34		
486.93	0.65		
490.95	0.39		
491.99	0.39		
502.95	0.61		
503.97	0.39		
504.97	1.23		
505.98	0.39		
506.99	0.31		
510.00	2.91 F		
511.01	0.81 F		
512.05	2.34 F		
513.05	0.59		
514.01	0.92		
515.00	0.30		
516.00	0.39		
517.06	0.78		

No.16



M.Wt. 560

EI⁺/CI⁺/CI⁻

GSB112049* x1 Bgd=20 22-JUN-90 13:47:08:03:35 70E
 BpM=0 I=7.1v Hm=769 TIC=207481000

EI⁺
 Acnt: PT= 0°
 Sys: ACE
 Cal: PFK1 JUNE


HMR
 MASS



GSB112053* x1 Bgd=21 22-JUN-90 13:47:08:03:51 70E
 BpM=0 I=2.1v Hm=564 TIC=47320000

CI⁺
 Acnt: PT= 0°
 Sys: ACE
 Cal: PFK1 JUNE ,


HMR
 MASS



GSB112055* x1 Bgd=41 22-JUN-90 13:47:08:04:07 70E
 BpM=0 I=5.6v Hm=563 TIC=49093000

CI⁻
 Acnt: PT= 0°
 Sys: ACE
 Cal: PFK1 JUNE





HMR
 MASS

EI⁺ Data

Mass	% Base		
41.93	7.45	546.29	1.62
42.95	1.35	547.30	0.16
43.95	31.78	557.71	0.04
44.97	7.97	558.01	0.05
45.98	7.41	558.11	0.10
57.01	1.62	558.27	0.04
58.02	90.75	560.12	64.24 F
59.02	3.05	560.37	100.00 F
70.06	2.42	561.32	28.08 F
71.07	0.78	562.33	4.58
214.14	8.08		
215.15	1.66		
514.20	10.04		
515.10	4.47 F		
515.29	4.95 F		
516.22	7.49		
517.22	1.82		
518.23	0.79		
519.15	0.16		
519.33	0.15		
522.29	0.24		
523.24	0.07		
525.13	0.07		
525.31	0.19		
527.20	0.17		
528.20	2.34		
529.21	2.57		
530.20	1.05		
531.24	0.24		
538.29	0.15		
539.25	0.17		
540.35	0.11		
541.29	4.91		
542.31	1.37		
543.28	0.53		
544.29	0.22		
545.27	4.43		

No.17

M.Wt. 576

EI⁺/CI⁺/CI⁻CI⁻ Data

Mass	% Base
40.95	0.49
41.93	1.21
43.94	0.47
55.97	3.49
56.96	1.12
57.95	1.17
71.95	0.53
85.95	0.42
126.76	100.00 0
127.88	0.42
147.82	0.51
169.81	0.34
419.62	0.56
421.65	0.32
431.62	0.36
530.52	0.51
532.53	0.55
533.57	0.34
535.58	1.86
536.54	2.39 F
537.55	5.57 F
538.54	1.89
539.56	1.36
555.54	10.10
556.54	3.25
557.55	4.24
558.55	0.84
559.52	1.07
560.48	0.51
561.51	1.03
575.52	71.71 F
576.54	17.57 F
577.54	2.93
578.54	0.50
591.56	0.32


No.18

M.Wt. 538

EI⁺/CI⁺/CI⁻


GS8113C029_o x1 Bgd=26 14-FEB-91 11:51:0:02:48 78E
 BpM=8 I=10v Hw=664 TIC=174747888
 G.SANDFORD

Cl⁻
 Rcnt: Sys:ACE
 PT= 0⁰ Cal: PFK13F1




GS8113C035_o x1 Bgd=31 14-FEB-91 11:51:0:03:08 78E
 BpM=8 I=1.2v Hw=576 TIC=63826888
 G.SANDFORD

EI⁺
 Rcnt: Sys:ACE
 PT= 0⁰ Cal: PFK13FEB



GS8113C038_o x1 Bgd=36 14-FEB-91 11:51:0:03:23 78E
 BpM=8 I=7.1v Hw=589 TIC=83859888
 G.SANDFORD

Cl⁻
 Rcnt: Sys:ACE
 PT= 0⁰ Cal: PFK13FEB

CI⁺ Data



Mass	% Base
43.89	7.11
44.89	0.80
45.90	8.41
48.90	0.48
51.91	0.55
57.87	1.68
59.88	0.55
69.84	0.31
445.10	0.55
463.11	0.45
465.11	0.58
481.05	3.78 F
482.59	1.09 F
492.04	0.40
493.05	0.55
501.13	1.64
518.23	1.09 F
521.12	1.74 F
523.98	1.64 F
538.08	3.00 F
539.13	100.00 F
540.13	27.20 F
541.14	4.25 F
542.82	0.55 F
577.16	0.55

No. 19


M.Wt. 425

EI⁺/CI⁺/CI⁻


GSB10#20* x1 Bgd=10 19-JUN-90 10:13:01:40 70E EI⁺
 BpM=0 I=3.0v Hm=452 TIC=90731000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK1JUNE



GSB10#25* x1 Bgd=6 19-JUN-90 10:13:02:00 70E CI⁻
 BpM=0 I=4.1v Hm=452 TIC=55909000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK1JUNE >



GSB10#30* x1 Bgd=11 19-JUN-90 10:13:02:20 70E CI⁻
 BpM=0 I=10v Hm=445 TIC=154423000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK1JUNE

EI⁺ Data


Mass	% Base	Mass	% Base
41.97	1.20	404.42	1.52
43.99	1.16	405.44	1.30
97.40	93.87	406.44	5.64
170.22	3.59	407.45	1.07
224.26	0.92	410.42	1.94
225.26	1.34	422.44	0.82
231.22	2.57	423.44	0.58
232.26	0.72	424.44	29.80
256.30	0.81	425.45	100.00
268.28	0.89	426.46	20.55
274.29	2.03	427.47	1.98
275.30	4.18	451.52	0.77
281.26	6.41		
282.31	19.95		
283.32	2.65		
293.28	0.55		
304.32	0.93		
305.33	1.18		
306.34	1.44		
309.31	0.54		
310.32	2.59		
311.31	1.04		
312.30	0.86		
323.31	0.50		
324.32	29.64		
325.33	28.28		
326.33	4.05		
331.32	0.84		
336.37	0.59		
354.37	1.36		
355.38	0.92		
356.39	2.58		
361.34	1.07		
374.40	4.46		
375.41	2.23		
381.37	0.70		
386.42	0.55		

No. 20

M.Wt. 435

EI⁺/CI⁺/CI⁻


GSD19020+ x1 Bgd=1 17-JAN-91 16:36:01:48 70E EI+
 BpM=0 I=2.6v Mn=456 TIC=351520000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK17JAN



GSD19022+ x1 Bgd=10 17-JAN-91 16:36:01:48 70E CI+
 BpM=0 I=10v Mn=457 TIC=123504000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK17JAN



GSD19020+ x1 Bgd=11 17-JAN-91 16:36:02:12 70E CI-
 BpM=0 I=9.1v Mn=455 TIC=105223000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK17JAN

CI⁺ Data



Mass	% Base
43.96	0.90
44.96	0.39
58.03	0.90
132.14	1.19
138.17	0.58
204.22	3.05
254.20	0.51
270.28	3.05
292.25	0.78
334.29	0.43
335.29	0.79
356.28	0.59
358.30	0.71
363.36	3.07
376.32	4.34
377.33	1.54
378.34	0.60
396.37	0.55
415.38	0.68
416.38	0.58
418.36	10.00
435.31	2.48 F
436.42	28.55 F
437.50	5.48 F
456.46	1.40

No. 21


M.Wt. 182

EI⁺/CI⁺/CI⁻


GSE1905+ x1 Bgd=1 12-MAR-91 12:09:0:00:45 78E EI+
 BpM=0 I=10v Hw=103 TIC=300000000 Acnt: Sys:ACE
 G.SANDFORD Cal: PFK11MAR



GSE19010+ x1 Bgd=7 12-MAR-91 12:09:0:01:09 78E CI-
 BpM=0 I=10v Hw=315 TIC=261903000 Acnt: Sys:ACE
 G.SANDFORD Cal: PFK11MAR






GSE19014+ x1 Bgd=13 12-MAR-91 12:09:0:01:20 78E CI-
 BpM=0 I=7.5v Hw=197 TIC=59056000 Acnt: Sys:ACE
 G.SANDFORD Cal: PFK11MAR

EI⁺ Data

Mass	% Base
40.01	6.66
41.01	19.60
43.04	2.15 F
44.98	11.51 F
45.02	2.00 F
45.99	36.95
50.99	3.73
52.00	8.63
52.99	6.63
55.03	2.06 F
56.98	2.49 F
61.98	2.01
62.98	6.81
63.99	18.09
65.00	18.39
66.00	5.53
70.97	13.05
89.95	5.34
90.97	5.85
91.99	46.90
92.99	3.38
108.97	2.77
109.97	3.60
135.95	3.38
141.94	12.81
142.94	2.98
153.93	2.18
154.94	30.42
155.94	5.60
180.93	6.68
181.94	100.00 O
182.94	28.61
183.94	1.81

No. 22




M.Wt. 332

EI⁺/CI⁺/CI⁻EI⁺ Data

Mass	% Base
45.02	0.59
57.06	0.52
60.02	0.46
69.05	0.37
81.91	0.39
83.06	0.39
92.01	6.99
93.02	0.89
96.06	0.38
97.07	0.39
118.01	2.02
119.01	0.32
141.00	0.39
149.00	0.92
163.01	1.01
185.97	0.62
192.99	4.18
195.96	1.64
235.98	0.54
236.19	0.55
243.00	11.60
244.00	2.12
263.01	6.90
264.01	1.77
*304.98	0.45
312.03	0.62
313.02	7.04
314.04	0.68
331.01	0.42
332.02	100.00
333.03	15.01
334.03	1.23
379.08	4.06
380.09	0.75
424.09	1.58
439.14	0.52
482.06	1.61

No.23

M.Wt. 482

EI⁺/CI⁺/CI⁻EI⁺ Data


Mass	% Base	Mass	% Base
40.03	1.65	413.11	1.62
41.03	5.75	414.11	1.16
46.01	2.49	415.13	0.25
52.02	2.18	425.09	0.09
53.01	1.94	433.05	0.16
63.01	1.66	434.06	0.13
64.02	7.11	444.11	1.17
65.02	6.75	445.11	0.39
66.03	3.45	446.12	0.14
68.99	62.68	454.09	0.47
70.00	1.62	455.09	0.09
76.00	4.35	463.11	18.03 F
91.03	4.21	464.15	3.46 F
92.04	73.22	465.15	1.37
93.03	5.09	466.16	0.27
99.99	3.23	478.72	0.06
107.00	2.19	482.08	100.00 F
108.01	1.62	483.11	29.01 F
118.05	21.36	484.24	3.33 F
119.03	1.87	486.04	0.06
126.00	10.71		
127.01	1.00		
133.01	1.06		
176.01	3.78		
196.02	2.71		
218.06	5.29		
221.02	3.78		
268.07	1.52		
286.07	1.60		
321.05	7.26		
371.04	1.49		
394.10	3.07		
412.09	0.08		

No. 24


M.Wt. 196

EI⁺/CI⁺/CI⁻


GSD13014* x1 Bgd=11 13-DEC-90 10:04:0:01:16 70E C1-
 BpM=0 I=3.5v Hn=444 TIC=34791000 Acnt: Sys:ACE
 G.SANDFORD PT= 0⁰ Cat: PFK12DEC HMR:
 MASS:



GSD13017* x1 Bgd=4 13-DEC-90 10:04:0:01:20 70E EI-
 BpM=0 I=1.5v Hn=446 TIC=33672000 Acnt: Sys:ACE
 G.SANDFORD PT= 0⁰ Cat: PFK12DEC HMR:
 MASS:



GSD13023* x1 Bgd=6 13-DEC-90 10:04:0:01:52 70E CI-
 BpM=0 I=10v Hn=464 TIC=150179000 Acnt: Sys:ACE
 G.SANDFORD PT= 0⁰ Cat: PFK12DEC HMR:
 MASS:

EI⁺ Data


Mass	% Base		
25.73	1.02	80.97	0.50
26.74	2.77	89.92	0.44
27.74	3.02	90.92	0.81
30.73	1.34	91.93	0.51
36.77	1.22	95.94	0.32
37.78	2.56	102.92	0.80
38.79	5.38	103.92	0.76
39.79	0.98	104.94	25.58
40.81	2.63	105.94	3.88
41.82	3.55	108.91	0.59
44.80	0.51	109.92	0.54
45.81	10.35	116.86	0.58
46.82	0.37	116.90	0.63
49.84	1.60	121.89	0.40
50.85	5.12	122.93	0.47
51.86	5.25	142.89	0.49
52.87	4.19	153.85	0.86
53.88	1.43	154.84	0.34
54.89	0.89	154.88	0.66
56.85	0.46	167.87	7.68
58.88	1.09	168.88	0.68
59.89	0.55	176.85	0.39
62.89	0.69	176.91	0.48
63.90	4.16	193.87	1.07
64.92	1.21	194.20	0.39
65.92	0.71	194.62	0.36
66.92	0.41	194.84	100.00
66.93	0.38	195.85	75.37
70.91	2.56	196.86	10.43
71.93	0.65	197.85	0.61
74.92	0.75	445.73	1.18
75.93	1.83	446.72	0.41
76.94	1.00		
77.95	12.18		
78.95	3.24		
79.97	10.24		

No.25


M.Wt. 346

EI⁺/CI⁻

GSD14#14* xl Bgd=11 13-DEC-98 10:33:01:17 78E C1-
 BpM=0 I=1.6v Hw=497 TIC=38412088 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK120EC



GSD14#18* xl Bgd=5 13-DEC-98 10:33:01:32 78E EI+
 BpM=0 I=5.6v Hw=682 TIC=327884888 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK120EC

CI⁻ Data



Mass	% Base		
25.71	2.95		
45.81	0.70	361.93	1.62
58.89	0.36	362.88	0.76
63.86	0.50	364.69	0.90
73.95	0.41	370.93	0.75
88.00	0.50	371.92	0.32
88.96	4.89	386.87	4.89
106.94	0.80		
108.96	0.38		
126.82	3.70		
126.92	0.52		
132.89	0.41		
141.94	0.72		
143.89	0.58		
151.86	0.54		
155.89	1.35		
156.88	0.97		
156.92	1.29		
157.90	7.34		
158.90	1.32		
323.84	0.86		
323.98	1.24		
324.78	0.78		
325.81	15.10 F		
325.96	11.19 F		
326.12	0.50		
326.87	6.06		
327.85	0.71		
328.00	0.70		
328.92	0.63		
329.01	0.41		
329.88	0.71		
342.83	0.50		
343.27	0.44		
343.78	0.73		
344.82	2.45		
345.85	100.00		
346.36	0.46		
346.77	17.13 F		
346.95	14.91 F		
347.71	0.97		
347.87	1.01		
348.01	1.02		

No. 26

M.Wt. 496


EI⁺/CI⁺/CI⁻

GSD1204+ x1 Bgd=1 12-DEC-98 16:36:00:36 70E EI⁺
 BpM=0 I=2.6v Hw=600 TIC=70182000 Acnt




GSD1208+ x1 Bgd=6 12-DEC-98 16:36:00:52 70E CI⁺
 BpM=0 I=18v Hw=498 TIC=74301000 Acnt:
 G.SANDFORD

Sys: ACE
 Cal: PFK12DEC
 PT= 0°



GSD12014+ x1 Bgd=12 12-DEC-98 16:36:00:16 70E CI⁻
 BpM=0 I=5.2v Hw=521 TIC=68667000 Acnt:
 G.SANDFORD

Sys: ACE
 Cal: PFK12DEC
 PT= 0°






EI⁺ Data

Mass	% Base		
26.75	1.32	132.93	0.33
27.75	1.62	139.90	0.60
37.79	0.66	175.85	2.98
38.80	3.80	177.89	0.63
40.82	1.14	195.84	2.04
41.83	2.98	202.86	0.44
45.82	1.49	203.88	0.55
49.85	0.49	204.87	0.48
50.86	2.38	210.89	0.32
51.87	3.02	211.87	0.91
52.89	1.77	231.86	1.57
53.89	0.71	241.84	0.31
54.90	1.49	256.85	0.47
63.91	2.26	281.80	1.07
64.92	0.83	298.80	0.31
65.94	0.86	299.79	0.50
66.94	0.56	320.81	0.79
68.91	14.92	325.82	0.75
75.94	1.85	326.81	0.32
76.96	0.82	337.77	0.75
77.96	8.61	338.79	0.40
78.97	3.94	357.76	0.51
79.98	7.92	370.67	1.52
80.99	0.45	375.73	1.90
90.95	0.52	406.71	12.52
91.96	0.66	407.72	2.25
99.91	0.94	425.68	5.96
103.95	1.49	426.69	56.89
104.95	32.11	427.70	10.01
105.96	14.98	428.69	0.62
106.95	0.91	453.72	0.49
107.92	1.02	456.72	0.32
125.89	3.25	476.66	21.93
126.91	0.51	477.68	4.13
129.93	0.65	478.68	0.78
130.94	0.70	494.60	47.88 F
131.94	3.30	495.64	100.00 F
		496.64	29.81
		497.64	3.25

No.27

M.Wt. 246

EI⁺/CI⁺CI⁺ Data



Mass	% Base
45.86	0.39
117.75	0.78
129.73	0.78
130.72	0.39
154.71	1.17
158.75	0.39
169.75	0.79
183.75	0.78
201.77	0.77
203.71	0.40
204.70	0.45
217.64	0.78
224.69	1.56
226.74	0.41
228.68	0.79
239.70	2.54
242.06	2.55 F
243.74	0.78 F
244.69	2.58 F
245.72	25.24 F
246.74	100.00 FO
247.76	17.23 F
248.94	1.56 F
261.78	0.39
288.78	0.39

No. 28

M.Wt. 396


EI⁺/CI⁺/CI⁻

GSD16#8* x1 Bgd=6 13-DEC-98 17:09:00:55 78E C1-
 BpM=0 I=18v Hn=548 TIC=389799888 Acnt: PT= 0⁰ Sys: ACE
 Cal: PFK12DEC




HMR:
 MASS:

GSD16#14* x1 Bgd=11 13-DEC-98 17:09:00:01:10 78E C1-
 BpM=0 I=1.9v Hn=546 TIC=516568888 Acnt: PT= 0⁰ Sys: ACE
 Cal: PFK12DEC



HMR:
 MASS:

GSD16#16* x1 Bgd=1 13-DEC-98 17:09:00:01:26 78E EI-
 BpM=0 I=5.7v Hn=547 TIC=388159888 Acnt: PT= 0⁰ Sys: ACE
 Cal: PFK12DEC



HMR:
 MASS:




EI⁺ Data

Mass	% Base							
68.89	8.41	F	99.91	1.25	171.92	2.06		
68.97	5.84	F	100.42	1.29	172.93	26.19		
69.97	3.77		100.92	4.59	173.93	3.66		
70.98	15.43		101.93	6.00	174.92	0.50		
71.44	0.80		102.94	11.31	226.83	0.49		
71.96	0.98		103.94	2.75	229.83	0.75		
72.45	1.07		112.97	2.19	230.88	0.30		
72.94	1.38		113.92	5.60	238.99	0.31		
73.92	3.64		114.93	10.60	244.82	1.07		
74.93	5.03		115.93	9.98	245.82	3.00		
75.93	4.84		116.93	5.29	326.78	6.72		
76.95	17.50		117.93	2.08	327.76	1.13		
77.44	0.90		125.90	2.08	344.70	0.42		
77.95	8.08		126.91	2.45	F	345.71	0.55	
78.45	1.78		127.01	1.61	F	370.67	1.38	
78.96	3.65		127.92	9.44		376.69	3.59	
79.97	0.69		128.93	6.32	F	377.70	0.95	
80.98	2.06		129.93	68.03	F	381.66	0.60	
81.99	1.67		130.94	100.00		392.70	0.39	
83.00	3.70		131.93	10.57		394.48	3.27	F
84.00	2.14		132.92	1.18		395.67	57.42	F
85.01	10.22		136.98	0.33		398.69	10.27	F
85.46	0.46		138.93	0.54		397.69	0.87	
85.95	1.45		139.91	1.67		445.77	1.14	
86.46	0.90		140.92	4.61				
86.93	2.82		141.91	2.87				
87.93	3.63		142.92	12.08				
88.94	15.95		143.92	6.75				
89.94	9.10		144.92	3.24				
90.95	4.58		153.90	2.46				
91.95	0.73		154.89	12.88				
92.96	0.73		155.90	23.03				
93.96	0.39		156.91	7.79				
94.97	1.15		157.92	75.31				
95.97	1.10		158.92	9.35				
96.99	3.07							
97.97	1.24							
98.99	2.95							

No. 29

M.Wt. 546

EI⁺/CI⁺/CI⁻



EI⁺ Data

Mass	% Base				
68.97	64.44	F	103.01	2.73	156.01
69.06	5.74	F	103.99	5.78	157.02
69.98	1.06	F	105.01	1.61	158.01
70.06	2.34	F	106.95	2.35	F
70.98	1.56	F	107.98	1.58	166.99
71.07	4.10	F	109.07	1.43	168.02
73.02	1.56		111.08	1.48	168.92
74.01	7.70		112.98	1.97	F
75.00	4.19		113.99	16.03	175.48
75.97	8.90	F	115.00	12.78	175.96
76.01	7.44	F	116.00	1.87	180.00
77.01	10.43		117.02	1.15	181.00
77.51	1.17		118.95	1.88	F
78.01	17.21		119.05	1.92	F
78.51	1.99		120.98	1.17	F
79.03	1.56		125.96	27.39	476.95
81.04	2.39	F	126.98	4.69	F
82.03	1.56		128.01	11.51	507.94
83.06	3.02		129.01	6.78	526.94
84.06	1.23		130.03	5.24	527.94
85.06	2.78	F	130.94	1.97	F
87.00	5.30		131.03	1.19	F
87.99	5.26		132.00	1.17	543.88
89.00	2.12		139.01	1.03	544.80
91.01	3.93		139.99	2.97	545.90
93.00	1.56		141.00	12.03	546.93
95.04	2.29		142.02	7.42	547.94
96.06	1.23		143.03	1.22	548.95
97.06	2.73		148.98	42.34	
98.98	1.05	F	149.98	3.93	
99.97	3.39		152.99	2.73	
101.00	9.72		153.96	4.27	F
102.00	4.81		154.18	1.17	F
			154.99	49.01	F



No.30

M.Wt. 263

EI⁺/CI⁺/CI⁻


GSD20034* x1 Bgd=31 25-JAN-91 11:40:04:30 70E
 BpM=0 I=3.0v Hn=313 TIC=50308000
 G.SANDFORD

EI⁺
 Sys:ACE
 Cal:PFK25JAN




GSD20038* x1 Bgd=21 25-JAN-91 11:40:04:46 70E
 BpM=0 I=10v Hn=440 TIC=176713000
 G.SANDFORD

CI⁺
 Sys:ACE
 Cal:PFK25JAN



GSD20041* x1 Bgd=27 25-JAN-91 11:40:04:50 70E
 BpM=0 I=7.8v Hn=445 TIC=200300000
 G.SANDFORD

CI⁻
 Sys:ACE
 Cal:PFK25JAN

EI⁺ Data



Mass	% Base	Mass	% Base
151.28	0.44	217.32	0.64
152.29	0.52	220.29	0.32
155.31	0.38	221.30	6.49
156.30	0.52	222.30	5.73
157.28	1.30	223.31	0.85
158.29	1.30	228.31	1.30
159.31	1.03	229.32	0.61
160.32	7.79	234.32	0.40
161.33	0.67	235.32	3.90
163.28	0.33	236.33	0.49
168.28	0.38	242.33	2.93
169.27	0.38	243.33	1.30
170.30	2.09	244.34	1.30
171.29	1.30	246.33	0.44
172.30	0.49	247.31	3.90
173.33	9.00	248.32	2.72
174.33	1.30	249.32	1.30
175.28	0.60	260.32	1.30
176.28	3.91	261.37	1.78
177.29	1.30	262.33	27.62
182.28	0.47	263.35	100.00
183.29	1.30	264.35	16.63
188.28	0.64	265.35	1.30
189.29	0.51		
190.30	0.30		
194.27	0.65		
195.28	1.66		
196.29	0.49		
197.30	0.49		
198.30	0.51		
201.28	1.30		
202.29	1.30		
203.30	1.70		
215.31	0.71		
216.31	0.34		

No. 31


M.Wt. 236

EI⁺/CI⁺

GSC52115 x1 Bgd=1 20-SEP-98 12:00:01:28 78E EI+
 BpM=0 I=4.0v Mw=236 TIC=110509000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK14SEP



GSC52120 x1 Bgd=6 20-SEP-98 12:00:01:48 78E CI+
 BpM=0 I=10v Mw=447 TIC=169529000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cal: PFK14SEP

EI⁺ Data


Mass	% Base				
40.04	1.18	105.13	0.38	235.27	97.22
41.06	2.24	110.62	0.36	236.28	100.00
42.06	7.72	115.12	0.36	237.28	13.70
43.07	1.29	116.13	1.10	238.29	0.89
44.08	2.48	117.14	0.60		
45.03	1.34	118.15	3.28		
46.04	8.33	119.16	1.00		
50.05	2.57	120.17	0.71		
51.06	2.81	127.13	0.37		
52.06	0.99	128.14	0.77		
55.09	0.59	129.14	3.46		
56.10	0.35	130.15	2.30		
57.11	1.12	131.16	1.63		
60.06	0.31	143.17	0.75		
62.06	1.13	144.18	0.40		
63.06	1.93	145.19	7.30		
64.06	1.95	146.19	1.25		
65.07	0.38	147.15	8.58		
66.08	0.73	148.16	0.67		
69.11	1.13	149.14	1.45		
71.05	1.71	170.21	0.70		
71.13	0.47	179.17	0.39		
72.58	0.56	192.19	9.07		
74.07	0.90	193.19	1.98		
75.07	2.87	194.21	1.58		
76.08	2.93	195.22	0.49		
77.09	1.81	205.20	0.35		
78.09	0.49	206.21	2.05		
81.13	0.34	207.22	0.84		
88.08	0.55	208.23	0.38		
89.10	0.84	219.23	9.05		
90.08	2.82	220.24	9.83		
91.09	0.79	221.24	3.37		
95.10	0.38	222.25	0.92		
101.10	0.47	233.25	0.91		
102.11	12.05	234.28	1.02		
103.12	2.37				
104.12	3.92				

No. 32


M.Wt. 386

EI⁺/CI⁺/CI⁻


GSC15#18* x1 Bgd=1 24-OCT-98 14:26:0:01:33 78E EI⁺
 BpM=0 I=2.8v Hn=446 TIC=3782888
 G.(ACE GOAL PORCHER).SANDFORD



GSC15#22* x1 Bgd=7 24-OCT-98 14:26:0:01:49 78E CI⁺
 BpM=0 I=7.5v Hn=447 TIC=7217888 Acnt: Sys:ACE
 G.(ACE GOAL PORCHER).SANDFORD PT= 0° Cal: PFK220CT



GSC15#28* x1 Bgd=11 24-OCT-98 14:26:0:02:13 78E CI⁻
 BpM=0 I=1.2v Hn=388 TIC=18631888 Acnt: Sys:ACE
 G.(ACE GOAL PORCHER).SANDFORD PT= 0° Cal: PFK220CT

EI⁺ Data



Mass	% Base	Mass	% Base
42.03	3.10	342.09	1.82
43.04	0.31	367.19	6.87
44.05	1.20	368.19	1.25
46.01	1.09	369.15	0.55
50.02	0.41	370.15	0.65
51.03	0.86	371.17	0.74
63.03	0.49	372.17	0.41
69.00	9.87	384.19	1.16
71.00	1.27	385.19	56.36
75.02	0.84	386.19	100.00
76.01	0.77	387.20	29.80
77.04	0.83	388.21	3.35
90.04	0.31	446.33	0.57
102.04	3.60		
103.05	0.94		
104.05	0.96		
116.05	0.64		
117.06	0.34		
118.07	0.60		
120.09	1.20		
121.09	2.55		
129.06	2.31		
130.06	1.84		
131.07	1.28		
143.07	0.42		
145.09	12.59		
146.09	3.32		
147.05	1.22		
172.10	0.41		
266.13	6.25		
267.13	0.91		
297.15	2.10		
298.17	0.96		
301.12	0.45		
316.15	9.12		
317.15	4.05		
318.16	0.58		

No. 33


M.Wt. 536

EI⁺/CI⁺/CI⁻


GSC16834 x1 Bgd=18 23-OCT-98 17:02:08:02:36 78E EI⁺
 BpM=0 I=10v Hn=613 TIC=1862828932
 G.SANDFORD



GSC16837 x1 Bgd=23 23-OCT-98 17:02:08:02:48 78E CI⁺
 BpM=0 I=10v Hn=553 TIC=711174976 Rcnt: Sys:RCE
 G.SANDFORD PT= 0° Cal: PFK220CT



GSC16845 x1 Bgd=41 23-OCT-98 17:02:08:03:19 78E CI⁻
 BpM=0 I=7.5v Hn=829 TIC=141258880 Rcnt: Sys:RCE
 G.SANDFORD PT= 0° Cal: PFK220CT

CI⁻ Data



Mass	% Base	Mass	% Base
126.83	0.97	497.98	1.08
156.91	1.26	501.95	71.22 F
168.90	0.40	502.97	14.00 F
175.90	1.85	503.98	1.38
194.89	0.99	518.00	1.37
211.86	0.56	517.00	1.69
261.94	0.95	518.02	1.73
344.90	0.95	519.01	0.32
351.95	0.39	520.96	0.30
352.96	2.01	522.00	4.11
353.97	0.30	523.02	1.02
358.96	0.91	534.85	1.29 F
366.95	1.80	536.00	100.00 F
367.97	1.17	537.01	19.45 F
385.97	1.50	538.02	2.15
387.89	1.63	532.03	0.62
393.89	0.34		
394.90	0.46		
396.89	0.40		
406.90	0.58		
407.23	0.95		
411.89	1.97		
412.88	3.53		
413.88	0.48		
415.88	11.93		
416.89	1.45		
429.90	0.63		
430.90	1.60		
431.89	3.12		
432.90	2.01		
459.96	0.36		
462.02	0.96		
463.95	0.68		
466.97	0.71		
469.94	0.56		
481.95	2.02		
482.96	0.50		
483.97	1.87		
484.97	0.41		

No. 34


M.Wt. 414

EI⁺/CI⁺

GSC19010• x1 Bgd=1 1-NOV-98 15:52:00:42 78E EI•
 BpM=0 I=2.9v Hn=565 TIC=224125000 Acnt: Sys:ACE
 PT= 0° Cal: PFK380CT



GSC19010• x1 Bgd=10 1-NOV-98 15:52:00:01:34 78E CI•
 BpM=0 I=2.8v Hn=416 TIC=38893000 Acnt: Sys:ACE
 PT= 0° Cal: PFK380CT

CI⁺ Data

Mass	% Base
44.04	3.92
44.08	1.96
45.02	1.96
45.04	1.96
46.04	1.96
46.06	1.96
46.07	1.96
58.05	1.96
58.08	1.96
59.05	1.49
60.04	1.96
61.03	1.96
72.07	1.96
74.05	1.96
88.06	1.96
90.02	1.96
102.02	1.96
102.08	1.96
106.03	1.96
106.08	1.96
116.06	1.96
121.03	1.96
122.06	3.92
123.03	1.96
134.06	2.02
149.08	1.96
150.08	91.74
150.34	1.96
151.09	10.11
152.04	1.96
397.06	1.96
415.12	12.89
416.15	2.18


No. 35

M.Wt. 564

Cl⁻

GSC17X038 x1 0gd=11 30-OCT-90 12:29:08:02:22 78E Cl⁻
 BpM=8 l=1.7v Mw=566 TIC=19514088 Acnt: Sys:ACE
 G.SANDFORD *X1=0 PT= 0° Cat: PFK380CT

HMR:
 MASS:

Cl⁻ Data

Mass	% Mass
393.93	1.22
395.02	1.47
411.92	4.95
412.92	6.07
413.99	1.81
415.89	1.33
431.90	3.34
432.90	3.05
515.98	4.60
516.98	0.32
526.02	2.13
536.02	0.46
544.01	7.56
545.01	3.47
546.03	3.81
547.05	0.49
562.01	2.27
563.04	1.86
564.02	100.00
565.02	22.30
566.01	2.07

No. 36


M.Wt. 444

EI⁺/CI⁺

GSC4050 x1 8gd=3 19-SEP-98 18:41:00:42 78E EI+
 BpM=0 I=7.5v Hm=577 TIC=298626016 Acnt: PT= 0°

Sys: ACE
 Cal: PFK14SEP


HMR:
 MASS:



GSC4020 x1 8gd=0 19-SEP-98 18:41:01:41 78E CI+
 BpM=0 I=10v Hm=578 TIC=174103000 Acnt: PT= 0°

Sys: ACE
 Cal: PFK14SEP

HMR:
 MASS:

CI⁺ Data


Mass	% Base
44.06	1.67
46.08	2.74
58.08	0.76
88.10	0.38
330.30	3.45
331.30	0.71
400.27	0.58
402.28	0.32
413.27	0.33
429.30	0.60
431.32	0.86
442.36	0.96
443.35	0.34
444.29	1.52 F
445.31	100.00 FO
446.33	27.20 F
447.34	10.34
448.34	2.51
449.35	0.41

No. 37


M.Wt. 236

EI⁺/CI⁺

GSC2200+ x1 Bgd=1 31-OCT-90 14:52:00:30 70E EI⁺
 BpM=0 I=6.0v Hm=354 TIC=240250000 Acnt: Sys:ACE
 PT= 0° Cal: PFK300CT



GSC2200+ x1 Bgd=6 31-OCT-90 14:52:00:54 70E CI⁺
 BpM=0 I=10v Hm=447 TIC=990010000 Acnt: Sys:ACE
 PT= 0° Cal: PFK300CT

EI⁺ Data



mass	base	mass	base	mass	base
69.00	0.58	104.06	0.58	161.07	0.58
69.07	1.73	105.08	2.91	162.07	1.77
70.09	0.58	106.08	0.58	167.07	0.58
71.01	8.08 F	110.05	0.58	168.07	0.58
71.10	1.36 F	110.55	0.58	169.08	0.58
72.03	2.90	111.14	0.49	171.10	0.58
73.05	0.58	113.08	0.58	176.09	0.58
74.03	1.73	114.10	0.58	181.09	1.15
75.03	1.73	115.05	0.58	189.09	0.58
76.04	3.04	115.08	0.58	196.12	0.58
77.05	39.99	116.04	2.30	206.08	1.36
78.05	5.39	117.05	2.50	207.08	24.10
79.07	1.15	118.07	4.26	208.09	5.80
80.07	0.58	119.09	2.03	209.10	0.60
81.09	0.82	120.11	4.81	217.12	0.58
82.08	0.58	121.09	1.15	219.09	0.58
83.10	1.15	122.07	2.06	220.09	0.58
85.12	0.70	128.08	0.82	221.10	100.00
87.06	0.58	129.08	0.81	222.11	12.80
88.04	0.58	131.09	0.58	223.13	1.15
89.05	0.65	132.05	0.58	233.13	0.58
90.05	4.04	134.06	0.58	234.16	0.58
91.06	4.61	135.06	1.73	235.13	14.98
92.06	1.17	136.07	0.39	236.14	46.86
93.06	0.64	139.07	0.64	237.15	6.33
94.06	1.15	143.06	1.23	238.16	0.58
95.08	0.58	144.08	1.33		
96.05	0.58	145.07	0.58		
97.12	0.58	147.06	0.72		
102.05	0.58	149.05	6.68		
103.07	1.25	150.06	0.58		
		156.09	0.90		
		157.07	0.58		
		158.08	0.58		
		159.07	1.73		
		160.06	0.58		

No. 38


M.Wt. 386

EI⁺/CI⁺/CI⁻


GSE15010* x1 Bgd=0 12-MAR-91 09:58:0:01:00 70E C1+
 BpM=0 I=0.2v Hn=536 TIC=250004000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cat: PFK11MAR



GSE15015* x1 Bgd=14 12-MAR-91 09:58:0:01:32 70E C1-
 BpM=0 I=2.0v Hn=386 TIC=25016000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cat: PFK11MAR



GSE15016* x1 Bgd=1 12-MAR-91 09:58:0:01:37 70E EI+
 BpM=0 I=0.1v Hn=462 TIC=367995000 Acnt: Sys:ACE
 G.SANDFORD PT= 0° Cat: PFK11MAR



EI⁺ Data



Mass	% Base				
41.03	1.33	120.02	20.65	385.91	76.16 F
42.03	6.38	121.02	3.40	386.93	21.57 F
46.00	9.77	121.98	5.53	387.93	2.45
50.00	6.32	127.98	1.20	462.95	1.47
51.01	27.25	128.98	1.59		
52.01	2.93	129.98	3.87		
53.01	2.07	130.99	4.52		
62.00	1.44	134.97	1.49		
63.00	5.31	142.98	2.77		
64.01	3.50	143.98	4.04		
65.02	9.99	144.99	1.09		
66.02	1.44	155.97	1.25		
68.97	20.25 F	161.96	2.41		
69.02	2.17 F	174.96	9.66		
70.98	13.87	175.96	4.96		
71.99	6.44	187.95	1.37		
73.99	1.52	188.96	4.26		
74.99	1.94	195.90	1.34		
75.99	3.67	227.89	2.16		
77.01	72.27	237.92	3.40		
78.01	8.73	266.88	1.44		
79.01	1.88	267.90	1.04		
88.99	1.01	268.91	2.14		
89.99	5.20	287.90	3.97		
91.00	7.36	288.90	1.97		
92.00	1.54	296.93	1.44		
94.00	1.19	301.92	2.88		
99.95	1.61	308.90	2.04		
101.98	1.03	316.94	3.39		
102.99	4.87	356.73	9.38 F		
104.00	20.93	357.89	9.24 F		
105.01	6.75	358.90	1.44		
106.01	2.93	366.93	10.78		
107.96	1.25	367.93	1.92		
115.98	1.54	370.88	100.00 F		
116.98	4.31	371.91	15.93 F		
118.00	12.42	372.91	1.27		
119.00	4.77	384.84	10.38 F		

No. 39


M.Wt. 252

EI⁺/CI⁺

GSD6020 x1 Bgd=16 26-NOV-98 12:19:01:41 78E EI+
 BpM=8 I=4.7v Hn=447 TIC=218445888 Acnt: Sys: ACE
 G.SANDFORD PT= 0° Cal: PFK26NOV





GSD6024 x1 Bgd=6 26-NOV-98 12:19:01:56 78E CI+
 BpM=8 I=18v Hn=279 TIC=232528888 Acnt: Sys: ACE
 G.SANDFORD PT= 0° Cal: PFK26NOV

EI⁺ Data

Mass	% Base				
69.06	2.76	115.04	0.33	235.02	0.66
70.07	1.14	116.00	1.21	236.03	0.84
70.99	3.64 F	117.01	1.33	237.04	1.07
71.07	2.07 F	118.01	0.44	251.05	0.48
72.00	1.29	119.05	0.43	252.06	9.30
73.02	1.11	120.03	2.95	253.07	1.34
74.01	0.91	121.01	5.74	293.96	1.30
75.01	1.05	122.02	1.47	325.94	0.36
76.02	3.35	123.02	0.64	446.17	1.49
77.03	10.28	128.02	0.51	447.18	0.49
78.03	4.28	129.04	0.57		
79.03	1.27	132.01	0.70		
80.05	0.36	134.03	0.53		
81.06	1.30	135.04	1.16		
82.07	0.95	137.98	0.59		
83.02	1.97 F	143.00	0.59		
83.07	1.70 F	144.04	0.70		
84.08	0.49	145.02	1.02		
85.09	1.26	149.00	30.23		
86.02	0.31	150.01	2.64		
86.96	0.94	151.01	0.88		
87.03	0.33	154.02	0.42		
89.03	0.39	161.02	0.30		
90.02	1.45	162.96	0.57		
91.04	2.51	167.01	1.32		
92.03	1.71	168.02	0.85		
93.02	2.01	172.03	0.84		
94.05	0.47	178.06	1.36		
95.06	1.52	182.96	2.12		
96.08	0.59	190.00	0.52		
97.08	1.68	193.02	0.90		
98.99	0.44	194.02	1.19		
102.02	0.56	205.06	0.53		
103.02	0.90	206.02	5.98		
104.02	2.63	207.03	1.56		
105.02	6.36	208.03	0.49		
106.03	1.15	209.04	1.92		
107.04	0.65	210.04	0.40		
108.04	0.49	221.04	100.00		
109.08	0.68	222.05	12.97		
111.10	0.89	223.07	2.59		
		233.06	0.33		

No. 40



M.Wt. 265

EI⁺/CI⁺EI⁺ Data

Mass	% Base				
69.04	0.99	123.47	0.43	222.97	0.34
70.97	3.42	124.48	0.73	229.97	0.56
71.05	0.45	128.98	0.50	232.93	0.59
71.97	3.75	129.98	0.69	233.94	6.87
73.01	1.21	130.99	2.02	234.95	3.64
74.00	0.73	131.48	2.31	235.96	0.59
74.99	1.15	131.99	2.64	245.99	0.35
75.99	1.97	132.49	3.44	247.95	6.13
77.00	6.87	133.00	3.06	248.96	4.58
78.00	2.95	134.02	1.86	249.96	22.67
79.01	1.24	135.01	2.03	250.97	3.53
79.99	0.36	143.98	1.89	261.96	0.32
81.03	0.40	144.98	0.98	262.98	0.77
81.88	0.39	146.00	0.43	263.95	33.96
82.01	0.38	147.01	2.68	264.96	100.00
87.98	0.32	148.02	1.75	265.96	14.67
88.99	0.55	148.95	3.80 F	266.99	0.90
89.99	2.66	149.03	2.00 F		
91.00	2.13	157.99	0.51		
92.00	1.61	159.00	0.45		
93.00	1.52	160.00	2.28		
100.00	0.55	160.99	0.41		
101.98	1.00	162.04	1.75		
102.98	0.65	164.05	0.58		
103.99	4.17	177.06	0.51		
104.98	12.87	184.99	0.40		
106.00	1.56	192.94	0.56		
107.02	0.87	193.94	0.64		
108.02	0.98	204.93	0.39		
115.96	1.31	205.94	6.61		
116.96	2.95	206.94	2.48		
118.00	3.52	207.95	0.56		
119.00	2.17	218.94	0.40		
120.02	4.21	219.94	1.31		
121.02	3.80	220.95	4.04		
122.03	0.83	221.96	0.59		

No. 41

M.Wt. 538

EI⁺/CI⁺EI⁺ Data

68.91	68.33	157.90	1.22
69.92	1.21	158.92	19.15
70.92	1.32	159.92	2.08
75.93	4.24	432.63	5.22
76.95	1.57	433.65	13.70
77.96	33.24	434.66	1.59
78.97	72.30	438.69	0.33
79.97	14.15	444.66	0.31
80.96	1.20	452.73	0.87
87.92	0.30	453.74	2.18
89.94	0.50	454.75	0.46
90.94	1.17	457.68	0.54
91.94	4.00	483.63	0.39
92.97	27.22	501.59	1.17
93.95	2.87	502.59	0.53
94.95	0.64	503.62	18.41
99.90	4.84	504.61	3.66
102.94	1.56	505.63	0.42
103.94	1.68	507.61	5.63
104.95	5.69	508.62	1.42
105.95	1.79	509.62	0.78
106.97	90.01	510.60	0.53
107.95	7.42	517.65	2.73
108.96	0.48	518.64	0.64
111.90	0.58	521.58	5.94 F
113.91	1.39	522.61	56.13 F
116.94	0.55	523.60	77.24 F
117.95	1.61	524.62	13.28 F
118.92	1.77	525.60	1.37
119.95	0.78	535.64	0.86
120.96	4.41	536.64	9.59 F
121.97	51.54	537.66	14.64 F
122.99	100.00	538.66	2.55
123.98	8.63	573.65	0.64
124.98	0.32	574.63	0.34
125.89	1.99		
126.89	2.86		
127.90	0.42		
129.93	0.59		
130.89	1.29		
131.94	7.81		
132.93	6.74		

APPENDIX FOUR

The Board of Studies in Chemistry requires that each postgraduate research thesis contains an appendix listing:-

- (1) all research colloquia, seminars and lectures arranged by the Department of Chemistry during the period of the author's residence as a postgraduate student;
- (2) lectures organised by Durham University Chemical Society;
- (3) all research conferences attended and papers presented by the author during the period when research for the thesis was carried out;
- (4) details of the postgraduate induction course.

**COLLOQUIA, LECTURES AND SEMINARS GIVEN BY INVITED SPEAKERS,
OCTOBER 1988 - SEPTEMBER 1991**

(Those attended are marked *)

- | | |
|-------------|--|
| 18.10.88 | Dr. J. Dingwall (Ciba Geigy)
<i>Phosphorous Containing Amino Acids: Biologically Active Natural and Unnatural Products</i> |
| 24.11.88 | Drs. R.R. Baldwin and R.W. Walker (Hull University)
<i>Combustion: Some Burning Problems</i> |
| 12.88 | Dr. G. Hardgrove (St. Olaf College, USA)
<i>Polymers in the Physical Chemistry Laboratory</i> |
| 25.1.89 | Dr. L. Harwood (Oxford University)
<i>Synthetic Approaches to Phorbols Via Intramolecular Furan Diels-Alder Reactions: Chemistry Under Pressure</i> |
| 2.2.89
* | Prof. L.D. Hall (Addenbrooke's Hospital, Cambridge)
<i>NMR - A Window to the Human Body</i> |
| 9.2.89
* | Prof. J.E. Baldwin (Oxford University)
<i>Recent Advances in the Bioorganic Chemistry of Penicillin Biosynthesis</i> |

- 15.2.89 Dr. A.R. Butler (St. Andrews University)
* *Cancer in Linxiam: The Chemical Dimension*
- 16.2.89 Prof. B.J. Aylett (Queen Mary College, London)
Silicon Based Chips: The Chemist's Contribution
- 1.3.89 Dr. R.J. Errington (Newcastle University)
Polymetalate Assembly in Organic Solvents
- 15.3.89 Dr. R. Aveyard (Hull University)
Surfactants at your Surface
- 20.4.89 Dr. M. Casey (Salford University)
Sulphoxides in Stereoselective Synthesis
- 27.4.89 Dr. D. Crich (University College, London)
Some Novel Uses of Free Radicals in Organic Synthesis
- 11.5.89 Dr. J. Frey (Southampton University)
Spectroscopy of the Reaction Path: Photodissociation Raman Spectra of NOCl
- 10.11.89 Prof. J.I.G. Cadogan (B.P.)
From Pure Science to Profit
- 17.10.89 Dr. F. Palmer (Nottingham University)
* *Thunder and Lightning*
- 25.10.89 Prof. C. Floriani (Lausanne University, Switzerland)
Molecular Aggregates - A Bridge Between Homogeneous and Heterogeneous Systems
- 1.11.89 Dr. J.P.S. Badyal (Durham University)
Breakthroughs in Heterogeneous Catalysis
- 9.11.89 Prof. N.N. Greenwood (Leeds University)
Novel Cluster Geometries in Metalloborane Chemistry
- 10.11.89 Prof. J.E. Bercaw (California Institute of Technology)
* *Synthetic and Mechanistic Approaches to Ziegler-Natta Polymerisation of Olefins.*

- 13.11.89 Dr. J. Becher (Odense University)
* *Synthesis of New Macrocyclic Systems using Heterocyclic Building Blocks*
- 16.11.89 Dr. D. Parker (Durham University)
* *Macrocycles, Drugs and Rock 'n' Roll*
- 29.11.89 Prof. D.J. Cole-Hamilton (St. Andrews University)
New Polymers from Homogeneous Catalysis
- 30.11.89 Dr. M.N. Hughes (King's College, London)
* *A Bug's Eye View of the Periodic Table*
- 4.12.89 Dr. D. Graham (B.P. Research Centre)
How Proteins Absorb on Interfaces
- 6.12.89 Dr. R.L. Powell (ICI)
* *The Development of CFC Replacements*
- 7.12.89 Dr. A. Butler (St. Andrews University)
* *The Discovery of Penicillin: Facts and Fancies*
- 13.12.89 Dr. J. Klinowski (Cambridge University)
Solid State NMR Studies of Zeolite Cages
- 15.12.89 Prof. R. Huisgen (Universitat Munchen)
* *Recent Mechanistic Studies of [2+2] Additions*
- 24.1.90 Dr. R.N. Perutz (York University)
Plotting the Course of C-H Activations with Organometallics
- 31.1.90 Dr. U. Dyer (Glaxo)
* *Synthesis and Conformation of C-Glycosides*
- 1.2.90 Prof. J.H. Holloway (Leicester University)
* *Noble Gas Chemistry*
- 7.2.90 Dr. D.P. Thompson (Newcastle University)
The role of Nitrogen in Extending Silicate Crystal Chemistry

- 8.2.90 Rev. R. Lancaster (Kimbolton Fireworks)
* *Fireworks - Principles and Practice*
- 12.2.90 Prof. L. Lunazzi (University of Bologna)
Application of Dynamic NMR to the Study of Conformational Isomerism
- 14.2.90 Prof. D. Sutton (Simon Fraser University, Vancouver B.C.)
Synthesis and Applications of Dinitrogen and Diazo Compounds of Rhenium and Iridium
- 15.2.90 Prof. L. Crombie (Nottingham University)
The Chemistry of Cannabis and Khat
- 21.2.90 Dr. C. Bleasdale (Newcastle University)
The Mode of Action of some Anti-tumour Agents
- 22.2.90 Prof. D.T. Clark (ICI Wilton)
* *Spatially Resolved Chemistry using Nature's Paradigm in the Advanced Materials Area*
- 28.2.90 Dr. R.K. Thomas (Oxford University)
Neutron Reflectometry from Surfaces
- 1.3.90 Dr. J.F. Stoddart (Sheffield University)
* *Molecular Lego*
- 8.3.90 Dr. A.K. Cheetham (Oxford University)
* *Chemistry of Zeolite Cages*
- 21.3.90 Dr. I. Powis (Nottingham University)
Spinning off in a huff: Photodissociation of Methyl Iodide
- 23.3.90 Prof. J.M. Bowman (Emory University)
Fitting Experiment with Theory in Ar-OH
- 9.7.90 Prof. L.S. German (USSR Academy of Sciences - Moscow)
* *New Syntheses in Fluoroaliphatic Chemistry: Recent Advances in the Chemistry of Fluorinated Oxiranes*

- 9.7.90 Prof. V.E. Platonov (USSR Academy of Sciences - Novosibirsk)
* *Polyfluoroindanes: Synthesis and Transformation*
- 9.7.90 Prof. I.N. Rozhkov (USSR Academy of Sciences - Moscow)
* *Reactivity of Perfluoroalkyl Bromides*
- 11.10.90 Dr. W.A. MacDonald (ICI Wilton)
* *Materials for the Space Age*
- 24.10.90 Dr. M. Bochmann (U.E.A.)
Synthesis, Reactions and Catalytic Activity of Cationic Titanium Alkyls
- 26.10.90 Prof. R. Soulen (South Western University, Texas)
* *Chemistry of some Fluorinated Cyclobutenes*
- 31.10.90 Dr. R. Jackson (Newcastle University)
* *New Synthetic Methods: α -aminoacids and Small Rings*
- 1.11.90 Dr. N. Logan (Nottingham University)
Rocket Propellants
- 6.11.90 Dr. P. Kocovsky (Uppsala)
* *Stereo-controlled Reactions Mediated by Transition and Non-Transition Metals*
- 7.11.90 Dr. D. Gerrard (B.P.)
Raman Spectroscopy for Industrial Analysis
- 7.11.90 Dr. W. Dolbier (Gainesville, Florida)
* *Rearrangements of bis CF₃ Vinyl Aromatics: a Route to 1,3,5-Hexatrienes*
- 8.11.91 Dr. S.K. Scott (Leeds University)
* *Clocks, Oscillations and Chaos*
- 14.11.90 Prof. T. Bell (SUNY, Stony Brook)
* *Functional Molecular Architecture and Molecular Recognition*
- 21.11.90 Prof. J. Pritchard (Queen Mary and Westfield College, London)
Copper Surfaces and Catalysts

- 28.11.90 Dr. B.J. Whitaker (Leeds University)
Two-dimensional Velocity Imaging of State-selected Reaction Products
- 29.11.90 Prof. D. Crout (Warwick University)
Enzymes in Organic Synthesis
- 5.12.90 Dr. P.G. Pringle (Bristol University)
* *Metal Complexes with Functionalised Phosphines*
- 13.12.90 Prof. A.H. Cowley (University of Texas)
New Organometallic Routes to Electronic Materials
- 15.1.91 Dr. B.J. Alder (Lawrence Livermore Labs., California)
Hydrogen in all its Glory
- 17.1.91 Dr. P. Sarre (Nottingham University)
Comet Chemistry
- 23.1.91 Prof. J.S. Higgins (Imperial College, London)
Rheology and Molecular Structure of Ionomer Solutions
- 24.1.91 Dr. P.J. Sadler (Birkbeck College, London)
Design of Inorganic Drugs: Precious Metals, Hypertension and HIV
- 30.1.91 Prof. E. Sinn (Hull University)
New Results in High T_C Superconductivity
- 31.1.91 Dr. D. Lacey (Hull University)
* *Liquid Crystals*
- 6.2.91 Dr. R. Bushby (Leeds University)
* *Biradicals and Organic Magnets*
- 14.2.91 Dr. M.C. Petty (Durham University)
* *Molecular Electronics*
- 20.2.91 Prof. B.L. Shaw (Leeds University)
* *New Chemistry with Transition Metal Multihydrides*
- 28.2.91 Dr. J. Brown (Oxford University)

Can Chemistry Provide Catalysts Superior to Enzymes?

- 6.3.91 Dr. C.M. Dobson (Oxford University)
NMR Studies of Dynamics in Molecular Crystals
- 7.3.91 Dr. J. Markam (ICI Pharmaceuticals)
* *DNA Fingerprinting*
- 24.4.91 Prof. R.R. Schrock (MIT)
* *Metal-ligand Multiple Bonds and Metathesis Initiators*
- 25.4.91 Prof. T. Hudlicky (Virginia Polytechnic Institute)
Biocatalysis and Symmetry Based Approaches to the Efficient Synthesis of Complex Natural Products
- 20.6.91 Prof. M.S. Brookhart (University of North Carolina)
Olefin Polymerisations, Oligomerisations and Dimerisations Using Electrophilic Late Transition Metal Catalysts
- 29.7.91 Dr. M.A. Brimble (Massey University, New Zealand)
* *Synthetic Studies Towards the Antibiotic Griseusin-A*

Research Conferences Attended

- Dec 88 Royal Society of Chemistry Perkin Division, One Day Meeting, York University.
- April 1989 North East Graduate Symposium, Durham University.
- 5.7.89 Royal Society of Chemistry Heterocyclic Group, Postgraduate Heterocyclic Symposium, Sheffield University.
- Aug. 89 European Symposium on Fluorine Chemistry, Leicester University.
- 13.12.89 Modern Aspects of Stereochemistry, One Day Meeting, Sheffield University
- 15.12.89 Royal Society of Chemistry Perkin Division, One Day Meeting, Durham University.

- 7.3.90 SCI Graduate Symposium, York University.
- 2.4.90 North East Graduate Symposium, Newcastle University.
- Sept 91 13th International Symposium on Fluorine Chemistry, Ruhr
Universität, Bochum, Germany.

FIRST YEAR INDUCTION COURSE

This course consists of a series of one hour lectures on the services available in the department.

Departmental Organisation - Dr. E.J.F. Ross

Safety Matters - Dr. M.R. Crampton

Electrical Appliances - Mr. B.T. Barker

Chromatography and Microanalysis - Mr. T.F. Holmes

Atomic Absorptiometry and Inorganic Analysis - Mr. R. Coult

Library Facilities - Mr. R.B. Woodward

Mass Spectroscopy - Dr. M. Jones

Nuclear Magnetic Resonance Spectroscopy - Dr. R.S. Matthews

Glass-blowing Techniques - Mr. R. Hart and Mr. G. Haswell

REFERENCES

1. R.D. Chambers, *"Fluorine in Organic Chemistry"*, John Wiley and Sons, New York, 1973.
2. M.R.C. Gerstenberger and A. Haas, *Angew. Chem., Int. Ed. Engl.*, 1981, **20**, 647.
3. S. Rozen and R. Filler, *Tetrahedron*, 1985, **41**, 1111.
4. J. Mann, *Chem. Soc. Rev.*, 1987, **16**, 381.
5. J. Emsley, *M. Sc. Thesis*, University of Durham, 1988.
6. C.L. Liotta and H.P. Harris, *J. Am. Chem. Soc.*, 1974, **96**, 2250.
7. G.C. Yakobson, *J. Fluorine Chem.*, 1985, **28**, 73.
8. K. Makuo and H. Yoshioka, *J. Fluorine Chem.*, 1987, **35**, 677.
9. Y. Yoshida and Y. Kimura, *Chem. Lett.*, 1988, 1355.
10. K. Makino and H. Yoshioka, *J. Fluorine Chem.*, 1987, **35**, 677.
11. M. Gingras and D.N. Harpp, *Tetrahedron Lett.*, 1988, **37**, 4669.
12. S. Dermiek and Y. Sason, *J. Org. Chem.*, 1985, **50**, 879.
13. G.L. Cantrell, U.S. Patent No. 4 849 552 (1989). *C. Abstr.* **112**: 118436f.
14. E. Kysela and R. Braden, German Patent DE 3 827 436 (1990). *C. Abstr.* **113**: 40127e.
15. J.H. Clark and D.J. MacQuarrie, *Tetrahedron Lett.*, 1987, **28**, 111.
16. N. Yazawa, H. Suzuki, Y. Yoshida, O. Furusawa, and Y. Kimura, *Chem. Lett.*, 1989, 2213.
17. Y. Yoshida, Y. Kimura, and M. Tomoi, *Chem. Lett.*, 1990, 769.
18. N. Ishikawa, T. Kitazume, T. Yamazaki, Y. Mochida, and T. Tatsuio, *Chem. Lett.*, 1981, 761.
19. N. Ishikawa, T. Kitazune, and M. Nakabayashi, *Chem. Lett.*, 1980, 1089.
20. Y. Kimura and H. Suzuki, *Tetrahedron Lett.*, 1989, **30**, 1271.
21. A. McKillop and D.W. Young, *Synthesis*, 1979, **401**, 481.
22. J.H. Clark, A.J. Hyde, and D.K. Smith, *J. Chem. Soc., Chem. Commun.*, 1986, 791.
23. J. Ichihara, T. Matsuo, T. Hanafusa, and T. Ando, *J. Chem. Soc., Chem. Commun.*, 1986, 793.
24. K. Takemoto, J. Ichihara, A. Hanabusa, and T. Ando, Japanese Patent 62 114 939 (1987).
25. H. Liu, H. Lu, P. Wang, P. Sun, and F. Zhang, *J. Chem. Soc., Perkin Trans. I*, 1990, 2624.
26. H. Liu, P. Wang, and P. Sun, *J. Fluorine Chem.*, 1989, **43**, 429.
27. J. Cousseau and P. Albert, *J. Org. Chem.*, 1989, **54**, 5380.
28. D.P. Cox, J. Terpinski, and W. Lawrynowicz, *J. Org. Chem.*, 1984, **49**, 3216.
29. J.H. Clark and D.K. Smith, *Tetrahedron Lett.*, 1985, **26**, 2233.
30. I. Latif, A. Malik, and W. Voelter, *Liebigs Ann. Chem.*, 1987, 617.
31. K.O. Christe, *Inorg Chem*, in press

32. K.O. Christe and W.W. Wilson, *J. Fluorine Chem.*, 1990, **47**, 117.
33. S.J. Brown and J.H. Clark, *J. Chem. Soc., Chem. Commun.*, 1985, 672.
34. A.A. Kolomeitsev, Y.L. Yagupol'skii, G.N. Koidan, A.P. Marchenko, and A.M. Pinchuk, *Zh. Org. Chim.*, 1990, **26**, 1143.
35. J. Leroy, E. Hebert, and C. Wakselman, *J. Org. Chem.*, 1979, **44**, 3406.
36. M. Hudlicky, "Organic Reactions", Vol 35, John Wiley and Sons, New York, 1986.
37. T. Hamatani, S. Matsubara, H. Matsuda, and M. Schlosser, *Tetrahedron*, 1988, **44**, 2875.
38. P.A. Messina, K.C. Mange, and W.J. Middleton, *J. Fluorine Chem.*, 1989, **42**, 137.
39. W.B. Farnham, D.A. Dixon, and J.C. Calabrese, *J. Am. Chem. Soc.*, 1988, **110**, 2607.
40. L.N. Markovskii, V.L. Tovstenko, E.G. Martynyuk, V.E. Pashinnik, Y.V. Shermolovich, and S.V. Iksanova, *Zh. Org. Chim.*, 1990, **26**, 1372.
41. S.S. Hecht and E.S. Rothman, *J. Org. Chem.*, 1973, **38**, 395.
42. French Patent 1 374 591 (1964), Ciba Ltd: *C. Abstr.*, **62**: P9212a.
43. M. van der Akken and F. Jellinek, *Recl. Trav. Chim. Pays-Bas*, 1967, **86**, 275.
44. R.F. Hirschmann, R. Miller, J. Wood, and R.E. Jones, *J. Am. Chem. Soc.*, 1956, **78**, 4956.
45. W.R. Dolbier, T.A. Gray, and K. Ohnishi, *Synthesis*, 1987, 956.
46. W.R. Dolbier, L. Celewicz, and K. Ohnishi, *Tetrahedron Lett.*, 1989, **30**, 4929.
47. G.A. Olah, J.T. Welch, Y.D. Vankar, M. Nojima, I. Kerekes, and J.A. Olah, *J. Org. Chem.*, 1979, **44**, 3872.
48. C.G. Bergstrom, R.T. Nicholson, and R.M. Dodson, *J. Org. Chem.*, 1963, **28**, 2633.
49. G.A. Olah and S.J. Kuhn, *J. Am. Chem. Soc.*, 1960, **82**, 2380.
50. D. Boenigk and D. Mootz, *J. Am. Chem. Soc.*, 1988, **110**, 2135.
51. G.A. Olah, *Acc. Chem. Res.*, 1980, **13**, 330.
52. T. Fukuhara, S. Sasaki, N. Yoneda, and A. Suzuki, *Bull. Chem. Soc. Jpn.*, 1990, **63**, 2058.
53. T. Fukuhara, N. Yoneda, K. Takamara, and A. Suzuki, *J. Fluorine Chem.*, 1991, **51**, 299.
54. Y. Girault, S. Geribaldi, M. Rouillard, and M. Azzaro, *Tetrahedron*, 1987, **43**, 2485.
55. G. Haufe, S. Lacombe, A. Laurent, and C. Rousset, *Tetrahedron Lett.*, 1983, **24**, 5877.
56. G. Alvernhe, A. Laurent, K. Touhami, R. Bartnik, and G. Mloston, *J. Fluorine Chem.*, 1985, **29**, 363.
57. S. Kanemoto, M. Shimizu, and H. Yoshioka, *Tetrahedron Lett.*, 1987, **28**, 663.
58. Y. Girault, M. Rouillard, M. Decouzan, and S. Geribaldi, *J. Fluorine Chem.*, 1990, **49**, 231.

59. G.A Olah and X.Y. Li, *Synlett.*, 1990, 267.
60. A. Beamer and W. Clarke, *Ber.*, 1879, **12**, 1067.
61. J.F.T. Berliner and R.M. Mann, *J. Phys. Chem.*, 1928, **32**, 1142.
62. R. Franz, *J. Fluorine Chem.*, 1980, **15**, 423.
63. M. Hudlicky, *J. Fluorine Chem.*, 1985, **28**, 461.
64. K.O. Christe and W.W. Wilson, *J. Fluorine Chem.*, 1990, **46**, 339.
65. J.H. Clark, E.M. Goodman, D.K. Smith, S.J. Brown, and J.M. Miller, *J. Chem. Soc., Chem. Commun.*, 1986, 657.
66. A.E. Bayliff and R.D. Chambers, *J. Chem. Soc., Perkin Trans. I*, 1988, 201.
67. C. Carmona, G. Eaton, and M.C.R Symons, *J. Chem. Soc., Chem. Commun.*, 1987, 873.
68. L. Gouin, J. Cousseau, and J.A.S. Smith, *J. Chem. Soc., Faraday Trans. II*, 1977, **73**, 1878.
69. A.M. Buswell, R.L. Maycock, and W. H. Rodebush, *J. Chem. Phys.*, 1940, **8**, 362.
70. J. Emsley, *Chem. Soc. Rev.*, 1980, 91.
71. S.W. Peterson and H. A. Levy, *J. Chem. Phys.*, 1952, **20**, 704.
72. K.M. Harmon, S.L. Madeira, and R.W. Carling, *Inorg. Chem.*, 1974, **13**, 1260.
73. I. Gennick, K.M. Harmon, and M.M. Potvin, *Inorg. Chem.*, 1977, **16**, 2033.
74. L. Andrews, S.R. Davis, and G.L. Johnson, *J. Phys. Chem.*, 1986, **90**, 4273.
75. M. Bimmler, M. Von Janta-Lipinski, P. Langen, and H. Plaul, East German Patent DD 221, 736.
76. G. Alvernhe, A. Laurent, and G. Haufe, *Synthesis*, 1987, 562.
77. I. Chehidi, M.M. Chaabouni, and A. Baklouti, *Tetrahedron Lett.*, 1989, **30**, 3167.
78. G. Haufe, G. Alvernhe, and A. Laurent, *Tetrahedron Lett.*, 1986, **27**, 4449.
79. G. Alvernhe, D. Anker, A. Laurent, G. Haufe, and C. Beguin, *Tetrahedron*, 1988, **44**, 3551.
80. G. Haufe, G. Alvernhe, and A. Laurent, *J. Fluorine Chem.*, 1990, **46**, 83.
81. G. Alvernhe, A. Laurent, and G. Haufe, French Patent, 2 616 145 (1988); *C. Abstr.* 112: 20544q.
82. G. Haufe, G. Alvernhe, D. Anker, A. Laurent, and C. Saluzzo, *Tetrahedron Lett.*, 1988, **29**, 2311.
83. C. Saluzzo, G. Alvernhe, and D. Anker, *J. Fluorine Chem.*, 1990, **47**, 467.
84. C. Saluzzo, G. Alvernhe, D. Anker, and G. Haufe, *Tetrahedron Lett.*, 1990, **31**, 663.
85. C. Saluzzo, G. Alvernhe, D. Anker, and G. Haufe, *Tetrahedron Lett.*, 1990, **31**, 2127.
86. G. Alvernhe, C.M. Ennakoua, S.M. Lacombe, and A.J. Laurent, *J. Org. Chem.*, 1981, **46**, 4938.
87. D. Picq, D. Anker, C. Rousset, and A. Laurent, *Tetrahedron Lett.*, 1983, **24**, 5619.

88. D. Picq, I. Drivas, G. Carret, D. Anker, and M. Abou-Assali, *Tetrahedron*, 1985, **41**, 2681.
89. P. Nebois, D. Picq, and D. Anker, *J. Carbohydr. Chem.*, 1990, **9**, 359.
90. D. Picq and D. Anker, *Carbohydr. Res.*, 1987, **166**, 309.
91. D. Picq and D. Anker, *J. Carbohydr. Chem.*, 1985, **4**, 113.
92. G. Alvernhe, A. Laurent, and G. Haufe, *J. Fluorine Chem.*, 1986, **34**, 147.
93. M. Muehlbacher and C.D. Poulter, *J. Org. Chem.*, 1988, **53**, 1026.
94. S. Takano, M. Yanase, and K. Ogasawara, *Chemistry Lett.*, 1989, 1689.
95. H. Amri and M.M. El Gaieo, *J. Fluorine Chem.*, 1990, **46**, 75.
96. J. Umezawa and K. Furuhashi, Japanese Patent JP 02, 167, 240 (1990).
97. M.M. Chaabouni and A. Baklouti, *Bull. Chim. Soc. Fr.*, 1989, 549.
98. J.H. Meurs, D.W. Sopher, and W. Eilenberg, *Angew. Chem., Int. Ed. Eng.*, 1989, **28**, 927.
99. N. Yoneda, T. Abe, T. Fukuhara, and A. Suzuki, *Chem. Lett.*, 1983, **7**, 1135.
100. N. Yoneda, S. Nagata, T. Fukuhara, and A. Suzuki, *Chem. Lett.*, 1984, 1241.
101. D. Haas, E. Kemnitz, M. Roennebeck, U. Schmidt, R. Kaden, and C. Henke, Patent Ger. (East) DD 268 170 (1989). *C. Abstr.*, **111**: P216313u.
102. N. Yoneda, *Tetrahedron*, 1991, **47**, 5329.
103. W.G. Brown and N.J. Letang, *J. Am. Chem. Soc.*, 1941, **63**, 358.
104. R.W. Alder, P.S. Bowmann, W.R.S. Steele, and D.R. Winterman, *J. Chem. Soc., Chem. Commun.*, 1968, 723.
105. H. Quast, W. Risler, and G. Dollscher, *Synthesis*, 1972, 558.
106. R.W. Alder, M.R. Bryce, N.C. Goode, N. Miller, and J. Owen, *J. Chem Soc., Perkin Trans I*, 1981, 2840.
107. H. Einspahr, J.-B. Robert, R.E. Marsh, and J.D. Roberts, *Acta. Crystallogr., Sect. B*, **29**, 1611.
108. M.R. Truker and B.L. Vickery, *J. Chem. Soc., Dalton Trans.*, 1972, 395.
109. E. Haselbach, A. Henriksson, F. Jachimowicz, and J. Wirz, *Helv. Chim. Acta.*, 1972, **55**, 167.
110. L.J. Altman, D. Laungani, G. Gunnarsson, H. Winnerstrom, and S. Forsen, *J. Am. Chem. Soc.*, 1978, **100**, 8264.
111. S.S. Pachkovskii and A.A. Voityuk, *Z. Fiz. Khim.*, 1990, **64**, 1592.
112. H.A. Staab and T. Saupe, *Angew. Chem. Int. Ed. Eng.*, 1988, **27**, 865.
113. R.W. Alder, *Chem. Rev.*, 1989, **89**, 1215.
114. F. Hibbert, *Acc. Chem. Res.*, 1984, **17**, 115.
115. E. Grech, Z. Malarski, and L. Sobczyk, *Chem. Phys. Lett.*, 1986, **128**, 259.
116. E. Grech, Z. Malarski, and L. Sobczyk, *J. Mol. Struct.*, 1985, **129**, 35.
117. B. Brzezinski, E. Grech, Z. Malarski, and L. Sobczyk, *J. Chem. Soc. Faraday Trans.*, 1990, **86**, 1777.
118. R.W. Alder and J.E. Anderson, *J. Chem. Soc., Perkin Trans. II*, 1973, 2086.

119. M.J. Diem, D.F. Burow, and J.L. Fry, *J. Org. Chem.*, 1977, **42**, 1801.
120. A. Ishida and T. Mukaiyama, *Chem. Lett.*, 1976, 1127.
121. A. Bayliff, *Ph. D., Thesis*, Durham University, 1986.
122. R.D. Chambers, G. Taylor, and R.L. Powell, *J. Chem. Soc., Perkin Trans. I*, 1980, 426.
123. R.D. Chambers, S. Partington, and D.B. Speight, *J. Chem. Soc., Perkin Trans. I*, 1974, 2673.
124. R.D. Chambers, J.A. Jackson, and R.A. Storey, *J. Chem. Soc., (C)*, 1968, 2221.
125. C.J. Drayton, W.T. Flowers, and R.N. Hazeldine, *J. Chem. Soc., (C)*, 1971, 2750.
126. R.L. Dressler and J.A. Young, *J. Org. Chem.*, 1967, **32**, 2004.
127. A.G. Pittman, D.L. Sharp, and R.E. Lundin, *J. Polymer Sci., A11*, 1966, **4**, 2637.
128. J.S. Brimacombe, A.B. Foster, R. Hems, J.H. Westwood, and L.D. Hall, *Can. J. Chem.*, 1970, **48**, 3946.
129. W.A. Szarek, G.W. Hay, and B. Doboszewski, *J. Chem. Soc., Chem. Commun.*, 1985, 663.
130. J.S. Brimacombe, P.A. Gent, and M. Stacey, *J. Chem. Soc. (C)*, 1968, 567.
131. L.N. Yakhontov, *Russ. Chem. Rev.*, 1984, **53**, 757.
132. I.B. Simon and V.P. Vvedenskii, *Zh. Obs. Khim.*, 1964, **34**, 4037.
133. S.J. Angyal and W.K. Warburton, *J. Chem. Soc.*, 1951, 2492.
134. P. Bosch, F. Camps, E. Chamarro, V. Gasol, and A. Guerrero, *Tetrahedron Lett.*, 1987, **28**, 4733.
135. J. March, *"Advanced Organic Chemistry"*, John Wiley and Sons, New York, 1985.
136. V. Kurasov, *Zh. Org. Khim.*, 1983, **19**, 590.
137. M.L. Hallensleben and H. Kansy, *Makromol. Chem.*, 1989, **190**, 231.
138. N.V. Vistorobskii and A.F. Pozharskii, *Z. Org. Chim.*, 1989, **25**, 2154.
139. G. Opitz, K. Rieth, and G. Rimmler, *Tetrahedron Lett.*, 1986, **27**, 167.
140. F. Terrier, J-C. Halle, M-J. Pouet, and M-P. Simonnin, *J. Org. Chem.*, 1986, **51**, 409.
141. M-P. Simonnin, J-C. Halle, M-J. Pouet, F. Terrier, and D. Davoust, *Can. J. Chem.*, 1986, **64**, 404.
142. R.D. Chambers, R.S. Matthews, G. Taylor, and R.L. Powell, *J. Chem. Soc., Perkin Trans. I*, 1980, 435.
143. R.D. Chambers and M.R. Bryce in *"Comprehensive Carbanion Chemistry"*, eds. E. Buncl and T. Durst, Elsevier, Amsterdam, 1987, ch. 5, p. 293.
144. M.R. Bryce, R.D. Chambers, and G. Taylor, *J. Chem. Soc., Perkin Trans. I*, 1984, 509.
145. M.W. Briscoe, R.D. Chambers, S.J. Mullins, T. Nakamura, and F.G. Drakesmith, *J. Chem. Soc., Chem. Commun.*, 1990, 1127.
146. T. Nakamura, these laboratories.
147. D.J. Cowley and I. Pasha, *J. Chem. Soc., Perkin Trans. 2.*, 1981, 918.
148. R.D. Chambers, *Adv. Heterocycl. Chem.*, 1981, **28**, 1.

149. J.M.E. Quirke, in *"Comprehensive Heterocyclic Chemistry"*, Eds. A.J. Boulton and A. McKillop, Pergamon, Oxford, 1984.
150. J.K. Chakrabarti, A.F. Cockerill, G.L.O. Davies, T.M. Hotten, D.M. Rackham, and D.E. Tupper, *J. Chem. Soc., Perkin Trans II*, 1974, 861.
151. R.A. Shaw and P. Ward, *J. Chem. Soc.*, 1965, 5446.
152. R.A. Shaw and P. Ward, *J. Chem. Soc. (B)*, 1968, 1431.
153. R.A. Shaw and P. Ward, Brit. Pat., 1, 119, 824 (1968). *C. Abstr.*, **69**: P87024.
154. R.A. Shaw and P. Ward, *J. Chem. Soc., (B)*, 1967, 123.
155. R.A. Shaw and P. Ward, *J. Chem. Soc. (B)*, 1969, 596.
156. T.L. Gilchrist, *"Heterocyclic Chemistry"*, Pitman, London, 1985.
157. J.K. Chakrabarti, R.W. Goulding, and A. Todd, *J. Chem. Soc. Perkin Trans I*, 1973, 2499.
158. J.K. Chakrabarti and D.E. Tupper, *J. Het. Chem.*, 1974, **11**, 417.
159. J.K. Chakrabarti and D.E. Tupper, *J. Org. Chem.*, 1975, **40**, 958.
160. J.R. Sandler, *J. Org. Chem.*, 1970, **35**, 3967.
161. R.A. Shaw and P. Ward, *J. Chem. Soc., Perkin Trans II*, 1973, 2075.
162. J.K. Chakrabarti and A. Todd, *Eur. J. Chem.-Chim. Ther.*, 1974, **9**, 146.
163. J.K. Chakrabarti and A. Todd, U.S. Pat., 3, 830, 915 (1974). *C. Abstr.*, **81**: P164750t.
164. J.K. Chakrabarti and A. Todd, Ger. Pat., 2, 117, 100 (1971). *C. Abstr.*, **76**: P46220u.
165. J.K. Chakrabarti and A. Todd, Brit. Pat., 1, 353, 581 (1970). *C. Abstr.*, **81**: P63688y.
166. J.K. Chakrabarti and D.E. Tupper, *Tetrahedron*, 1975, **31**, 1879.
167. D.J. Chadwick, in *"Comprehensive Heterocyclic Chemistry"*, Eds. C.W. Bird and G.W.H. Cheeseman, Pergamon, Oxford, 1984.
168. H-O. Kalinowski, S. Berger, and S. Braun, *"¹³C NMR Spectroscopy"*, John Wiley and Sons, New York, 1988.
169. D.J. Williams, *Angew. Chem. Int. Ed. Engl.*, 1984, **23**, 690.
170. W. Rettig, *Angew. Chem., Int. Ed. Engl.*, 1986, **25**, 971.
171. L.S. Pu, *J. Chem. Soc., Chem. Commun.*, 1991, 429.
172. R.D. Chambers, P.D. Philpot, and P.L. Russell, *J. Chem. Soc., Perkin Trans. I*, 1977, 1605.
173. C.H. Dungan and J.R. Van Wazer, *"Compilation of Reported ¹⁹F NMR Chemical Shifts"*, Wiley-Interscience, New York, 1970.

